



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Comparative efficacy of treatment strategies for hepatocellular carcinoma: systematic review and network meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-021269
Article Type:	Research
Date Submitted by the Author:	01-Jan-2018
Complete List of Authors:	<p>Tian, Guo; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China.</p> <p>Yang, Shigui; The State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, School of Medicine, Zhejiang University, The Key Laboratory of Infectious Diseases</p> <p>Yuan, Jinqiu; The Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong, Division of Epidemiology</p> <p>Threapleton, Diane; Chinese University of Hong Kong, School of Public Health & Primary Care</p> <p>Zhao, Qiyu ; Department of Ultrasound Medicine, the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China, Department of Ultrasound Medicine</p> <p>Chen, Fen ; Department of Hepatobiliary Pancreatic Surgery, First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China., Department of Hepatobiliary Pancreatic Surgery</p> <p>Cao, Hongcui ; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University</p> <p>Jiang, Tian'an; First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China., Department of Ultrasonography</p> <p>Li, Lanjuan; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University,</p>
Keywords:	resection, radiofrequency ablation, microwave ablation, transcatheter arterial chemoembolization, percutaneous ethanol injection, hepatocellular carcinoma

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

**Comparative efficacy of treatment strategies for hepatocellular carcinoma:
systematic review and network meta-analysis**

Guo Tian MD ^{1*}, Shigui Yang PhD ^{1*}, Jinqiu Yuan PhD ^{2,3}, Diane Threapleton PhD ²,
Qiyu Zhao PhD ⁴, Fen Chen PhD ⁵, Hongcui Cao PhD ¹, Tian'an Jiang PhD ⁴, Lanjuan
Li PhD ¹

¹State Key Laboratory for Diagnosis and Treatment of Infectious Diseases,
Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases,
The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou
310003, China.

²Division of Epidemiology, The Jockey Club School of Public Health and Primary
Care, The Chinese University of Hong Kong, Hong Kong;

³Shenzhen Municipal Key Laboratory for Health Risk Analysis, Shenzhen Research
Institute of The Chinese University of Hong Kong, Shenzhen, Guangdong Province,
China

⁴Department of Ultrasonography, First Affiliated Hospital, Zhejiang University
School of Medicine, Hangzhou 310003, China.

⁵Department of Hepatobiliary Pancreatic Surgery, First Affiliated Hospital, Zhejiang
University School of Medicine, Hangzhou 310003, China.

Correspondence to: Lanjuan Li, State Key Laboratory for Diagnosis and Treatment of
Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of

Infectious Diseases, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China (ljli@zju.edu.cn); Or Tian'an Jiang, Department of Ultrasound, First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou 310003, China (tiananjiang@zju.edu.cn); Or Hongcui Cao, State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China (hccao@zju.edu.cn).

*These authors contributed equally to this work.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Electronic word count: 2803 (excluding Tables and References)

Number of figures and tables: 7 figures and 15 tables

List of abbreviations in order of appearance: HCC: hepatocellular carcinoma; RES: resection; RFA: radiofrequency ablation; MWA: microwave ablation; TACE: transcatheter arterial chemoembolization; PEI: percutaneous ethanol injection; GRADE: Grading of Recommendations Assessment, Development and Evaluation; OR: odds ratio; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses; TR: TACE plus RFA; OS: overall survival; MCMC: Markov Chain Monte Carlo; CrI: credible interval; SUCRA: surface under the cumulative ranking curve LPS: lipopolysaccharide; TNF α : tumor necrosis factor α ; IL: interleukin; TGF β : transforming growth factor β .

Conflict of interest: The authors have declared that no competing interests regarding the publication of this paper.

Data sharing statement: No additional data are available.

Financial support: This study was supported by the opening foundation of the State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital of Medical College, Zhejiang University, grant NO. 2015KF06; This study was supported by the Foundation of Zhejiang Health Committee (2017KY346).

Author Contributions:

1. Conceived and designed the experiments: Hongcui Cao, Tian'an Jiang, Lanjuan Li

2. Performed the experiments: Guo Tian, Shigui Yang, Jinqiu Yuan, Diane Threapleton, Qiyu Zhao, Fen Chen, Tian'an Jiang
3. Analyzed the data: Guo Tian, Shigui Yang, Jinqiu Yuan, Qiyu Zhao
4. Contributed reagents/materials/analysis tools: Qiyu Zhao, Fen Chen
5. Wrote the manuscript: Guo Tian, Shigui Yang, Tian'an Jiang
6. Critically revised and approved the final version of manuscript: Diane Threapleton, Hongcui Cao, Tian'an Jiang, Lanjuan Li
7. Study supervision: Hongcui Cao, Tian'an Jiang, Lanjuan Li

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objective: Hepatocellular carcinoma (HCC) is the 3rd leading cause of cancer death worldwide. We conducted network meta-regression within a bayesian framework to compare and rank different treatment strategies for HCC through direct and indirect evidence from international studies.

Methods and analyses: We pooled the odds ratio (OR) for 1-, 3- and 5-year overall survival, based on lesions of size < 3 cm, 3-5 cm and ≤ 5 cm, using five therapeutic options including resection (RES), radiofrequency ablation (RFA), microwave ablation (MWA), transcatheter arterial chemoembolization (TACE) plus RFA (TR) and percutaneous ethanol injection (PEI).

Results: We identified 62 studies, including 23893 patients. After adjustment for study design, and in the full sample of studies, the treatments were ranked as follows for 5-year survival: 1) RES, 2) TR, 3) RFA, 4) MWA, and 5) PEI. The ranks were similar for 1 and 3-year survival, with RES and TR being the highest ranking treatments. In both smaller (<3cm) and larger tumors (3-5cm), RES and TR were also the two highest ranking treatments. There was little evidence of inconsistency between direct and indirect evidence.

Conclusion: The comparison of different treatment strategies for HCC indicated that RES is associated with longer survival. However, many of the between-treatment comparisons were not statistically significant and, for now, selection of strategies for treatment will depend patient and disease characteristics. Additionally, much of the evidence was provided by non randomised studies and knowledge gaps still exist.

More head-to-head comparisons between both RES and TR, or other approaches, will be necessary to confirm these findings.

Key words: resection; radiofrequency ablation; microwave ablation; transcatheter arterial chemoembolization; percutaneous ethanol injection; hepatocellular carcinoma.

Strengths and limitations of this study:

1. We conducted network meta-regression within a bayesian framework to compare and rank different treatment strategies for HCC through direct and indirect evidence from international studies.
2. We pooled the odds ratio (OR) for 1-, 3- and 5-year overall survival, based on lesions of size < 3 cm, 3-5 cm and ≤ 5 cm, using five therapeutic options including resection (RES), radiofrequency ablation (RFA), microwave ablation (MWA), transcatheter arterial chemoembolization (TACE) plus RFA (TR) and percutaneous ethanol injection (PEI).
3. The comparison of different treatment strategies for HCC indicated that RES is associated with longer survival.
4. A major limitation is in the inclusion of non-randomised studies, in which selection bias is likely to confound observations. Selection of treatment is likely to be based on individual or tumor characteristics, and thus these factors will bias and confound observations of survival.
5. All included studies did not report our primary outcome of interest (5-year survival) and this was a particular limitation among randomised studies.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Cancer was the second leading cause of death in 2013, behind cardiovascular disease, and in 2013 more than 8 million people died from cancer globally ¹⁻³. Hepatocellular carcinoma (HCC) is the 6th most common cancer worldwide and the 3rd leading cause of cancer death, with 5-year overall survival rates under 12% ^{4,5}.

Hepatic resection (RES) is the traditional choice for patients with HCC, without cirrhosis and with good remaining liver function ⁶. Despite nearly 70% 5-year survival, recurrence rates with surgery are high ⁷. Repeated hepatectomies to lengthen survival are not often appropriate owing to multiple-site tumor recurrence or patient background of liver cirrhosis ^{8,9}. Many locoregional therapies have been developed including ablative treatments such as percutaneous ethanol injection (PEI), radiofrequency ablation (RFA), or microwave ablation (MWA), and trans-arterial therapies such as transcatheter arterial chemoembolization (TACE) or transarterial chemotherapy infusion (TACI). Locoregional therapies are minimally invasive and therefore are cheaper and faster to recover from, as compared to resection. Such approaches may be appropriate for patients with unresectable, small or multiple carcinomas or those with severe cirrhosis. However, there may be a greater risk of recurrence because of incomplete destruction of cancer cells at the treatment margin, as seen with RFA ¹⁰.

Selection of treatment strategy is determined by liver function, tumour stage and patient performance status ⁷, but much uncertainty still remains surrounding the comparative efficacy of different treatment approaches. A recent review of

international guidelines for HCC found similarities but also some discrepancy in treatment allocation recommendations because of regional classification differences, secondary to a lack of solid or high-level evidence ¹¹. A recent review of therapies also revealed that there was no consensus on whether surgery or ablation was better for small tumors ⁷. Some discrepancy in prevalence and treatment outcomes may remain in different regions because of local biology, available resources or expertise and access to care ¹¹. However, if we ever hope to achieve standardized and evidence-based therapy for HCC, the unanswered question surrounding relative treatment efficacy of RES compared to ablative locoregional therapies must be resolved.

Traditional meta-analysis is limited by existing head-to-head treatment comparisons within included studies. It is therefore not possible to gauge the relative benefit of two treatments that have never been directly compared in studies. Real-life treatment-decisions are hindered by gaps in existing evidence, but network meta-analysis enables integration of direct and indirect comparisons to provide estimates for relative comparisons across many treatments ¹². In order to investigate comparative effectiveness among RES and common locoregional ablative therapies, we performed a systematic review and network meta-analysis.

Search Strategy

We conducted a systematic review and report findings in accordance with PRISMA for Network Meta-Analyses (PRISMA-NMA) ¹³ (PRISMA NMA Checklist).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The following databases were searched: PubMed, Embase, Web of science and Scopus, up to December 2015, using these keywords: resection, surgery, hepatectomy, radiofrequency ablation, transarterial chemoembolization, microwave thermal ablation, ethanol injection, liver, cancer, tumor (Additional file 1: Text S1). No language restrictions were used. Bibliographies from other relevant review articles were cross-examined for potential missed studies. Disagreement was resolved by a third reviewer. Citations were downloaded into reference management software and duplicate citations were electronically or manually removed.

We systematically included the studies using the following criteria: 1) original data from prospective or retrospective cohort studies and randomized clinical trials (RCTs) in humans; 2) reporting at least two treatments, including resection or any local ablative therapy (RES, RFA, MWA, PEI, or TACE+RFA (TR)); 3) mean lesion size ≤ 5 cm; 4) evaluating overall survival rate not less than one year after first or recurrent treatments. Conference abstracts and case reports were excluded, as were older publications from studies with multiple publications. Studies were excluded where participants had received combinations of the 5 included treatment approaches, such that outcomes could not be ascribed to individual therapies.

Data Extraction and Study Quality

Two investigators independently extracted and cross-checked the data from the eligible studies: author, year, study design, country, disease type, inclusion criteria, treatment style, study size, gender, age, tumor size, follow-up duration, treatment

complications and survival outcomes. If in disagreement, a third reviewer adjudicated. The level of evidence was appraised using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) guidance¹⁴, which was classified into four levels of high, moderate, low, and very low. The quality score was downgraded according to 5 domains, including risk of bias, inconsistency, indirectness, imprecision, and publication bias while scores were upgraded according to large effect, appropriate control for plausible confounding, and dose-response gradient.

Data Analysis

Network meta-analysis was used if a closed evidence loop was available. When possible, pair-wise direct head-to-head comparisons were conducted to calculate the pooled odds ratio (OR) and its 95% confidence interval (CI). Between-study heterogeneity was evaluated using the tau-squared statistic (τ^2)¹⁵. A node-splitting analysis was applied to check the consistency between direct evidence (existing real reported comparisons) and indirect evidence (estimated treatment comparisons) for their agreement on a specific node¹⁶. Bayesian network meta-analysis with Markov Chain Monte Carlo (MCMC), through a consistency model, was utilized to estimate the pooled ORs and its 95% credible interval (CrI) for the direct and indirect comparisons. The inconsistency model was used to check for heterogeneity due to chance imbalance in the distribution of effect modifiers. Consistency in every closed loop was checked by the loop-specific approach in order to estimate whether

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

treatment survival effects were disturbed by variance in the distribution of potential confounding factors among the studies. In order to compare and rank survival rates of different treatments we examined all studies first and then separately assessed smaller (<3cm) and larger (3-5cm) tumors. Random-effect meta-regression models were used, with and without adjustment for study design (cohort or RCT) and subgroup analyses were also conducted for RCTs in order to examine treatment effectiveness. We appraised the ranking probabilities for all therapies for each intervention and the treatment hierarchy was ordered by the surface under the cumulative ranking curve (SUCRA) ¹⁷. Sensitivity analysis was conducted to remove each study, in turn, and estimate the treatment effect in the remaining studies. Funnel plots were utilized to check the possible presence of publication bias or small-study bias ¹⁸. In this study, we used Bayesian MCMC simulations by WinBUGS 1.4 and graphically presented the results using Stata 13.

Results

Study Characteristics

After screening, 62 relevant studies in 61 articles were identified, of which 18 were randomized controlled trials and 44 were cohort studies ¹⁹⁻⁸⁰. We excluded 61571 duplicate or non-relevant citations (Figure 1). The summary characteristics of these studies are shown in Additional file 1: Table S1. Overall, 23893 patients of mean age from 46 to 73.5 years, with approximately 29236 tumors, were assigned to receive RES, RFA, MWA, TR and PEI, and the mean follow-up ranged from 1.5 to

5.7 years.

Network Meta-Analysis Results

Ten possible treatment comparisons among the five interventions were examined in the included studies. Comparable survival estimates were made for each treatment (per 1000 patients) and the survival OR among each of the treatment comparisons, according to follow-up duration, are presented in Additional file 1: Table S2, along with estimation of the quality of evidence using GRADE criteria.

Across the range of treatment comparisons and follow-up durations, evidence was graded between low and high quality. Evidence was often graded as low quality owing to publication bias and graded as high quality owing to a larger number of participants in direct comparisons.

Survival probabilities (estimated using Meanrank) and ranks for the five treatments in patients with tumours <3cm, 3-5cm or ≤ 5 cm (with and without adjustment for study design) are graphically displayed in Figures 2-5, and numerical details are given in Additional file 1: Table S3-S4. RES was consistently associated with greater survival (rank 1) compared to MWA, RFA, TR and PEI for the 5-year survival estimates. The ranks were similar for 1 and 3-year survival with RES or TR being ranked as 1 or 2 in most analyses. After adjustment for study design, and in the full sample of available studies (n=40), the treatments were ranked as follows for 5-year survival: 1) RES, 2) TR, 3) RFA, 4) MWA, and 5) PEI (Table S4).

Efficacy comparisons from network meta-regression for all treatments are

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

summarized in Table 1 and 2, according to follow-up duration and initial tumor size. Compared to RES, 5-year survival in all studies (trials and observational studies) for all tumours ≤ 5 cm, estimated from network comparisons, was 0.47 (95%CrI 0.22 to 0.87) for PEI, 0.79 (95%CrI 0.24 to 1.92) for TR, 0.56 (95%CrI 0.23 to 1.14) for MWA and 0.56 (95%CrI 0.27 to 0.99) for RFA (Table 2). When examining the comparisons across all treatments, the only significant difference for tumours < 3 cm was for 5-year survival, and a significantly worse survival was observed for PEI compared to RES 0.46 (95%CrI 0.18 to 0.95). For tumours between 3 and 5 cm, no significant differences were observed at 5-year survival, but significantly worse 3-year survival was observed with PEI, MWA and RFA compared to RES (Table 2). Despite smaller number of studies in analyses of only RCTs, the pairwise comparisons showed similar results. However, all relative rankings should be interpreted with caution because most network meta-regression comparisons did not suggest a statistically significant difference between treatments. Detailed results of each comparison for survival rates are shown in Additional file 1: Table S5-S10.

Loop-specific methods detected no inconsistency between the pairwise and network meta-analysis for most closed loops in the network (Additional file 1: Figure S1). However, inconsistency was observed between direct and indirect comparisons for the following loops: lesions < 3 cm: RES-RFA-TR, PEI-RES-RFA, MWA-RES-RFA; lesions 3-5cm: MWA-RES-RFA, RES-RFA-TR; and lesions ≤ 5 cm: RES-RFA-TR). In addition, tests for inconsistency were carried out (Additional file 1: Table S11-S13), which indicated a close relationship of between-trial heterogeneity

and inconsistency between “direct” and “indirect” evidence.

Sensitivity Analysis and publication bias

No significant change was observed when any one study was deleted. Funnel plots indicated that the included studies in each group were distributed symmetrically around the vertical line ($x=0$), suggesting that no obvious evidence of publication bias or small-sample effect existed in this network (Additional file 1: Figure S2).

Discussion

There are many techniques for attaining a large ablated zone and complete necrosis of HCC and this comprehensive review addresses two of the more common treatments, namely resection and ablation. In this network meta-analysis, of the five examined therapies, the pooled data showed RES ranked best in full sample analysis with or without adjustment for study design. In both smaller ($<3\text{cm}$) and larger tumors (3-5cm) RES remained the highest ranking treatment. However, most of the individual treatment comparisons were not statistically significant and thus, RES may not be superior to all other therapies. Our evidence indicates locoregional therapies and particularly RES or TR (TACE+RFA) are associated with longer survival.

Our observation of better survival outcomes with TR may be through the advantage of dual mechanisms. With TR, TACE induces hypoxic injury on cancer cells through occlusion of blood vessels and is followed by local ablation. This combination therapy may result in a larger ablated zone⁸¹, reduce the possibility of micrometastasis and recurrence, and thus, result in better survival outcomes than RFA

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

alone.

While being more invasive, and despite risk of complications, RES was associated with better survival outcomes after 1 year, 3 years and 5 years. This may be due to removal of larger sections of liver than can be targeted with locoregional therapies, thus removing a larger area of potentially cancerous cells. Additionally, rat models indicate that the liver has the potential to quickly restore its original size after partial hepatectomy. This may be mediated via interactions of lipopolysaccharide (LPS), tumor necrosis factor (TNF) α , interleukin (IL)-6, and transforming growth factor β (TGF β)⁸². However, evidence from rat models and human studies indicates that resection success is associated with resection size and regeneration is stunted with larger resections⁸³⁻⁸⁵. The safe limit for remnant liver volume in normal liver is approximately 30% of total liver volume, but this is estimated to rise to 40-50% in those with liver disease^{83 86}. Liver resection is recognised as the most efficient treatment for HCC but is only applicable for less than 30% of all patients (Morise 2014). However, developments in preoperative imaging techniques, laproscopic surgery and newly developing combinations with chemotherapy may extend its application to more advanced tumors⁸⁶. Furthermore, the consistent associations observed with all studies and only in RCTs indicates that patient selection bias in the observational studies does not wholly explain the better survival outcomes with RES.

Overall, we found PEI was associated with shorter survival than the other four therapies, a finding which is supported in previous studies^{20 29}. One study reported RFA was superior to PEI in achieving short- and long-term survival outcomes,

1
2
3 although PEI and RFA showed similar 5-year survival in lesions <3 cm⁵¹. The
4
5 possible reason why PEI is less effective than RFA may be because lesions often have
6
7 a thick capsule and therefore ethanol may not distribute through tissues.
8
9

10
11 There are several limitations in this study. Firstly, a major limitation is in the
12
13 inclusion of non-randomised studies, in which selection bias is likely to confound
14
15 observations. Selection of treatment is likely to be based on individual or tumor
16
17 characteristics, and thus these factors will bias and confound observations of survival.
18
19 Secondly, this study included both RCTs and observational studies, in which study
20
21 designs and type of data collection may not be comparable. However, findings were
22
23 consistent among both study designs. Thirdly, all included studies did not report our
24
25 primary outcome of interest (5-year survival) and this was a particular limitation
26
27 among randomised studies. Fourthly, for many individual comparisons, there were
28
29 either no direct comparisons or comparisons from only a small number of studies. The
30
31 lack of evidence may increase the risk of bias, which could enlarge or undervalue
32
33 effect size, and may explain the small inconsistency seen between direct and
34
35 estimated comparisons. Thus, we should be cautious in interpreting treatment
36
37 rankings for the different survival times and for different size lesions. While adverse
38
39 events from treatments may differ (not evaluated in detail in this review), by
40
41 examining overall survival outcomes in our review, we have taken account of both
42
43 long-term potential benefits and harms from treatments. The focus of these findings
44
45 should therefore be on the overall observation that RES or TR may be superior in
46
47 terms of survival, rather than focusing on specific OR values for individual treatment
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

comparisons.

In conclusion, the findings of the current bayesian network meta-analysis indicate that RES or TR may be among the most effective therapeutic approaches for HCC for 5-year survival in both smaller (< 3cm) and larger (3-5cm) lesions. However, evidence was of variable quality, and the majority of evidence came from non randomised studies, which are prone to selection bias and knowledge gaps still exist. For not, at the individual level, selection of strategies should depend on patient and clinical characteristics. To facilitate generation of evidence-based recommendations for HCC therpy, and to standardize treatment approaches, further head-to-head comparisons, especially of resection and ablative therapies, are required from high-quality RCTs, with long follow-up for survival outcomes.

Conflict of interests

The authors have declared that no competing interests regarding the publication of this paper.

Data sharing statement

No additional data are available.

Acknowledgments

This study was supported by the opening foundation of the State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center

for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital of Medical College, Zhejiang University, grant NO. 2015KF06; This study was supported by the Foundation of Zhejiang Health Committee (2017KY346).

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Collaborators GMaCoD. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;385:117-71
2. Lozano R, Naghavi M, Foreman K, *et al.* Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095-128
3. Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: Global Burden of Disease Study. *Lancet* 1997;349:1269-76
4. El-Serag HB. Hepatocellular carcinoma. *The New England journal of medicine* 2011;365:1118-27
5. Ferlay J, Shin HR, Bray F, *et al.* Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *International journal of cancer* 2010;127:2893-917
6. Forner A, Llovet JM, Bruix J. Hepatocellular carcinoma. *Lancet* 2012;379:1245-55
7. Raza A, Sood GK. Hepatocellular carcinoma review: current treatment, and evidence-based medicine. *World journal of gastroenterology* 2014;20:4115-27
8. Kishi Y, Hasegawa K, Sugawara Y, *et al.* Hepatocellular carcinoma: current management and future development-improved outcomes with surgical resection. *International journal of hepatology* 2011;2011:728103
9. Mazzaferro V, Bhoori S, Sposito C, *et al.* Milan criteria in liver transplantation

- for hepatocellular carcinoma: an evidence-based analysis of 15 years of experience. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society* 2011;17 Suppl 2:S44-57
10. Shan, Xue-Mei, Ding, *et al.* Radiofrequency ablation of hepatocellular carcinoma sized > 3 and ≤ 5 cm: Is ablative margin of more than 1 cm justified? *World journal of gastroenterology* 2011;19:7389-98
11. Yu SJ. A concise review of updated guidelines regarding the management of hepatocellular carcinoma around the world: 2010-2016. *Clinical & Molecular Hepatology* 2016;22:7-17
12. Chaimani A, Higgins JP, Mavridis D, *et al.* Graphical tools for network meta-analysis in STATA. *PloS one* 2013;8:e76654
13. Hutton B, Salanti G, Caldwell DM, *et al.* The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Annals of internal medicine* 2015;162:777-84
14. Puhan MA, Schunemann HJ, Murad MH, *et al.* A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis. *Bmj* 2014;349:g5630
15. Higgins JP, Green S. Cochrane Handbook for Systematic Reviews of Interventions. *Naunyn-Schmiedebergs Archiv für experimentelle Pathologie und Pharmakologie* 2009;2011:S38

16. Dias S, Welton NJ, Caldwell DM, *et al.* Checking consistency in mixed treatment comparison meta-analysis. *Statistics in medicine* 2010;29:932-44

17. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology* 2011;64:163-71

18. Jin ZC, Zhou XH, He J. Statistical methods for dealing with publication bias in meta-analysis. *Statistics in medicine* 2015;34:343-60

19. Zhang Z, Wu M, Chen H, *et al.* [Percutaneous radiofrequency ablation combined with transcatheter arterial chemoembolization for hepatocellular carcinoma]. *Zhonghua wai ke za zhi [Chinese journal of surgery]* 2002;40:826-9

20. Lencioni RA, Allgaier HP, Cioni D, *et al.* Small hepatocellular carcinoma in cirrhosis: randomized comparison of radio-frequency thermal ablation versus percutaneous ethanol injection. *Radiology* 2003;228:235-40

21. Lin SM, Lin CJ, Lin CC, *et al.* Radiofrequency ablation improves prognosis compared with ethanol injection for hepatocellular carcinoma < or =4 cm. *Gastroenterology* 2004;127:1714-23

22. Vivarelli M, Guglielmi A, Ruzzenente A, *et al.* Surgical resection versus percutaneous radiofrequency ablation in the treatment of hepatocellular carcinoma on cirrhotic liver. *Annals of surgery* 2004;240:102-7

23. Cho CM, Tak WY, Kweon YO, *et al.* [The comparative results of radiofrequency ablation versus surgical resection for the treatment of

- hepatocellular carcinoma]. *The Korean journal of hepatology* 2005;11:59-71
24. Hong SN, Lee SY, Choi MS, *et al.* Comparing the outcomes of radiofrequency ablation and surgery in patients with a single small hepatocellular carcinoma and well-preserved hepatic function. *Journal of clinical gastroenterology* 2005;39:247-52
25. Huang GT, Lee PH, Tsang YM, *et al.* Percutaneous ethanol injection versus surgical resection for the treatment of small hepatocellular carcinoma: a prospective study. *Annals of surgery* 2005;242:36-42
26. Lin SM, Lin CJ, Lin CC, *et al.* Randomised controlled trial comparing percutaneous radiofrequency thermal ablation, percutaneous ethanol injection, and percutaneous acetic acid injection to treat hepatocellular carcinoma of 3 cm or less. *Gut* 2005;54:1151-6
27. Lu MD, Xu HX, Xie XY, *et al.* Percutaneous microwave and radiofrequency ablation for hepatocellular carcinoma: a retrospective comparative study. *Journal of gastroenterology* 2005;40:1054-60
28. Montorsi M, Santambrogio R, Bianchi P, *et al.* Survival and recurrences after hepatic resection or radiofrequency for hepatocellular carcinoma in cirrhotic patients: a multivariate analysis. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2005;9:62-7; discussion 67-8
29. Shiina S, Teratani T, Obi S, *et al.* A randomized controlled trial of radiofrequency ablation with ethanol injection for small hepatocellular

carcinoma. *Gastroenterology* 2005;129:122-30

30. Chen MS, Li JQ, Zheng Y, *et al.* A prospective randomized trial comparing percutaneous local ablative therapy and partial hepatectomy for small hepatocellular carcinoma. *Annals of surgery* 2006;243:321-8

31. Lu MD, Kuang M, Liang LJ, *et al.* [Surgical resection versus percutaneous thermal ablation for early-stage hepatocellular carcinoma: a randomized clinical trial]. *Zhonghua yi xue za zhi* 2006;86:801-5

32. Cho YB, Lee KU, Suh KS, *et al.* Hepatic resection compared to percutaneous ethanol injection for small hepatocellular carcinoma using propensity score matching. *Journal of gastroenterology and hepatology* 2007;22:1643-9

33. Gao W, Chen MH, Yan K, *et al.* Therapeutic effect of radiofrequency ablation in unsuitable operative small hepatocellular carcinoma. *Chinese Journal of Medical Imaging Technology* 2007

34. Lupo L, Panzera P, Giannelli G, *et al.* Single hepatocellular carcinoma ranging from 3 to 5 cm: radiofrequency ablation or resection? *HPB : the official journal of the International Hepato Pancreato Biliary Association* 2007;9:429-34

35. Zhou T, Qiu YD, Kong WT. Comparing the effect of radiofrequency ablation and surgical resection for the treatment of small hepatocellular. *Journal of Hepatobiliary Surgery* 2007

36. Abu-Hilal M, Primrose JN, Casaril A, *et al.* Surgical resection versus radiofrequency ablation in the treatment of small unifocal hepatocellular

- carcinoma. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2008;12:1521-6
37. Brunello F, Veltri A, Carucci P, *et al.* Radiofrequency ablation versus ethanol injection for early hepatocellular carcinoma: A randomized controlled trial. *Scandinavian journal of gastroenterology* 2008;43:727-35
38. Guglielmi A, Ruzzenente A, Valdegamberi A, *et al.* Radiofrequency ablation versus surgical resection for the treatment of hepatocellular carcinoma in cirrhosis. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2008;12:192-8
39. Hiraoka A, Horiike N, Yamashita Y, *et al.* Efficacy of radiofrequency ablation therapy compared to surgical resection in 164 patients in Japan with single hepatocellular carcinoma smaller than 3 cm, along with report of complications. *Hepato-gastroenterology* 2008;55:2171-4
40. Ohmoto K, Yoshioka N, Tomiyama Y, *et al.* Comparison of therapeutic effects between radiofrequency ablation and percutaneous microwave coagulation therapy for small hepatocellular carcinomas. *Journal of gastroenterology and hepatology* 2009;24:223-7
41. Sakaguchi H, Seki S, Tsuji K, *et al.* Endoscopic thermal ablation therapies for hepatocellular carcinoma: a multi-center study. *Hepatology research : the official journal of the Japan Society of Hepatology* 2009;39:47-52
42. Santambrogio R, Opocher E, Zuin M, *et al.* Surgical resection versus laparoscopic radiofrequency ablation in patients with hepatocellular carcinoma

and Child-Pugh class a liver cirrhosis. *Annals of surgical oncology* 2009;16:3289-98

43. Shibata T, Isoda H, Hirokawa Y, *et al.* Small hepatocellular carcinoma: is radiofrequency ablation combined with transcatheter arterial chemoembolization more effective than radiofrequency ablation alone for treatment? *Radiology* 2009;252:905-13

44. Ueno S, Sakoda M, Kubo F, *et al.* Surgical resection versus radiofrequency ablation for small hepatocellular carcinomas within the Milan criteria. *Journal of hepato-biliary-pancreatic surgery* 2009;16:359-66

45. Xiang-Yang BU, Wang Y, Zhong GE, *et al.* Comparison of radiofrequency ablation and surgical resection for small primary liver carcinoma. *Chinese Archives of General Surgery* 2009

46. Guo WX, Zhai B, Lai EC, *et al.* Percutaneous radiofrequency ablation versus partial hepatectomy for multicentric small hepatocellular carcinomas: a nonrandomized comparative study. *World journal of surgery* 2010;34:2671-6

47. Huang J, Yan L, Cheng Z, *et al.* A randomized trial comparing radiofrequency ablation and surgical resection for HCC conforming to the Milan criteria. *Annals of surgery* 2010;252:903-12

48. Kagawa T, Koizumi J, Kojima S, *et al.* Transcatheter arterial chemoembolization plus radiofrequency ablation therapy for early stage hepatocellular carcinoma: comparison with surgical resection. *Cancer* 2010;116:3638-44

- 1
2
3
4 49. Morimoto M, Numata K, Kondou M, *et al.* Midterm outcomes in patients with
5
6 intermediate-sized hepatocellular carcinoma: a randomized controlled trial for
7
8 determining the efficacy of radiofrequency ablation combined with
9
10 transcatheter arterial chemoembolization. *Cancer* 2010;116:5452-60
11
12
13 50. Azab M, Zaki S, El-Shetey AG, *et al.* Radiofrequency ablation combined with
14
15 percutaneous ethanol injection in patients with hepatocellular carcinoma. *Arab*
16
17 *journal of gastroenterology : the official publication of the Pan-Arab*
18
19 *Association of Gastroenterology* 2011;12:113-8
20
21
22
23 51. Giorgio A, Di Sarno A, De Stefano G, *et al.* Percutaneous radiofrequency
24
25 ablation of hepatocellular carcinoma compared to percutaneous ethanol
26
27 injection in treatment of cirrhotic patients: an Italian randomized controlled
28
29 trial. *Anticancer research* 2011;31:2291-5
30
31
32
33 52. Hung HH, Chiou YY, Hsia CY, *et al.* Survival rates are comparable after
34
35 radiofrequency ablation or surgery in patients with small hepatocellular
36
37 carcinomas. *Clinical gastroenterology and hepatology : the official clinical*
38
39 *practice journal of the American Gastroenterological Association*
40
41 2011;9:79-86
42
43
44
45 53. Nishikawa H, Inuzuka T, Takeda H, *et al.* Comparison of percutaneous
46
47 radiofrequency thermal ablation and surgical resection for small hepatocellular
48
49 carcinoma. *BMC gastroenterology* 2011;11:143
50
51
52
53 54. Yun WK, Choi MS, Choi D, *et al.* Superior long-term outcomes after surgery
54
55 in child-pugh class a patients with single small hepatocellular carcinoma
56
57
58
59
60

compared to radiofrequency ablation. *Hepatology international* 2011;5:722-9

55. Zhang J, Liu HC, Zhou L. The effectiveness of radiofrequency ablation for the treatment of liver cancer. *Journal of Hepatobiliary Surgery* 2011;19:30-33

56. Zhao M, Wang JP, Li W, *et al.* [Comparison of safety and efficacy for transcatheter arterial chemoembolization alone and plus radiofrequency ablation in the treatment of single branch portal vein tumor thrombus of hepatocellular carcinoma and their prognosis factors]. *Zhonghua yi xue za zhi* 2011;91:1167-72

57. Feng K, Yan J, Li X, *et al.* A randomized controlled trial of radiofrequency ablation and surgical resection in the treatment of small hepatocellular carcinoma. *Journal of hepatology* 2012;57:794-802

58. Peng ZW, Lin XJ, Zhang YJ, *et al.* Radiofrequency ablation versus hepatic resection for the treatment of hepatocellular carcinomas 2 cm or smaller: a retrospective comparative study. *Radiology* 2012;262:1022-33

59. Peng ZW, Zhang YJ, Liang HH, *et al.* Recurrent hepatocellular carcinoma treated with sequential transcatheter arterial chemoembolization and RF ablation versus RF ablation alone: a prospective randomized trial. *Radiology* 2012;262:689-700

60. Signoriello S, Annunziata A, Lama N, *et al.* Survival after locoregional treatments for hepatocellular carcinoma: a cohort study in real-world patients. *TheScientificWorldJournal* 2012;2012:564706

61. Wang JH, Wang CC, Hung CH, *et al.* Survival comparison between surgical

- resection and radiofrequency ablation for patients in BCLC very early/early stage hepatocellular carcinoma. *Journal of hepatology* 2012;56:412-8
62. Desiderio J, Trastulli S, Pasquale R, *et al.* Could radiofrequency ablation replace liver resection for small hepatocellular carcinoma in patients with compensated cirrhosis? A 5-year follow-up. *Langenbeck's archives of surgery / Deutsche Gesellschaft fur Chirurgie* 2013;398:55-62
63. Ding J, Jing X, Liu J, *et al.* Comparison of two different thermal techniques for the treatment of hepatocellular carcinoma. *European journal of radiology* 2013;82:1379-84
64. Guo WX, Sun JX, Cheng YQ, *et al.* Percutaneous radiofrequency ablation versus partial hepatectomy for small centrally located hepatocellular carcinoma. *World journal of surgery* 2013;37:602-7
65. Hasegawa K, Kokudo N, Makuuchi M, *et al.* Comparison of resection and ablation for hepatocellular carcinoma: a cohort study based on a Japanese nationwide survey. *Journal of hepatology* 2013;58:724-9
66. Iida H, Aihara T, Ikuta S, *et al.* A comparative study of therapeutic effect between laparoscopic microwave coagulation and laparoscopic radiofrequency ablation. *Hepato-gastroenterology* 2013;60:662-5
67. Imai K, Beppu T, Chikamoto A, *et al.* Comparison between hepatic resection and radiofrequency ablation as first-line treatment for solitary small-sized hepatocellular carcinoma of 3 cm or less. *Hepatology research : the official journal of the Japan Society of Hepatology* 2013;43:853-64

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

68. Kim JW, Shin SS, Kim JK, *et al.* Radiofrequency ablation combined with transcatheter arterial chemoembolization for the treatment of single hepatocellular carcinoma of 2 to 5 cm in diameter: comparison with surgical resection. *Korean journal of radiology* 2013;14:626-35

69. Lai EC, Tang CN. Radiofrequency ablation versus hepatic resection for hepatocellular carcinoma within the Milan criteria--a comparative study. *International journal of surgery* 2013;11:77-80

70. Lin ZZ, Shau WY, Hsu C, *et al.* Radiofrequency ablation is superior to ethanol injection in early-stage hepatocellular carcinoma irrespective of tumor size. *PloS one* 2013;8:e80276

71. Peng ZW, Zhang YJ, Chen MS, *et al.* Radiofrequency ablation with or without transcatheter arterial chemoembolization in the treatment of hepatocellular carcinoma: a prospective randomized trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2013;31:426-32

72. Tohme S, Geller DA, Cardinal JS, *et al.* Radiofrequency ablation compared to resection in early-stage hepatocellular carcinoma. *HPB : the official journal of the International Hepato Pancreato Biliary Association* 2013;15:210-7

73. Wong KM, Yeh ML, Chuang SC, *et al.* Survival comparison between surgical resection and percutaneous radiofrequency ablation for patients in Barcelona Clinic Liver Cancer early stage hepatocellular carcinoma. *Indian journal of gastroenterology : official journal of the Indian Society of Gastroenterology* 2013;32:253-7

74. Zhang L, Wang N, Shen Q, *et al.* Therapeutic efficacy of percutaneous radiofrequency ablation versus microwave ablation for hepatocellular carcinoma. *PloS one* 2013;8:e76119
75. Abdelaziz A, Elbaz T, Shousha HI, *et al.* Efficacy and survival analysis of percutaneous radiofrequency versus microwave ablation for hepatocellular carcinoma: an Egyptian multidisciplinary clinic experience. *Surgical endoscopy* 2014;28:3429-34
76. Shi J, Sun Q, Wang Y, *et al.* Comparison of microwave ablation and surgical resection for treatment of hepatocellular carcinomas conforming to Milan criteria. *Journal of gastroenterology and hepatology* 2014;29:1500-7
77. Yang HJ, Lee JH, Lee DH, *et al.* Small single-nodule hepatocellular carcinoma: comparison of transarterial chemoembolization, radiofrequency ablation, and hepatic resection by using inverse probability weighting. *Radiology* 2014;271:909-18
78. Zhang T, Li K, Luo H, *et al.* [Long-term outcomes of percutaneous microwave ablation versus repeat hepatectomy for treatment of late recurrent small hepatocellular carcinoma: a retrospective study]. *Zhonghua yi xue za zhi* 2014;94:2570-2
79. Pompili M, De Matthaeis N, Saviano A, *et al.* Single hepatocellular carcinoma smaller than 2 cm: are ethanol injection and radiofrequency ablation equally effective? *Anticancer research* 2015;35:325-32
80. Xu J, Zhao Y. Comparison of percutaneous microwave ablation and

laparoscopic resection in the prognosis of liver cancer. *International journal of clinical and experimental pathology* 2015;8:11665-9

81. Goldberg SN, Giron GD, Lukyanov AN, *et al.* Percutaneous tumor ablation: increased necrosis with combined radio-frequency ablation and intravenous liposomal doxorubicin in a rat breast tumor model. *Radiology* 2002;222:797-804

82. Taub R. Liver regeneration: from myth to mechanism. *Nature reviews Molecular cell biology* 2004;5:836-47

83. Martins PN, Theruvath TP, Neuhaus P. Rodent models of partial hepatectomies. *Liver international : official journal of the International Association for the Study of the Liver* 2008;28:3-11

84. Kele PG, De BM, Ej VDJ, *et al.* Early hepatic regeneration index and completeness of regeneration at 6 months after partial hepatectomy. *British Journal of Surgery* 2012;99:1113–19

85. Preziosi ME, Monga SP. Update on the Mechanisms of Liver Regeneration. *Seminars in Liver Disease* 2017;37:141

86. Morise Z, Kawabe N, Tomishige H, *et al.* Recent advances in liver resection for hepatocellular carcinoma. *Frontiers in Surgery* 2014;1:21

File legends:

Figure 1 Flow chart of search.

Figure 2 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in RCTs.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments.

i Lesions < 3 cm.

ii Lesions 3-5 cm.

iii Lesions \leq 5 cm.

Figure 3 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in all studies.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments.

i Lesions < 3 cm.

ii Lesions 3-5 cm.

iii Lesions \leq 5 cm.

Figure 4 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in RCTs

A Lesions < 3 cm

1
2
3
4 B Lesions 3-5 cm

5
6 C Lesions \leq 5 cm (full sample).
7

8
9
10
11 **Figure 5 Treatment ranks for 1-year, 3-year and 5-year survival rates, according**
12
13 **to lesion size in all studies.**
14

15
16 A Lesions < 3 cm
17
18 B Lesions 3-5 cm
19
20
21 C Lesions \leq 5 cm (full sample).
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Odds ratios (95% credible interval) according to network meta-analyses for efficacy of treatments for all pairwise comparisons in randomized controlled trials.

<3cm for 1-year survival					
PEI					
10.75 (0.01-29.11)	TR				
0.08 (0-0.42)	1.42 (0-5.94)	MWA			
0.68 (0.28-1.36)	13.24 (0.02-55.15)	154.8 (1.74-590.10)	RFA		
0.68 (0.19-1.76)	15.61 (0.02-54.78)	161.8 (1.39-581.00)	1.01 (0.40-2.14)	RES	
<3cm for 3-year survival					
PEI					
1.29 (0.13-4.99)	TR				
NA	NA	MWA			
0.88 (0.44-1.79)	1.64 (0.20-5.84)	NA	RFA		
0.75 (0.28-1.89)	1.44 (0.14-5.50)	NA	0.86 (0.40-1.68)	RES	
<3cm for 5-year survival					
PEI					
NA	TR				
NA	NA	MWA			
0.93 (0.08-3.85)	NA	NA	RFA		
0.49 (0.04-2.02)	NA	NA	0.71 (0.10-2.47)	RES	
3-5cm for 1-year survival					
PEI					
NA	TR				
NA	NA	MWA			
NA	3.40 (0.64-11.93)	NA	RFA		
NA	1.00 (0-5.00)	NA	0.25 (0-1.47)	RES	
3-5cm for 3-year survival					
PEI					
NA	TR				
NA	NA	MWA			
NA	3.98 (0.71-15.22)	NA	RFA		
NA	1.14 (0-6.20)	NA	0.24 (0-1.25)	RES	
3-5cm for 5-year survival					
PEI					
NA	TR				
NA	NA	MWA			
NA	7.64 (0.14-42.49)	NA	RFA		
NA	12.87 (0.02-44.43)	NA	1.05 (0.03-5.33)	RES	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

≤5cm for 1-year survival

PEI					
0.26 (0.06-0.69)	TR				
0.24 (0.03-0.81)	1.26 (0.14-4.73)	MWA			
0.65 (0.32-1.14)	3.3 (1.05-8.21)	4.62 (0.85-15.59)	RFA		
0.42 (0.14-0.98)	2.15 (0.49-6.46)	2.75 (0.52-9.18)	0.65 (0.28-1.31)	RES	

≤5cm for 3-year survival

PEI					
0.49 (0.13-1.33)	TR				
1.25 (0.11-5.36)	3.25 (0.24-14.23)	MWA			
0.83 (0.39-1.73)	2.09 (0.81-4.65)	1.71 (0.17-6.61)	RFA		
0.66 (0.23-1.78)	1.69 (0.47-4.87)	1.18 (0.16-4.30)	0.80 (0.36-1.69)	RES	

≤5cm for 5-year survival

PEI					
1.51 (0.02-7.71)	TR				
NA	NA	MWA			
0.90 (0.08-3.65)	3.59 (0.14-18.06)	NA	RFA		
0.49 (0.04-2.03)	2.96 (0.05-14.70)	NA	0.72 (0.11-2.48)	RES	

The reference treatment (1.00) for all comparisons is listed to the right hand side
RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection;

Table 2 Odds ratios (95% credible interval) according to network meta-analyses for efficacy of treatments for all pairwise comparisons in all studies

3cm for 1-year survival					
PEI	TR	MWA	RFA	RES	
0.56 (0.07-2.13)					
0.55 (0.18-1.29)	1.96 (0.21-7.87)				
0.69 (0.39-1.13)	2.45 (0.33-8.72)	1.51 (0.60-3.11)			
0.71 (0.24-1.60)	2.51 (0.26-9.65)	1.55 (0.41-4.10)	1.03 (0.42-2.07)		
3cm for 3-year survival					
PEI	TR	MWA	RFA	RES	
0.59 (0.15-1.67)					
1.01 (0.45-2.00)	2.35 (0.54-6.80)				
0.95 (0.59-1.47)	2.21 (0.60-5.76)	1.02 (0.54-1.76)			
0.80 (0.33-1.68)	1.87 (0.40-5.56)	0.87 (0.31-1.96)	0.85 (0.40-1.62)		
3cm for 5-year survival					
PEI	TR	MWA	RFA	RES	
1.06 (0.19-3.41)					
0.90 (0.38-1.83)	1.37 (0.23-4.59)				
0.81 (0.48-1.28)	1.24 (0.25-3.80)	1.00 (0.50-1.77)			
0.46 (0.18-0.95)	0.72 (0.11-2.48)	0.58 (0.18-1.33)	0.58 (0.24-1.11)		
3-5cm for 1-year survival					
PEI	TR	MWA	RFA	RES	
0.21 (0.04-0.56)					
0.60 (0.09-1.94)	3.46 (0.57-11.35)				
0.50 (0.17-1.13)	2.92 (1.14-6.65)	1.25 (0.31-3.46)			
0.10 (0-0.63)	0.56 (0-3.31)	0.24 (0-1.61)	0.19 (0-1.18)		
3-5cm for 3-year survival					
PEI	TR	MWA	RFA	RES	
0.30 (0.03-1.06)					
0.90 (0.08-3.36)	3.48 (0.62-11.64)				
0.57 (0.10-1.83)	2.37 (0.90-5.53)	1.01 (0.25-2.72)			
0.09 (0-0.44)	0.36 (0.01-1.73)	0.15 (0-0.77)	0.14 (0.01-0.68)		
3-5cm for 5-year survival					
PEI	TR	MWA	RFA	RES	
6.11 (0-3.02)					
1.88 (0.04-5.54)	13.88 (0.19-50.64)				
0.79 (0.05-2.64)	7.08 (0.25-26.41)	1.25 (0.18-3.84)			
1.88 (0.01-3.18)	14.49 (0.05-27.29)	1.79 (0.03-5.39)	0.91 (0.05-4.18)		

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

≤5cm for 1-year survival

PEI					
0.30 (0.11-0.63)	TR				
0.91 (0.41-1.79)	3.51 (1.78-8.52)	MWA			
0.78 (0.51-1.13)	3.01 (1.33-6.15)	0.95 (0.48-1.67)	RFA		
0.61 (0.26-1.25)	2.35 (0.74-5.96)	0.73 (0.28-1.55)	0.78 (0.37-1.49)	RES	

≤5cm for 3-year survival

PEI					
0.52 (0.25-0.96)	TR				
1.03 (0.56-1.77)	2.16 (0.99-4.16)	MWA			
0.92 (0.63-1.32)	1.93 (1.05-3.29)	0.94 (0.58-1.44)	RFA		
0.71 (0.37-1.30)	1.50 (0.64-3.08)	0.72 (0.36-1.32)	0.78 (0.44-1.29)	RES	

≤5cm for 5-year survival

PEI					
0.71 (0.26-1.57)	TR				
0.90 (0.47-1.58)	1.50 (0.52-3.46)	MWA			
0.85 (0.57-1.22)	1.42 (0.58-2.96)	1.01 (0.60-1.59)	RFA		
0.47 (0.22-0.87)	0.79 (0.24-1.92)	0.56 (0.23-1.14)	0.56 (0.27-0.99)	RES	

The reference treatment (1.00) for all comparisons is listed to the right hand side

- RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection.




Figure 1 Flow chart of search.

254x190mm (300 x 300 DPI)




Figure 2 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in RCTs. Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments.

i Lesions < 3 cm.
ii Lesions 3-5 cm.
iii Lesions ≤ 5 cm.

227x223mm (300 x 300 DPI)




Figure 3 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in all studies.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments.

i Lesions < 3 cm.

ii Lesions 3-5 cm.

iii Lesions ≤ 5 cm.

88x79mm (300 x 300 DPI)




Figure 4 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in RCTs
A Lesions < 3 cm
B Lesions 3-5 cm
C Lesions ≤ 5 cm (full sample).

118x117mm (300 x 300 DPI)




Figure 5 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in all studies.
A Lesions < 3 cm
B Lesions 3-5 cm
C Lesions ≤ 5 cm (full sample).

118x117mm (300 x 300 DPI)

Text S1.

Search strategy:

Pubmed (1950-present)

1. ("TACE" OR "transarterial chemoembolization")
2. ("RFA" OR "radiofrequency ablation" OR "RF ablation" OR "radiofrequency thermal ablation" OR "RTA")
3. (PEI OR "ethanol injection" OR "ethanol ablation" OR "alcohol ablation")
4. ("microwave ablation" OR "microwave thermal ablation" OR MWA)
5. (liver OR hepato*)
6. (neoplas* OR cancer OR tumor OR tumour OR carcinoma OR oncolog*)
7. 1 OR 2 OR 3 OR 4
8. 5 AND 6 AND 7
9. "Ablation Techniques"[Mesh]
10. "Embolization"[Mesh]
11. "Liver Neoplasms"[Mesh]
12. 9 OR 10
13. 12 AND 11
14. 8 OR 13
15. (resection OR surgery OR hepatectomy)
16. (ablation OR injection OR embolization)
17. 5 AND 6 AND 15 AND 16
18. "Hepatectomy"[Mesh]
19. 12 AND 18 AND 11
20. 17 OR 19
21. 14 OR 20

Embase(1980-present)

1. 'TACE':ab,ti
2. 'transarterial chemoembolization':ab,ti
3. 1 OR 2
4. 'rfa':ab,ti

5. 'radiofrequency ablation':ab,ti
6. 'rf ablation':ab,ti
7. 'radiofrequency thermal ablation':ab,ti
8. 'rta':ab,ti
9. 4 OR 5 OR 6 OR 7 OR 8
10. 'PEI':ab,ti
11. ' ethanol injection ':ab,ti
12. ' ethanol ablation ':ab,ti
13. ' alcohol ablation ':ab,ti
14. 10 OR 11 OR 12 OR 13
15. ' microwave ablation ':ab,ti
16. ' microwave thermal ablation ':ab,ti
17. ' MWA ':ab,ti
18. 15 OR 16 OR 17
19. ' liver':ab,ti
20. ' hepato*':ab,ti
21. 19 OR 20
22. ' neoplas*':ab,ti
23. ' cancer ':ab,ti
24. ' tumor ':ab,ti
25. ' tumour ':ab,ti
26. ' carcinoma ':ab,ti
27. ' oncolog*':ab,ti
28. 22 OR 23 OR 24 OR 25 OR 26 OR 27
29. 3 OR 9 OR 14 OR 18
30. 21 AND 28 AND 29
31. ' resection':ab,ti
32. ' surgery':ab,ti
33. ' hepatectomy':ab,ti
34. 31 OR 32 OR 33
35. ' ablation':ab,ti

- 36. ' injection':ab,ti
- 37. ' embolization':ab,ti
- 38. 35 OR 36 OR 37
- 39. 34 AND 38 AND 21 AND 28
- 40. 30 OR 39

Scoups

- 1. TITLE-ABS-KEY (“TACE”)
- 2. TITLE-ABS-KEY (“transarterial chemoembolization”)
- 3. 1 OR 2
- 4. TITLE-ABS-KEY ("RFA")
- 5. TITLE-ABS-KEY (“radiofrequency ablation”)
- 6. TITLE-ABS-KEY ("RF ablation")
- 7. TITLE-ABS-KEY ("radiofrequency thermal ablation")
- 8. TITLE-ABS-KEY ("RTA")
- 9. 4 OR 5 OR 6 OR 7 OR8
- 10. TITLE-ABS-KEY (“PEI”)
- 11. TITLE-ABS-KEY (“ethanol injection”)
- 12. TITLE-ABS-KEY (“ethanol ablation”)
- 13. TITLE-ABS-KEY (“alcohol ablation”)
- 14. 10 OR 11 OR 12 OR 13
- 15. TITLE-ABS-KEY ("microwave ablation")
- 16. TITLE-ABS-KEY ("microwave thermal ablation ")
- 17. TITLE-ABS-KEY ("MWA")
- 18. 15 OR 16 OR 17
- 19. TITLE-ABS-KEY ("liver ")
- 20. TITLE-ABS-KEY ("hepato*")
- 21. 19 OR 20
- 22. TITLE-ABS-KEY ("neoplas*")
- 23. TITLE-ABS-KEY ("cancer")

24. TITLE-ABS-KEY ("tumor")
25. TITLE-ABS-KEY ("tumour")
26. TITLE-ABS-KEY ("carcinoma")
27. TITLE-ABS-KEY ("oncolog*")
28. 22 OR 23 OR 24 OR 25 OR 26 OR 27
29. 3 OR 9 OR 14 OR 18
30. 29 AND 21 AND 28
31. TITLE-ABS-KEY ("resection")
32. TITLE-ABS-KEY ("surgery")
33. TITLE-ABS-KEY ("hepatectomy")
34. 31 OR 32 OR 33
35. TITLE-ABS-KEY ("ablation")
36. TITLE-ABS-KEY ("injection")
37. TITLE-ABS-KEY ("embolization")
38. 35 OR 36 OR 37
39. 34 AND 38 AND 21 AND 28
40. 30 OR 39

Web of science

1. TS=(ablation)
2. TS=(embolization)
3. 1 OR 2
4. TS=(hepatectomy)
5. TS=(liver neoplasms)
6. 3 AND 4 AND 5
7. TI=(resection)
8. TI=(surgery)
9. TI=(hepatectomy)
10. 7 OR 8 OR 9
11. TI=(ablation)

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
12. TI=(injection)
13. TI=(embolization)
14. 11 OR 12 OR 13
15. TI=(liver)
16. TI=(hepato*)
17. 15 OR 16
18. TI=(neoplas*)
19. TI=(cancer)
20. TI=(tumor)
21. TI=(tumour)
22. TI=(carcinoma)
23. TI=(oncolog*)
24. 18 OR 19 OR 20 OR 21 OR 22 OR 23
25. 10 AND 14 AND 17 AND 24
26. 3 AND 5
27. TI=(TACE)
28. TI=("transarterial chemoembolization")
29. 27 OR 28
30. TI=(RFA)
31. TI=("radiofrequency ablation")
32. TI=("RF ablation")
33. TI=("radiofrequency thermal ablation")
34. TI=(RTA)
35. 30 OR 31 OR 32 OR 33 OR 34
36. TI=(PEI)
37. TI=("ethanol injection")
38. TI=("ethanol ablation")
39. TI=("alcohol ablation")
40. 36 OR 37 OR 38 OR 39
41. TI=("microwave ablation")
42. TI=("microwave thermal ablation")

43. TI=(MWA)
 44. 41 OR 42 OR 43
 45. 29 OR 35 OR 40 OR 44
 46. 46 AND 17 AND 24
 47. 6 OR 25 OR 26 OR 46

Table S1.
Summary of the studies included in the network meta-analysis.

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Zhang 2002 ¹⁹	Prospective cohort	China	HCC	0.3-2	RFA	15(15)	13/2	61.8 (38-78)	4.1 (2.4-6.0)	NA	0.80(1y)	0.80(1y)	NA
					TR	15(15)	12/3	57.8 (39-72)	4.6 (2.3-7.1)	NA	1.00(1y)	1.00 (1y)	NA
Lencioni 2003 ²⁰	RCT	Italy	HCC	1.9±0.8	RFA	52(69)	36/16	67±6 (52-78)	2.8±0.6	1.00(1y)	NA	1.00(1y)	15 pain and 10 fever
					PEI	50(73)	30/20	69±7.4 (40-82)	2.8±0.8	0.96(1y)	NA	0.96(1y)	13 pain and 5 fever
Lin 2004 ²¹	RCT	China	HCC	2±0.9	RFA	52(69)	35/17	62±11	2.9±0.8	0.76(3y)	NA	0.35(3y)	1 transient pleural effusion
					PEI	52(67)	34/18	59±10	2.8±0.8	0.66(3y)	NA	0.17(3y)	1 pain
Vivarelli 2004 ²²	Retrospective cohort	Italy	HCC	2.4	RES	79(92)	57/22	65.2±8.2 (43-81)	≤3/3.1-5 (21/58)	0.81(3y)	0.59(3y)	0.65(3y)	NA
					RFA	79(112)	67/12	67.8±8.7 (41-88)	≤3/3.1-5 (22/57)	0.50(3y)	0.25(3y)	0.33(3y)	NA

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Cho 2005 ²³	Retrospect ive cohort	Korea	HCC	0.1-3	RES	61	48/13	57	3.4±1.0	NA	0.77(3y)	0.77(3y)	2 bleeding, 1 intraabdominal abscess, 1 wound infection
					RFA	99	76/23	58	3.1±0.8	NA	0.80(3y)	0.80(3y)	1 chest wall metastasis, 1 cholecystitis, 1 iatrogenic burn, 1 ileus, 1 hepatic infarction
Huang 2005 ²⁵	RCT	China	HCC	1-4.9	RES	38(42)	27/11	59±11.4	≤2/2.1-3 (24/14)	0.82	NA	0.82	NA
					PEI	38(46)	19/19	63±10.9	≤2/2.1-3 (21/17)	0.45	NA	0.45	NA
Hong 2005 ²⁴	Retrospect ive cohort	Korea	HCC	2.9(0.4-4.6)	RES	93	69/24	49.2±9.9	2.5±0.8	0.84(3y)	NA	0.84(3y)	NA
					RFA	55	41/14	59.1±9.6	2.4±0.6	0.73(3y)	NA	0.73(3y)	NA
Lin 2005 ²⁶	RCT	China	HCC	2.3±1	RFA	62(78)	40/22	61±10	2.5±1	0.74(3y)	NA	0.74(3y)	2 haemothorax, 1 gastric bleeding and perforation
					PEI	62(76)	39/23	60±8	2.3±0.8	0.60(3y)	NA	0.60(3y)	1 pain
Lu 2005 ²⁷	Retrospect ive cohort	China	HCC	2.1±1.1	RFA	53(72)	43/10	54.5±11.7 (24-74)	2.6±1.2 (1.0-6.1)	0.38(3y)	NA	0.38(3y)	2 skin burn, 1 puncture wound infection
					MWA	49(98)	44/5	50.1±13.7 (24-74)	2.5±1.2 (0.9-7.2)	0.51(3y)	NA	0.51(3y)	2 puncture wounds, 2 subcapsular hematoma
Montorsi 2005 ²⁸	Prospectiv e cohort	Italy	HCC	2.1	RES	40	33/7	67±9	<5cm	NA	NA	0.73(3y)	NA
					RFA	58	43/15	67±6		NA	NA	0.60(3y)	NA
Shiina 2005 ²⁹	RCT	Japan	HCC	3.1(0.6-4.3)	RFA	118(184)	79/39	≤65/>65 (44/74)	≤2/>2 (45/73)	NA	NA	0.61(3y)	1 transient jaundice, 1 skin burn, 1 hepatic infarction, 3 neoplastic seeding

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					PEI	114(188)	87/27	≤65/>65 (41/73)	≤2/>2 (57/57)	NA	NA	0.45(3y)	1 abscess2 neoplastic seeding
Chen 2006 ³⁰	RCT	China	HCC	2.4±1	RES	90	75/15	49.4±10.9	≤3/3.1-5 (42/48)	0.53	NA	0.53	2 liver failure, 2 gastrointestinal bleeding, 27 ascites
					RFA	71	56/15	51.9±11.2	≤3/3.1-5 (37/34)	0.58	NA	0.58	3 skin burn
Lu 2006 ³¹	RCT	China	Early HCC	1.8	RES	54(56)	37/17	49±14	3.2±1.0	NA	NA	0.86 (3y)	3 wound infection, 1 gastrointestinal bleeding
					RFA	51(57)	42/9	55±13	2.7±1.0	NA	NA	0.87 (3y)	1 peritoneal bleeding, 1 neoplastic seeding
Cho 2007 ³²	Retrospect ive cohort	Korea	HCC	5.7	RES	130(145)	103/27	56.3±8.8	≤2/2.1-3 (43/87)	0.66	NA	0.66	NA
					PEI	249(275)	181/68	57.7±9.7	≤2/2.1-3 (169/80)	0.49	NA	0.49	NA
Gao 2007 ³³	Retrospect ive cohort	China	HCC	4.6	RES	34(37)	28/6	51.5 (38-67)	2.58±0.41	0.76	NA	0.76	12 fever, 5 ascites
					RFA	53(84)	41/12	57.1 (31-81)	2.45±0.37	0.62	NA	0.62	2 bleeding, 1 fistula, 1 wound infection, 6 fever, 9 ascites
Lupo 2007 ³⁴	Retrospect ive cohort	Italy	HCC	2.6	RES	42	33/9	67(28-80)	4.0(3-5)	NA	0.43	0.43	2 urine infection, 1 bilioma, 1 pleural effusion, 1 renal failure, 1 intra-abdominal bleeding
					RFA	60	47/13	68(42-85)	3.65(3-5)	NA	0.32	0.32	2 liver failure, 1 hepatic abscess, 2 pleural effusion, 1 cutaneous metastasis
Zhou 2007 ³⁵	Retrospect ive cohort	China	HCC	0.5-5.9	RES	40(42)	35/5	53±13	≤2/2.1-5 (7/33)	NA	NA	0.75	NA

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	47(54)	37/10	57 ±14	≤2/2.1-5 (8/39)	NA	NA	0.19	NA
Abu-Hilal 2008 ³⁶	Retrospective cohort	Italy and China	Early HCC	3.6	RES	34	26/8	67	3.8(1.3-5)	NA	0.56	0.56	3 hepatic failure
					RFA	34	27/7	65	3(2-5)	NA	0.56	0.56	1 artero-portal fistula
Brunello 2008 ³⁷	RCT	Italy	Early HCC	2.2	RFA	70(89)	49/20	70.3±8.1	1.27±0.54	0.60(3y)	NA	0.60(3y)	1 haemoperitoneum 1 right haemothorax
					PEI	69(88)	43/27	69.0±7.7	1.27±0.57	0.58(3y)	NA	0.58(3y)	1 haemoperitoneum 1 death
Guglielmi 2008 ³⁸	Retrospective cohort	Italy	HCC	2.3	RES	91(113)	73/18	≤65/>65 (47/44)	≤3/3.1-6 (31/60)	0.55	0.43	0.48	33 postoperative complications
					RFA	109(153)	88/21	≤65/>65 (38/71)	≤3/3.1-6 (32/77)	0.28	0.14	0.20	11 postoperative complications
Hiraoka 2008 ³⁹	Retrospective cohort	Japan	HCC	2.5	RES	59	44/15	62.4±10.6	2.27±0.55	0.59	NA	0.59	1 death, 2 abscess
					RFA	105	76/29	69.4±9.1	1.98±0.52	0.59	NA	0.59	1 biloma, 2 dermatitis
Bu 2009 ⁴⁵	Retrospective cohort	China	HCC	2.9(0.5-6)	RES	42(46)	36/6	53.93±10.74	≤3/3.1-5 (14/28)	0.57	0.46	0.50	1 postoperative hemorrhage, 3 pleural effusions, 2 subdiaphragmatic effusion
					RFA	46(54)	40/6	55.89±7.37	≤3/3.1-5 (20/26)	0.50	0.31	0.37	4 pleural effusions, 1 postoperative hemorrhage, 1 skin burn

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Ohmoto 2009 ⁴⁰	Retrospect ive cohort	Japan	HCC	2.8±2	RFA	34(37)	25/9	67 (44-78)	1.6 (0.7-2.0)	0.71	NA	0.71	2 pain, 4 fever, 1 bile duct injury, 1 pleural effusion, 1 skin burns, 1 vagovagal reflex
					MWA	49(56)	41/8	64 (38-75)	1.7 (0.8-2.0)	0.37	NA	0.37	11 pain, 17 fever, 9 bile duct injury, 8 pleural effusion, 5 ascites, 4 skin burns, 2 vagovagal reflex, 2 abscess, 2 intrapertitoneal bleeding, 1 hepatic infarction, 1 portal thrombus, 1 biliary peritonitis
Sakaguchi 2009 ⁴¹	Retrospect ive cohort	Japan	HCC	0.1-5	Laparosco pic /thorasc opic RFA	249	169/80	65.6±8.9	2.48±0.89	0.57	NA	0.57	1 frequent premature ventricular contractions, 1 liver decompensation
					Laparosco pic /thorasc opic MWA	142	107/35	64.9±7.8	2.28±0.74	0.63	NA	0.63	1 breath holding and incomplete intestinal obstruction, 2 liver decompensation
Santambrogio 2009 ⁴²	Prospectiv e cohort	Italy	HCC	3.2	RES	78	55/23	68±8	2.87±1.21	0.54	NA	0.54	15 extra-hepatic complications
					Laparosco pic RFA	74	59/15	68±7	2.63±1.07	0.41	NA	0.41	14 extra-hepatic complications
Shibata 2009 ⁴³	RCT	Japan	HCC	2.5±1.2	RFA	43(44)	33/10	69.8±8 (44-87)	1.6±0.5 (0.8-2.6)	0.84(3y)	NA	0.84(3y)	1 pseudoaneurysm

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					TR	46(49)	31/15	67.2±8.9 (45-83)	1.7±0.6 (0.9-3.0)	0.85(3y)	NA	0.85(3y)	1 hepatic infarction
Ueno 2009 ⁴⁴	Retrospective cohort	Japan	HCC	3(0.3-7.9)	RES	123(136)	82/41	67(28-85)	2.7±0.1	0.81	0.72	0.80	NA
					RFA	155(209)	100/55	66(40-79)	2.0±0.1	0.38	0.78	0.63	NA
Guo 2010 ⁴⁶	Retrospective cohort	China	HCC	2.5	RES	73(155)	57/16	50.0 (17.0-68.0)	≤3/3.1-5 (30/43)	0.27	0.47	0.44	1 postoperative hemorrhage, 5 abscess, 3 infected ascites, 1 liver failure, 4 pleural effusion
					RFA	86(211)	63/23	52.5 (26.0-80.0)	≤3/3.1-5 (42/44)	0.33	0.16	0.21	1 postoperative hemorrhage, 1 bile leak, 1 abscess, 1 infected ascites, 3 pleural effusion
Huang 2010 ⁴⁷	RCT	China	HCC	3.87	RES	115(144)	85/30	55.91±12.68	≤3/3.1-5 (45/44)	0.82	0.73	0.76	1 hepatic failure, 13 ascites, 5 effusion, 9 bile leakage, 2 postoperative bleeding, 2 gastrointestinal bleeding
					RFA	115(147)	79/36	56.57±14.30	≤3/3.1-5 (57/27)	0.61	0.52	0.55	1 gastric perforation, 2 hemorrhage, 1 malignant seeding, 1 hepatic infarction
Kagawa 2010 ⁴⁸	Retrospective cohort	Japan	Early HCC	4.2	RES	55(69)	40/15	66.1±8.4	≤2/2.1-5 (9/46)	0.42	NA	0.42	2 deaths, 1 liver failure, 1 pleural effusion, 1 pneumonia, 2 biliary leakage
					TR	62(79)	39/23	67.5±8.4	≤2/2.1-5 (19/43)	0.29	NA	0.29	1 duodenal perforation, 1 hemothorax
Morimoto	RCT	Japan	HCC	2.7	RFA	18(25)	12/6	73 (48-84)	3.7±0.6	NA	0.78(3y)	0.78(3y)	5 pain, 2 pleural effusion

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
2010 ⁴⁹					TR	19(21)	15/4	70 (57-78)	3.6±0.7	NA	0.95(3y)	0.95(3y)	1 pain, 1 pleural effusion
Azab 2011 ⁵⁰	RCT	Egypt	HCC	1.5	RFA	30(33)	75/15	46-77	<5cm	NA	NA	0.90	5 superficial burn, 17 transient pain, 3 portal vein thrombosis, 7 fever, 1 ascites
					PEI	30(32)				NA	NA	0.83	2 portal vein thrombosis, 3 fever, 3 ascites
Giorgio 2011 ⁵¹	RCT	Italy	HCC	1.8	RFA	142	105/37	70±2 (68-74)	2.34±0.45 (1.1-3)	0.70	NA	0.70	1 major complication
					PEI	143	102/41	72±6 (68-79)	2.27±0.48 (1.3-2.9)	0.68	NA	0.68	3 major complication
Hung 2011 ⁵²	Retrospective cohort	China	Early HCC	3.5±2	RES	229	184/45	60.07±12.56	2.88±1.06	0.77	NA	0.77	NA
					RFA	190	121/69	67.42±11.45	2.37±0.92	0.67	NA	0.67	NA
Nishikawa 2011 ⁵³	Retrospective cohort	Japan	HCC	3.3	RES	69	50/19	67.4±9.7	2.68±0.49	0.74	NA	0.74	2 bile leakage, 2 ascites, 1 acute respiratory distress syndrome, 1 gastrointestinal bleeding
					RFA	162	95/67	68.4±8.7	1.99±0.62	0.63	NA	0.63	1 biloma, 1 ascites, 1 intra-abdominal bleeding
Yun 2011 ⁵⁴	Retrospective cohort	Korea	HCC	3.5(0.1-9.1)	RES	215	171/44	51.7±9.7	2.1±0.5	0.94	NA	0.94	NA
					RFA	255	197/58	57.0±9.9	2.1±0.5	0.87	NA	0.87	NA

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Zhang 2011 ⁵⁵	Retrospect ive cohort	China	HCC	0.5-3.5	RES	103(117)	78/25	56.4±15.2	<5cm	NA	NA	0.35(3y)	12 wound infection, 5 postoperative hemorrhage, 2 hepatic failure, 15 pleural effusions, 6 pleural effusions
					RFA	85(106)	62/23	58.5±12.9	<5cm	NA	NA	0.39(3y)	2 gallbladder cardiac reflex, 4 postoperative hemorrhage, 3 pleural effusions
Feng 2012 ⁵⁷	RCT	China	HCC	3	RES	84(116)	75/9	47 (18-76)	2.6±0.8	0.62(3y)	NA	0.62(3y)	7 pleural effusion, 3 pneumonia, 1 effusion plus infection, 3 wound infection or dehiscence, 1 biliary fistula, 2 abdominal bleeding, 1 pneumothorax or hemothorax
					RFA	84(120)	79/5	51 (24-83)	2.4±0.6	0.55(3y)	NA	0.55(3y)	5 pleural effusion, 1 liver abscess, 2 abdominal bleeding
Peng 2012 ⁵⁸	Retrospect ive cohort	China	Recurr ent HCC	4.9	RES	74	65/9	51.5±12.1 (24-75)	1.1±0.5 (0.8-2.0)	0.62	NA	0.62	1 liver failure, 2 gastrointestinal bleeding, 1 peritoneal bleeding, 1 intestinal obstruction, 1 spontaneous bacterial peritonitis, 1 persistent jaundice, 31 ascites
					RFA	71	63/8	53.1±12.1 (28-74)	1.2±0.6 (0.9-2.0)	0.72	NA	0.72	1 gastrointestinal bleeding, 1 persistent jaundice, 12 ascites

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Peng 2012 ⁵⁹	RCT	China	Recurrent HCC	3.3±1.8	RFA	70(76)	55/15	55.1±9.5 (22-75)	≤3/3.1-5 (46/24)	NA	0.17	0.36	1 persistent jaundice, 1 ascites, 22 fever, 45 pain, 4 vomiting
					TR	69(74)	59/9	57.5±10.0 (19-75)	≤3/3.1-5 (41/28)	NA	0.39	0.46	1 liver failure, 1 ascites, 27 fever, 50 pain, 42 vomiting
Signoriello 2012 ⁶⁰	Retrospective cohort	Italy	HCC	0.1-9	RES	34(44)	30/4	62±7	≤3/3.1-5/>5.1 (13/9/4)	NA	NA	0.29	NA
					RFA	50(74)	40/10	68±7	≤3/3.1-5/>5.1 (24/11/7)	NA	NA	0.15	NA
					PEI	256(349)	188/68	67±8	≤3/3.1-5/>5.1 (143/43/12)	NA	NA	0.20	NA
a. Wang 2012 ⁶¹	Retrospective cohort	China	Early HCC	2.5	RES	52	38/14	≤60 (35)	NA	NA	NA	0.92	NA
					RFA	91	60/31	≤60 (40)		NA	NA	0.73	NA
b. Wang 2012 ⁶²	Retrospective cohort	China	Early HCC	2.5	RES	208	168/40	≤60 (113)	≤2/2.1-5 (6/202)	NA	NA	0.77	NA
					RFA	254	161/93	≤60 (85)	≤2/2.1-5 (60/194)	NA	NA	0.57	NA
Desiderio 2013 ⁶²	Retrospective cohort	Italy	HCC	4.3(2.3-5)	RES	52(94)	37/15	65.6±4.8	≤3	0.46	NA	0.46	2 hepatic failure, 1 biliary fistula, 2 hemoperitoneum, 9 ascites
					RFA	44(81)	35/9	64.4±6.5		0.36	NA	0.36	6 pain, 7 fever
Ding 2013 ⁶³	Retrospective cohort	China	HCC	2.3±1.3	RFA	85(98)	68/17	58.64±8.52 (40-77)	2.38±0.81 (1.0-4.8)	0.82(3y)	NA	0.82(3y)	1 frequent premature ventricular contractions, 1 liver decompensation

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					MWA	113(131)	85/28	59.06±11.68 (30-86)	2.55±0.89 (0.8-5.0)	0.78(3y)	NA	0.78(3y)	1 breath holding and incomplete intestinal obstruction, 2 liver decompensation
Guo 2013 ⁶⁴	Retrospect ive cohort	China	HCC	2.7	RES	102(129)	94/8	51.5(18-75)	≤3/3.1-5 (75/27)	NA	NA	0.63	5 postoperative hemorrhage, 3 bile leak, 4 abscess, 3 infected ascites, 1 liver failure, 4 pleural effusion
					RFA	94(125)	78/16	56(19-75)	≤3/3.1-5 (62/32)	NA	NA	0.50	1 postoperative hemorrhage, 2 bile leak, 1 abscess, 1 infected ascites, 3 pleural effusion
Hasegawa 2013 ⁶⁵	Retrospect ive cohort	Japan	HCC	2.2	RES	5361(646 1)	3967/139 4	66 (48-77)	2.3 (1.2-3)	0.71	NA	0.71	NA
					RFA	5548(741 2)	3569/197 9	69 (52-80)	2 (1-3)	0.61	NA	0.61	NA
					PEI	2059(283 6)	1303/756	69 (52-80)	1.7 (1-3)	0.56	NA	0.56	NA
Iida 2013 ⁶⁶	Retrospect ive cohort	Japan	HCC	0.1-7.5	Laparosco pic RFA	18(27)	NA	73.5±4.0	2.1±0.5	0.78	NA	0.78	1 abscess
					Laparosco pic MWA	40(56)		70.1±6.6	2.0±0.9	0.78	NA	0.78	1 abscess
Imai 2013 ⁶⁷	Retrospect ive cohort	Japan	HCC	4.1	RES	101	75/26	63.3±9.7	2.14±0.55	0.87	NA	0.87	NA
					RFA	82	46/36	67.6±8.5	1.87±0.50	0.60	NA	0.60	NA

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Kim 2013 ⁶⁸	Retrospective cohort	Korea	Early HCC	0.1-4.2	RES	47	36/11	58.8±10.7	3.66±0.76	NA	0.85(3y)	0.85(3y)	2 pleural effusion, 2 pneumonia, 1 hepatic failure, 1 hepatic abscess, 1 mechanical ileus
					TR	37	31/6	61.7±11.1	3.46±0.75	NA	0.78(3y)	0.78(3y)	1 bile duct dilatation
Lai 2013 ⁶⁹	Retrospective cohort	China	HCC	2.9±1.5	RES	80	55/25	60.8±9.9	2.9±1.1	0.71	NA	0.71	NA
					RFA	31	19/12	63.1±12.8	1.8±0.6	0.84	NA	0.84	NA
Lin 2013 ⁷⁰	Retrospective cohort	China	Early HCC	3.4	RFA	658	393/265	64.7±10.5	2.4±1.1 (0.8-9.5)	0.60	0.50	0.55	NA
					PEI	378	243/135	63.5±12.1	2.0±0.9 (0.4-7.0)	0.50	0.28	0.40	NA
Peng 2013 ⁷¹	RCT	China	HCC	0.6-5.2	RFA	95(133)	71/24	55.3±13.3	3.39±1.35	NA	0.59(3y)	0.59(3y)	51 pain, 26 fever, 29 vomiting, 4 ascites, 2 pleural effusion, 1 skin burn, 1 abdominal infection, 1 small intestinal obstruction
					TR	94(137)	75/19	53.3±11	3.47±1.44	NA	0.67(3y)	0.67(3y)	57 pain, 33 fever, 40 vomiting, 5 ascites, 3 pleural effusion, 1 skin burn, 1 bile duct stenosis, 1 gastric hemorrhage
Tohme 2013 ⁷²	Retrospective cohort	America	Early HCC	2.4	RES	50(62)	31/19	66.3±1	3.07±1.17	0.48	NA	0.48	3 pleural effusion, 1 pneumonia, 1 myocardial infarction, 2 biloma, 2 ileus, 1 ascites, 1 hyperbilirubinaemia >6, 1 renal insufficiency, 2 encephalopathy

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	60(75)	38/22	65.6±12	2.36±0.94	0.35	NA	0.35	1 oesophagitis, 3 encephalopathy, 1 cholangitis, 2 ascites, 1 renal insufficiency, 1 pneumonia
Wong 2013 ⁷³	Retrospect ive cohort	China	Early HCC	0.1-5	RES	46	30/16	55.1±12	2.1±0.6	0.85	NA	0.85	2 fever, 1 increased serum alanine aminotransferase level, 2 atelectasis, 2 biloma
					RFA	36	18/18	63.5±13	1.9±0.6	0.72	NA	0.72	None
Zhang 2013 ⁷⁴	Retrospect ive cohort	China	HCC	2.2±1	RFA	78(97)	64/14	54±10.5 (30-80)	≤3/3.1-5 (47/31)	0.43	0.39	0.41	1 persistent jaundice, 1 biliary fistula
					MWA	77(105)	67/10	54±9.5 (26-76)	≤3/3.1-5 (36/41)	0.58	0.29	0.39	1 hemothorax and intrahepatic hematoma, 1 peritoneal hemorrhage
Abdelaziz 2014 ⁷⁵	RCT	Egypt	Early HCC	2.3	RFA	45(52)	31/14	56.8±7.3	2.95±1.03	0.68(1y)	NA	0.68(1y)	2 subcapsular hematoma, 1 thigh burn, 2 pleural effusion
					MWA	66(76)	48/18	53.6±5	2.9±0.97	0.96(1y)	NA	0.96(1y)	1 subcapsular hematoma, 1 abdominal wall skin burn
Shi 2014 ⁷⁶	Retrospect ive cohort	China	HCC	3.8	RES	107(126)	87/20	54.5±9.9	≤3/3.1-5 (37/54)	0.73	0.57	0.60	NA
					MWA	117(143)	93/24	56.6±9.2	≤3/3.1-5 (40/56)	0.65	0.52	0.52	NA

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Yang 2014 ⁷⁷	Retrospective cohort	Korea	HCC	0.1-7	RES	52	38/14	55.7±10.6	≤2/2.1-5 (21/31)	0.94	NA	0.94	2 pneumonia, 1 wound infection, 1 biliary anastomotic leak, 1 portal vein thrombosis, 1 nausea, 1 delirium, 4 ascites
					RFA	79	59/20	57.2±9.2	≤2/2.1-5 (36/43)	0.86	NA	0.86	1 vomiting, 1 ascites, 6 abdominal pain, 2 nausea, 1 sinus bradycardia
Zhang 2014 ⁷⁸	Retrospective cohort	China	Recurrent HCC	2.7	RES	27(29)	25/2	47±13	3.2±1.0	NA	NA	0.63	NA
					MWA	39(46)	37/2	52±13	2.7±1.1	NA	NA	0.62	NA
Pompili 2015 ⁷⁹	Retrospective cohort	Italy	Early HCC	2.8	RFA	136	75/61	68 (41-85)	1.8 (1-2)	0.63	NA	0.63	2 ascites, 1 pleural effusion, 1 hemobilia
					PEI	108	90/18	68.5 (34-86)	1.95 (0.8-2)	0.65	NA	0.65	1 hemobilia, 1 portal vein thrombosis
Xu 2015 ⁸⁰	RCT	China	HCC	0.1-3	Laparoscopic RES	45	34/11	58.3±3.1 (26-78)	3.6±0.7 (1-5)	NA	0.38(3y)	0.38(3y)	3 bile leakage, 3 pleural effusion, 2 postoperative hemorrhage
					MWA	45	32/13	57.9±3.4 (27-76)	3.8±0.9 (2-5)	NA	0.33(3y)	0.33(3y)	1 bile leakage, 1 pleural effusion, 1 postoperative hemorrhage

HCC: hepatocellular carcinoma;

1 BCLC: Barcelona Clinic Liver Cancer;
2 RES: resection;
3 RFA: radiofrequency ablation;
4 MWA: microwave ablation;
5 TR: transcatheter arterial chemoembolization and radiofrequency ablation;
6 PEI: percutaneous ethanol injection;
7 RCT: randomized controlled trial;
8 NA: not available.
9
10
11
12

13 **Table S2.**
14 **Quality assessment of included studies using GRADE framework.**
15
16

Intervention/Comparator	Illustrative comparative risks* (per 1000, 95% CI)			Relative effect of survival time (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Comparator	Assumed survival risk	Corresponding survival risk with intervention			
1-year OS rate						
ES/MWA	923	984 (932 to 997)		OR 5.25 (1.15 to 23.97)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
FA/MWA	947	944 (902 to 968)		OR 0.94 (0.52 to 1.71)	990 (6 studies)	⊕ ⊕ ⊖ ⊖ low
ES/PEI	835	802 (674 to 889)		OR 0.80 (0.41 to 1.58)	519 (3 studies)	⊕ ⊕ ⊖ ⊖ low
FA/PEI	944	963 (906 to 1000)		OR 1.02 (0.96 to 1.09)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
ES/RFA	932	945 (931 to 956)		OR 1.25 (0.99 to 1.60)	5006 (30 studies)	⊕ ⊕ ⊕ ⊕ high

RES/TR	939	904 (765 to 965)	OR 0.61 (0.21 to 1.79)	201 (2 studies)	⊕ ⊕ ⊖ ⊖ low
2					
3					
RFA/TR	938	802 (310 to 978)	OR 0.27 (0.03 to 2.90)	31 (1 study)	⊕ ⊕ ⊖ ⊖ low
4					
5					
6					
3-year OS rate					
7					
RES/MWA	712	734 (623 to 822)	OR 1.12 (0.67 to 1.87)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
8					
9					
10					
11					
RFA/MWA	736	779 (717 to 828)	OR 1.26 (0.91 to 1.73)	987 (6 studies)	⊕ ⊕ ⊖ ⊖ low
12					
13					
14					
RES/PEI	499	536 (421 to 645)	OR 1.16 (0.73 to 1.83)	519 (3 studies)	⊕ ⊕ ⊖ ⊖ low
15					
16					
17					
RFA/PEI	729	748 (657 to 822)	OR 1.10 (0.71 to 1.71)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
18					
19					
20					
RES/RFA	785	851 (823 to 875)	OR 1.57 (1.28 to 1.93)	15906 (30 studies)	⊕ ⊕ ⊕ ⊖ moderate
21					
22					
23					
RES/TR	798	760 (618 to 860)	OR 0.80 (0.41 to 1.55)	201 (2 studies)	⊕ ⊕ ⊖ ⊖ low
24					
25					
26					
27					
RFA/TR	737	611 (516 to 704)	OR 0.56 (0.38 to 0.85)	454 (4 studies)	⊕ ⊕ ⊕ ⊖ moderate
28					
29					
5-year OS rate					
30					
31					
RES/MWA	545	607 (492 to 712)	OR 1.29 (0.81 to 2.07)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
32					
33					
34					
RFA/MWA	545	609 (442 to 756)	OR 1.30 (0.66 to 2.58)	687 (4 studies)	⊕ ⊕ ⊖ ⊖ low
35					
36					
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					

RES/PEI	293	436 (334 to 545)	OR 1.87 (1.21 to 2.90)	519 (3 studies)	⊕ ⊕ ⊕ ⊖ moderate
RFA/PEI	533	496 (368 to 624)	OR 0.86 (0.51 to 1.45)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
RES/RFA	601	744 (705 to 779)	OR 1.93 (1.59 to 2.34)	15154 (25 studies)	⊕ ⊕ ⊕ ⊖ moderate
RES/TR	290	419 (251 to 607)	OR 1.76 (0.82 to 3.78)	117 (1 study)	⊕ ⊕ ⊖ ⊖ low
RFA/TR	464	356 (222 to 523)	OR 0.64 (0.33 to 1.27)	139 (1 study)	⊕ ⊕ ⊕ ⊖ moderate

The absolute and relative risk of survival with treatments*. GRADE: Grading of Recommendations, Assessment, Development and Evaluation. *The results presented in the Table S1 were built around the assumption of a consistent relative effect. The implications of this effect for populations were considered at different baseline risks. Based on the assumed risks, corresponding risks after an intervention were calculated using the meta-analytic risk ratio.

Table S3.
Ranking treatments of 1-, 3-year and 5-year survival rate of the lesions < 3 cm, 3-5 cm and ≤ 5 cm in RCT.

Treatment	1-year			3-year			5-year		
	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank
< 3cm	12			10			4		
RES		2	3.06		1	1.80		1	1.25
RFA		3	3.21		3	2.56		2	2.08
MWA		1	1.14		NA	NA		NA	NA
TR		4	3.22		2	2.38		NA	NA
PEI		5	4.36		4	3.26		3	2.68

3-5cm	4			4			2		
RES		1	1.17		1	1.19		1	1.69
RFA		3	2.88		3	2.91		3	2.60
MWA		NA	NA		NA	NA		NA	NA
TR		2	1.94		2	1.90		2	1.71
PEI		NA	NA		NA	NA		NA	NA
All tumours (\leq 5cm)	18			14			5		
RES		3	2.78		2	2.43		1	1.68
RFA		4	3.91		3	3.52		3	2.75
MWA		1	1.62		4	3.10		NA	NA
TR		2	1.79		1	1.68		2	2.09
PEI		5	4.90		5	4.27		4	3.48

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection.

Table S4.

Ranking treatments of 1-, 3-year and 5-year survival rate of the lesions < 3 cm, 3-5 cm and \leq 5 cm in all studies.

Treatment	1-year			3-year			5-year		
	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank
< 3cm	44			42			31		
RES		3	3.02		2	2.49		1	1.35
RFA		4	3.16		3	3.44		2	3.03
MWA		2	2.19		4	3.52		4	3.31
TR		1	2.05		1	1.66		3	3.18

1	PEI	5	4.58	5	3.89	5	4.13
2	3-5cm	17		16		11	
3	RES	1	1.23	1	1.10	1	1.93
4	RFA	4	3.52	3	3.43	3	3.18
5	MWA	3	3.46	4	3.72	4	3.43
6	TR	2	1.97	2	2.10	2	1.94
7	PEI	5	4.82	5	4.66	5	4.53
8							
9	All tumours (\leq 5cm)	62		57		40	
10	RES	2	2.34	2	2.18	1	1.32
11	RFA	3	3.27	3	3.48	3	3.36
12	MWA	4	3.78	4	3.98	4	3.51
13	TR	1	1.10	1	1.27	2	2.45
14	PEI	5	4.52	5	4.10	5	4.36
15							
16							
17							
18							
19							
20							
21							
22							
23							
24							
25							
26							
27							
28							
29							
30							
31							
32							
33							
34							
35							
36							
37							
38							
39							
40							
41							
42							
43							
44							
45							
46							

RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection.

Table S5.
Survival rates (1-year, 3-year and 5-year) for small lesion (<3cm) treatment comparisons estimated by direct and network meta-analysis in RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	1.01 (0.40-2.14)	0.98 (0.77-1.26)
MWA vs RES	161.8 (1.39-581.0)	NA
TR vs RES	15.61 (0.02-54.78)	NA
PEI vs RES	0.68 (0.19-1.76)	1.03 (0.54-1.94)

MWA vs RFA	154.8 (1.74-590.1)	1.42 (0.63-3.19)
TR vs RFA	13.24 (0.02-55.15)	1.00 (0.56-1.80)
PEI vs RFA	0.68 (0.28-1.36)	0.97 (0.78-1.19)
TR vs MWA	1.42 (0-5.94)	NA
PEI vs MWA	0.08 (0-0.42)	NA
PEI vs TR	10.75 (0.01-29.11)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.86 (0.40-1.68)	0.92 (0.71-1.19)
MWA vs RES	NA	NA
TR vs RES	1.44 (0.14-5.50)	NA
PEI vs RES	0.75 (0.28-1.89)	1.21 (0.59-2.15)
MWA vs RFA	NA	NA
TR vs RFA	1.64 (0.20-5.84)	1.01 (0.55-1.87)
PEI vs RFA	0.88 (0.44-1.79)	0.91 (0.71-1.17)
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	1.29 (0.13-4.99)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.71 (0.10-2.47)	0.93 (0.62-1.37)
MWA vs RES	NA	NA
TR vs RES	NA	NA
PEI vs RES	0.49 (0.04-2.02)	0.55 (0.26-1.15)
MWA vs RFA	NA	NA
TR vs RFA	NA	NA
PEI vs RFA	0.93 (0.08-3.85)	0.97 (0.66-1.40)
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA

Table S6.

Survival rates (1-year, 3-year and 5-year) for lesion (3-5cm) treatment comparisons estimated by direct and network meta-analysis in

RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.25 (0-1.47)	0.89 (0.45-1.77)
MWA vs RES	NA	NA
TR vs RES	1.00 (0-5.0)	NA
PEI vs RES	NA	NA
MWA vs RFA	NA	NA
TR vs RFA	3.40 (0.64-11.93)	1.10 (0.78-1.55)
PEI vs RFA	NA	NA
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.24 (0-1.25)	0.70 (0.34-1.45)
MWA vs RES	NA	NA
TR vs RES	1.14 (0-6.20)	NA
PEI vs RES	NA	NA
MWA vs RFA	NA	NA
TR vs RFA	3.98 (0.71-15.22)	1.29 (0.87-1.89)
PEI vs RFA	NA	NA
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA
5-year OS rate for treatment vs reference		
RFA vs RES	1.05 (0.03-5.33)	0.71 (0.32-1.57)
MWA vs RES	NA	NA
TR vs RES	12.87 (0.02-44.43)	NA
PEI vs RES	NA	NA

MWA vs RFA	NA	NA
TR vs RFA	7.64 (0.14-42.49)	1.93 (0.53-7.06)
PEI vs RFA	NA	NA
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA

Table S7.

Survival rates (1-year, 3-year and 5-year) for lesion (≤ 5 cm) treatment comparisons estimated by direct and network meta-analysis in RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.65 (0.28-1.31)	0.96 (0.77-1.20)
MWA vs RES	2.75 (0.52-9.18)	0.98 (0.54-1.78)
TR vs RES	2.15 (0.49-6.46)	NA
PEI vs RES	0.42 (0.14-0.98)	1.03 (0.54-1.94)
MWA vs RFA	4.62 (0.85-15.59)	1.42 (0.63-3.19)
TR vs RFA	3.3 (1.05-8.21)	1.09 (0.84-1.43)
PEI vs RFA	0.65 (0.32-1.14)	0.95 (0.80-1.14)
TR vs MWA	1.26 (0.14-4.73)	NA
PEI vs MWA	0.24 (0.03-0.81)	NA
PEI vs TR	0.26 (0.06-0.69)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.80 (0.36-1.69)	0.87 (0.69-1.10)
MWA vs RES	1.18 (0.16-4.30)	0.88 (0.39-1.98)
TR vs RES	1.69 (0.47-4.87)	NA
PEI vs RES	0.66 (0.23-1.78)	1.12 (0.59-2.15)
MWA vs RFA	1.71 (0.17-6.61)	NA
TR vs RFA	2.09 (0.81-4.65)	1.20 (0.90-1.60)

1	PEI vs RFA	0.83 (0.39-1.73)	0.84 (0.66-1.07)
2	TR vs MWA	3.25 (0.24-14.23)	NA
3	PEI vs MWA	1.25 (0.11-5.36)	NA
4	PEI vs TR	0.49 (0.13-1.33)	NA
5	5-year OS rate for treatment vs reference		
6	RFA vs RES	0.72 (0.11-2.48)	0.85 (0.61-1.17)
7	MWA vs RES	NA	NA
8	TR vs RES	2.96 (0.05-14.7)	NA
9	PEI vs RES	0.49 (0.04-2.03)	0.55 (0.26-1.15)
10	MWA vs RFA	NA	NA
11	TR vs RFA	3.59 (0.14-18.06)	1.30 (0.70-2.41)
12	PEI vs RFA	0.90 (0.08-3.65)	0.97 (0.66-1.40)
13	TR vs MWA	NA	NA
14	PEI vs MWA	NA	NA
15	PEI vs TR	1.51 (0.02-7.71)	NA

OR: odds ratio;
RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection;
NA: not available.

Table S8.
Survival rates (1-year, 3-year and 5-year) for small lesion (<3cm) treatment comparisons estimated by direct and network meta-analysis in all studies.

Intervention	OR (95% CI)	
	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		

RFA vs RES	1.03 (0.42-2.07)	1.00(0.95-1.05)
MWA vs RES	1.55 (0.41-4.10)	1.00(0.53-1.89)
TR vs RES	2.51 (0.26-9.65)	1.00(0.56-1.80)
PEI vs RES	0.71 (0.24-1.60)	1.00 (0.93-1.07)
MWA vs RFA	1.51 (0.60-3.11)	1.02 (0.85-1.23)
TR vs RFA	2.45 (0.33-8.72)	1.00(0.56-1.80)
PEI vs RFA	0.69 (0.39-1.13)	0.99 (0.93-1.06)
TR vs MWA	1.96 (0.21-7.87)	NA
PEI vs MWA	0.55 (0.18-1.29)	NA
PEI vs TR	0.56 (0.07-2.13)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.85 (0.40-1.62)	0.94 (0.90-0.99)
MWA vs RES	0.87 (0.31-1.96)	0.96 (0.49-1.87)
TR vs RES	1.87 (0.40-5.56)	1.17 (0.67-2.04)
PEI vs RES	0.80 (0.33-1.68)	1.00 (0.71-1.40)
MWA vs RFA	1.02 (0.54-1.76)	1.00 (0.82-1.22)
TR vs RFA	2.21 (0.60-5.76)	1.01 (0.55-1.87)
PEI vs RFA	0.95 (0.59-1.47)	0.97 (0.90-1.03)
TR vs MWA	2.35 (0.54-6.80)	NA
PEI vs MWA	1.01 (0.45-2.00)	NA
PEI vs TR	0.59 (0.15-1.67)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.58 (0.24-1.11)	0.86 (0.81-0.90)
MWA vs RES	0.58 (0.18-1.33)	0.89 (0.44-1.79)
TR vs RES	0.72 (0.11-2.48)	0.69 (0.34-1.42)
PEI vs RES	0.46 (0.18-0.95)	0.79 (0.73-0.85)
MWA vs RFA	1.00 (0.50-1.77)	1.02 (0.78-1.33)
TR vs RFA	1.24 (0.25-3.80)	NA
PEI vs RFA	0.81 (0.48-1.28)	0.92 (0.85-0.99)
TR vs MWA	1.37 (0.23-4.59)	NA
PEI vs MWA	0.90 (0.38-1.83)	NA

PEI vs TR	1.06 (0.19-3.41)	NA
-----------	------------------	----

Table S9.
Survival rates (1-year, 3-year and 5-year) for lesion (3-5cm) treatment comparisons estimated by direct and network meta-analysis in all studies.

Intervention	OR (95%CI)	
	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.19 (0-1.18)	0.96 (0.78-1.17)
MWA vs RES	0.24 (0-1.61)	NA
TR vs RES	0.56 (0-3.31)	1.02 (0.55-1.88)
PEI vs RES	0.10 (0-0.63)	NA
MWA vs RFA	1.25 (0.31-3.46)	0.98 (0.49-1.95)
TR vs RFA	2.92 (1.14-6.65)	1.11 (0.80-1.54)
PEI vs RFA	0.50 (0.17-1.13)	0.89 (0.66-1.20)
TR vs MWA	3.46 (0.57-11.35)	NA
PEI vs MWA	0.60 (0.09-1.94)	NA
PEI vs TR	0.21 (0.04-0.56)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.14 (0.01-0.68)	0.78 (0.62-0.98)
MWA vs RES	0.15 (0-0.77)	1.02 (0.57-1.81)
TR vs RES	0.36 (0.01-1.73)	0.92 (0.48-1.75)
PEI vs RES	0.09 (0-0.44)	NA
MWA vs RFA	1.01 (0.25-2.72)	0.60 (0.26-1.36)
TR vs RFA	2.37 (0.90-5.53)	1.29 (0.87-1.89)
PEI vs RFA	0.57 (0.10-1.83)	0.71 (0.50-1.00)
TR vs MWA	3.48 (0.62-11.64)	NA
PEI vs MWA	0.90 (0.08-3.36)	NA
PEI vs TR	0.30 (0.03-1.06)	NA
5-year OS rate for treatment vs reference		

RFA vs RES	0.91 (0.05-4.18)	0.62 (0.45-0.85)
MWA vs RES	1.79 (0.03-5.39)	0.90 (0.48-1.69)
TR vs RES	14.49 (0.05-27.29)	NA
PEI vs RES	1.88 (0.01-3.18)	NA
MWA vs RFA	1.25 (0.18-3.84)	0.57 (0.21-1.51)
TR vs RFA	7.08 (0.25-26.41)	2.36 (0.66-8.37)
PEI vs RFA	0.79 (0.05-2.64)	0.56 (0.37-0.84)
TR vs MWA	13.88 (0.19-50.64)	NA
PEI vs MWA	1.88 (0.04-5.54)	NA
PEI vs TR	6.11 (0-3.02)	NA

Table S10.

Survival rates (1-year, 3-year and 5-year) for lesion (≤ 5 cm) treatment comparisons estimated by direct, indirect and network meta-analysis in all studies.

Intervention	OR (95%CI)	
	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.78 (0.37-1.49)	0.99 (0.95-1.04)
MWA vs RES	0.73 (0.28-1.55)	0.95 (0.71-1.27)
TR vs RES	2.35 (0.74-5.96)	1.04 (0.70-1.55)
PEI vs RES	0.61 (0.26-1.25)	1.01 (0.74-1.39)
MWA vs RFA	0.95 (0.48-1.67)	1.01 (0.85-1.21)
TR vs RFA	3.01 (1.33-6.15)	1.10 (0.85-1.43)
PEI vs RFA	0.78 (0.51-1.13)	0.98 (0.93-1.05)
TR vs MWA	3.51 (1.78-8.52)	0.91 (0.70-1.18)
PEI vs MWA	0.91 (0.41-1.79)	NA
PEI vs TR	0.30 (0.11-0.63)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.78 (0.44-1.29)	0.93 (0.89-0.98)
MWA vs RES	0.72 (0.36-1.32)	0.96 (0.69-1.32)

1	TR vs RES	1.50 (0.64-3.08)	1.06 (0.69-1.61)
2	PEI vs RES	0.71 (0.37-1.30)	0.93 (0.86-1.00)
3	MWA vs RFA	0.94 (0.58-1.44)	0.95 (0.78-1.16)
4	TR vs RFA	1.93 (1.05-3.29)	1.20 (0.90-1.60)
5	PEI vs RFA	0.92 (0.63-1.32)	0.95 (0.89-1.01)
6	TR vs MWA	2.16 (0.99-4.16)	NA
8	PEI vs MWA	1.03 (0.56-1.77)	NA
9	PEI vs TR	0.52 (0.25-0.96)	NA
10	5-year OS rate for treatment vs reference		
12	RFA vs RES	0.56 (0.27-0.99)	0.84 (0.80-0.89)
13	MWA vs RES	0.56 (0.23-1.14)	0.90 (0.61-1.31)
14	TR vs RES	0.79 (0.24-1.92)	0.69 (0.34-1.42)
16	PEI vs RES	0.47 (0.22-0.87)	0.79 (0.73-0.85)
17	MWA vs RFA	1.01 (0.60-1.59)	0.97 (0.75-1.25)
18	TR vs RFA	1.42 (0.58-2.96)	1.30 (0.70-2.41)
19	PEI vs RFA	0.85 (0.57-1.22)	0.91 (0.84-0.98)
21	TR vs MWA	1.50 (0.52-3.46)	NA
22	PEI vs MWA	0.90 (0.47-1.58)	NA
23	PEI vs TR	0.71 (0.26-1.57)	NA

OR: odds ratio;
RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection;
NA: not available.

Table S11.
Posterior summaries from random effects consistency and inconsistency models for small lesion (<3cm) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.55	0.21	(0.15-1.00)	0.38	0.23	(0.02-0.88)
τ	11.06	88.80	(1.00-43.58)	4020	78840	(1.28-2366.00)
resdev	90.04	13.04	(66.16-117.10)	94.65	12.94	(70.06-120.70)
pD	59.96			57.5		
DIC	402.44			404.59		
3-year OS rate for treatment vs reference						
σ	0.59	0.14	(0.34-0.88)	0.6	0.14	(0.36-0.91)
τ	3.74	10.43	(1.29-8.74)	3.29	1.92	(1.21-8.05)
resdev	92.02	14.19	(66.64-122.10)	90.7	13.92	(65.64-120.00)
pD	70.71			71.74		
DIC	517.72			517.43		
5-year OS rate for treatment vs reference						
σ	0.53	0.12	(0.32-0.80)	0.55	0.13	(0.34-0.84)
τ	4.19	2.29	(1.57-9.74)	3.82	2.02	(1.42-8.83)
resdev	63.99	11.47	(43.52-88.24)	63.55	11.37	(43.39-87.90)
pD	54.24			54.99		
DIC	411.73			412.03		

Table S12.

Posterior summaries from random effects consistency and inconsistency models for lesion (3-5cm) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency Model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.28	0.25	(0.01-0.92)	0.38	0.34	(0.02-1.28)
τ	42220	1.30E+06	(1.19-19650.00)	19500.00	720600.00	(0.62-4178.00)

resdev	28.90	6.96	(17.25-44.41)	32.18	7.36	(19.64-48.32)
pD	22.80			24.59		
DIC	152.25			157.31		
3-year OS rate for treatment vs reference						
σ	0.62	0.27	(0.17-1.24)	0.67	0.31	(0.14-1.40)
τ	9.02	65.04	(0.66-35.66)	49.29	1164.00	(0.51-48.58)
resdev	32.36	8.17	(18.39-50.07)	32.62	8.22	(18.52-50.51)
pD	28.02			28.65		
DIC	187.98			188.88		
5-year OS rate for treatment vs reference						
σ	0.80	0.46	(0.14-1.94)	0.60	0.42	(0.04-1.64)
τ	49.88	1159	(0.27-49.16)	5839.00	185600.00	(0.37-748.40)
resdev	22.54	6.73	(11.29-37.43)	22.57	6.519	(11.45-36.90)
pD	20.62			19.84		
DIC	132.23			131.49		

Table S13.
Posterior summaries from random effects consistency and inconsistency models for lesion ($\leq 5\text{cm}$) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency Model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.49	0.13	(0.26-0.77)	0.29	0.14	(0.05-0.58)
τ	5.30	3.72	(1.70-14.33)	83.27	806.8	(2.94-391.70)
resdev	129.2	14.99	(101.40-160)	133.1	14.50	(105.70-162.80)
pD	84.95			78.28		
DIC	606.94			604.11		
3-year OS rate for treatment vs reference						
σ	0.50	0.09	(0.33-0.70)	0.47	0.096	(0.29-0.67)

τ	4.51	1.83	(2.08-9.02)	5.28	2.59	(2.24-11.80)
resdev	124	15.64	(95.16-156.40)	124.5	15.89	(95.35-157.50)
pD	93.89			93.37		
DIC	723.55			723.53		
5-year OS rate for treatment vs reference						
σ	0.44	0.10	(0.26-0.65)	0.44	0.1	(0.26-0.67)
τ	6.25	3.60	(2.38-14.90)	6.08	4.01	(2.25-14.87)
resdev	86.73	13.53	(62.35-115.40)	85.74	13.55	(61.39-114.40)
pD	67.86			68.84		
DIC	544.41			544.41		

sd: standard deviation;

CI: Credible Interval

σ : between-trial standard deviation

τ^2 : between-trial variance

resdev: residual deviance




pD: effective number of parameters

DIC: deviance information criterion

1 **Figure S1.**

2
3 **Results of the consistency test for closed loop at 1-year, 3-year, and 5-year survival rate of the lesions < 3 cm, 3-5 cm and ≤ 5 cm.**


- 4
5
6 i Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions < 3 cm
- 7
8
9 ii Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions 3-5 cm
- 10
11 iii Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions ≤ 5 cm
- 12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46



1 **Figure S2.**

2
3 **Assessment of publication bias using funnel plot.**

- 4
5
6 i Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions < 3 cm.
7
8
9 ii Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions 3-5 cm.
10
11 iii Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions \leq 5 cm
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

For peer review only

PRISMA NMA Checklist of Items to Include When Reporting A Systematic Review Involving a Network Meta-analysis

Section/Topic	Item #	Checklist Item	Reported on Page #
TITLE			
Title	1	Identify the report as a systematic review <i>incorporating a network meta-analysis (or related form of meta-analysis)</i> .	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: Background: main objectives Methods: data sources; study eligibility criteria, participants, and interventions; study appraisal; and <i>synthesis methods, such as network meta-analysis</i> . Results: number of studies and participants identified; summary estimates with corresponding confidence/credible intervals; <i>treatment rankings may also be discussed. Authors may choose to summarize pairwise comparisons against a chosen treatment included in their analyses for brevity.</i> Discussion/Conclusions: limitations; conclusions and implications of findings. Other: primary source of funding; systematic review registration number with registry name.	5,6
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known, <i>including mention of why a network meta-analysis has been conducted.</i>	7,8
Objectives	4	Provide an explicit statement of questions being addressed, with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	8

METHODS

Protocol and registration	5	Indicate whether a review protocol exists and if and where it can be accessed (e.g., Web address); and, if available, provide registration information, including registration number.	8,9
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale. <i>Clearly describe eligible treatments included in the treatment network, and note whether any have been clustered or merged into the same node (with justification).</i>	9,10
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	9
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	9,10,Figure1, Additional file 1: Text S1
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	9,10
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	10
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	11
Geometry of the network	S1	Describe methods used to explore the geometry of the treatment network under study and potential biases related to it. This should include how the evidence base has been graphically summarized for presentation, and what characteristics were compiled and used to describe the evidence base to readers.	11
Risk of bias within individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	11,12

Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means). <i>Also describe the use of additional summary measures assessed, such as treatment rankings and surface under the cumulative ranking curve (SUCRA) values, as well as modified approaches used to present summary findings from meta-analyses.</i>	11,12
Planned methods of analysis	14	Describe the methods of handling data and combining results of studies for each network meta-analysis. This should include, but not be limited to: <ul style="list-style-type: none"> • <i>Handling of multi-arm trials;</i> • <i>Selection of variance structure;</i> • <i>Selection of prior distributions in Bayesian analyses; and</i> • <i>Assessment of model fit.</i> 	11,12
Assessment of Inconsistency	S2	Describe the statistical methods used to evaluate the agreement of direct and indirect evidence in the treatment network(s) studied. Describe efforts taken to address its presence when found.	10,11,12
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	10,11,12
Additional analyses	16	Describe methods of additional analyses if done, indicating which were pre-specified. This may include, but not be limited to, the following: <ul style="list-style-type: none"> • Sensitivity or subgroup analyses; • Meta-regression analyses; • <i>Alternative formulations of the treatment network; and</i> • <i>Use of alternative prior distributions for Bayesian analyses (if applicable).</i> 	11,12

RESULTS†

Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	11,12
Presentation of network structure	S3	Provide a network graph of the included studies to enable visualization of the geometry of the treatment network.	12,13,Figure2-3
Summary of network geometry	S4	Provide a brief overview of characteristics of the treatment network. This may include commentary on the abundance of trials and randomized patients for the different interventions and pairwise comparisons in the network, gaps of evidence in the treatment network, and potential biases reflected by the network structure.	12,13,
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	11,12, Table1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment.	
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: 1) simple summary data for each intervention group, and 2) effect estimates and confidence intervals. <i>Modified approaches may be needed to deal with information from larger networks.</i>	12,13, Figure2-5
Synthesis of results	21	Present results of each meta-analysis done, including confidence/credible intervals. <i>In larger networks, authors may focus on comparisons versus a particular comparator (e.g. placebo or standard care), with full findings presented in an appendix. League tables and forest plots may be considered to summarize pairwise comparisons.</i> If additional summary measures were explored (such as treatment rankings), these should also be presented.	12,13,Figure4-5, Additional file 1: Table S1-S13
Exploration for inconsistency	S5	Describe results from investigations of inconsistency. This may include such information as measures of model fit to compare consistency and inconsistency models, <i>P</i> values from statistical tests, or summary of inconsistency estimates from different parts of the treatment network.	12,13

Risk of bias across studies	22	Present results of any assessment of risk of bias across studies for the evidence base being studied.	12,13, Additional file 1: Figure S1-S2
Results of additional analyses	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression analyses, <i>alternative network geometries studied, alternative choice of prior distributions for Bayesian analyses</i> , and so forth).	12,13
DISCUSSION			
Summary of evidence	24	Summarize the main findings, including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy-makers).	14-16
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review level (e.g., incomplete retrieval of identified research, reporting bias). <i>Comment on the validity of the assumptions, such as transitivity and consistency. Comment on any concerns regarding network geometry (e.g., avoidance of certain comparisons).</i>	16
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	17
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. This should also include information regarding whether funding has been received from manufacturers of treatments in the network and/or whether some of the authors are content experts with professional conflicts of interest that could affect use of treatments in the network.	17

PICOS = population, intervention, comparators, outcomes, study design.

* Text in italics indicates wording specific to reporting of network meta-analyses that has been added to guidance from the PRISMA statement.

† Authors may wish to plan for use of appendices to present all relevant information in full detail for items in this section.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

For peer review only

BMJ Open

Comparative efficacy of treatment strategies for hepatocellular carcinoma: systematic review and network meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-021269.R1
Article Type:	Research
Date Submitted by the Author:	26-Mar-2018
Complete List of Authors:	<p>Tian, Guo; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China.</p> <p>Yang, Shigui; The State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, School of Medicine, Zhejiang University, The Key Laboratory of Infectious Diseases</p> <p>Yuan, Jinqiu; The Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong, Division of Epidemiology</p> <p>Threapleton, Diane; Chinese University of Hong Kong, School of Public Health & Primary Care</p> <p>Zhao, Qiyu ; Department of Ultrasound Medicine, the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China, Department of Ultrasound Medicine</p> <p>Chen, Fen ; Department of Hepatobiliary Pancreatic Surgery, First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China., Department of Hepatobiliary Pancreatic Surgery</p> <p>Cao, Hongcui ; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University</p> <p>Jiang, Tian'an; First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China., Department of Ultrasonography</p> <p>Li, Lanjuan; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University,</p>
Primary Subject Heading:	Oncology
Secondary Subject Heading:	Palliative care, Radiology and imaging, Surgery
Keywords:	resection, radiofrequency ablation, microwave ablation, transcatheter arterial chemoembolization, percutaneous ethanol injection, hepatocellular carcinoma

SCHOLARONE™
Manuscripts

For peer review only

**Comparative efficacy of treatment strategies for hepatocellular carcinoma:
systematic review and network meta-analysis**

Guo Tian MD ^{1*}, Shigui Yang PhD ^{1*}, Jinqiu Yuan PhD ^{2,3}, Diane Threapleton PhD ²,
Qiyu Zhao PhD ⁴, Fen Chen PhD ⁵, Hongcui Cao PhD ¹, Tian'an Jiang PhD ⁴, Lanjuan
Li PhD ¹

¹State Key Laboratory for Diagnosis and Treatment of Infectious Diseases,
Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases,
The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou
310003, China.

²Division of Epidemiology, The Jockey Club School of Public Health and Primary
Care, The Chinese University of Hong Kong, Hong Kong;

³Shenzhen Municipal Key Laboratory for Health Risk Analysis, Shenzhen Research
Institute of The Chinese University of Hong Kong, Shenzhen, Guangdong Province,
China

⁴Department of Ultrasonography, First Affiliated Hospital, Zhejiang University
School of Medicine, Hangzhou 310003, China.

⁵Department of Hepatobiliary Pancreatic Surgery, First Affiliated Hospital, Zhejiang
University School of Medicine, Hangzhou 310003, China.

Correspondence to: Lanjuan Li, State Key Laboratory for Diagnosis and Treatment of
Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Infectious Diseases, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China (ljli@zju.edu.cn); Or Tian'an Jiang, Department of Ultrasound, First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou 310003, China (tiananjiang@zju.edu.cn); Or Hongcui Cao, State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China (hccao@zju.edu.cn).

*These authors contributed equally to this work.

Electronic word count: 2803 (excluding Tables and References)

Number of figures and tables: 7 figures and 15 tables

List of abbreviations in order of appearance: HCC: hepatocellular carcinoma; RES: resection; RFA: radiofrequency ablation; MWA: microwave ablation; TACE: transcatheter arterial chemoembolization; PEI: percutaneous ethanol injection; GRADE: Grading of Recommendations Assessment, Development and Evaluation; OR: odds ratio; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses; TR: TACE plus RFA; OS: overall survival; MCMC: Markov Chain Monte Carlo; CrI: credible interval; SUCRA: surface under the cumulative ranking curve LPS: lipopolysaccharide; TNF α : tumor necrosis factor α ; IL: interleukin; TGF β : transforming growth factor β .

Conflict of interest: The authors have declared that no competing interests regarding the publication of this paper.

Data sharing statement: Because this is a meta-analysis, it is not available for Patient Consent. All data in this network meta-analysis have been provided in either the main manuscript or additional file.

Financial support: This study was supported by the opening foundation of the State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital of Medical College, Zhejiang University, grant NO. 2015KF06; This study was supported by the Foundation of Zhejiang Health Committee (2017KY346).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Author Contributions:

1. Conceived and designed the experiments: Hongcui Cao, Tian'an Jiang, Lanjuan Li
2. Performed the experiments: Guo Tian, Shigui Yang, Jinqiu Yuan, Diane Threapleton, Qiyu Zhao, Fen Chen, Tian'an Jiang
3. Analyzed the data: Guo Tian, Shigui Yang, Jinqiu Yuan, Qiyu Zhao
4. Contributed reagents/materials/analysis tools: Qiyu Zhao, Fen Chen
5. Wrote the manuscript: Guo Tian, Shigui Yang, Tian'an Jiang
6. Critically revised and approved the final version of manuscript: Diane Threapleton, Hongcui Cao, Tian'an Jiang, Lanjuan Li
7. Study supervision: Hongcui Cao, Tian'an Jiang, Lanjuan Li

Abstract

Objective: Hepatocellular carcinoma (HCC) is the 3rd leading cause of cancer death worldwide. We conducted network meta-regression within a bayesian framework to compare and rank different treatment strategies for HCC through direct and indirect evidence from international studies.

Methods and analyses: We pooled the odds ratio (OR) for 1-, 3- and 5-year overall survival, based on lesions of size < 3 cm, 3-5 cm and ≤ 5 cm, using five therapeutic options including resection (RES), radiofrequency ablation (RFA), microwave ablation (MWA), transcatheter arterial chemoembolization (TACE) plus RFA (TR) and percutaneous ethanol injection (PEI).

Results: We identified 62 studies, including 23893 patients. After adjustment for study design, and in the full sample of studies, the treatments were ranked in order of good to bad as follows for 5-year survival: 1) RES, 2) TR, 3) RFA, 4) MWA, and 5) PEI. The ranks were similar for 1 and 3-year survival, with RES and TR being the highest ranking treatments. In both smaller (< 3 cm) and larger tumors (3-5cm), RES and TR were also the two highest ranking treatments. There was little evidence of inconsistency between direct and indirect evidence.

Conclusion: The comparison of different treatment strategies for HCC indicated that RES is associated with longer survival. However, many of the between-treatment comparisons were not statistically significant and, for now, selection of strategies for treatment will depend patient and disease characteristics. Additionally, much of the evidence was provided by non randomised studies and knowledge gaps still exist.

More head-to-head comparisons between both RES and TR, or other approaches, will be necessary to confirm these findings.

Key words: resection; radiofrequency ablation; microwave ablation; transcatheter arterial chemoembolization; percutaneous ethanol injection; hepatocellular carcinoma.

Strengths and limitations of this study:

1. We conducted network meta-regression within a bayesian framework to compare and rank different treatment strategies for HCC through direct and indirect evidence from international studies.
2. We pooled the odds ratio (OR) for 1-, 3- and 5-year overall survival, based on lesions of size < 3 cm, 3-5 cm and \leq 5 cm, using five therapeutic options including resection (RES), radiofrequency ablation (RFA), microwave ablation (MWA), transcatheter arterial chemoembolization (TACE) plus RFA (TR) and percutaneous ethanol injection (PEI).
3. The comparison of different treatment strategies for HCC indicated that RES is associated with longer survival.
4. A major limitation is in the inclusion of non-randomised studies, in which selection bias is likely to confound observations. Selection of treatment is likely to be based on individual or tumor characteristics, and thus these factors will bias and confound observations of survival.
5. All included studies did not report our primary outcome of interest (5-year survival) and this was a particular limitation among randomised studies.

Introduction

Cancer was the second leading cause of death in 2013, behind cardiovascular disease, and in 2013 more than 8 million people died from cancer globally ¹⁻³. Hepatocellular carcinoma (HCC) is the 6th most common cancer worldwide and the 3rd leading cause of cancer death, with 5-year overall survival rates under 12% ^{4,5}.

Hepatic resection (RES) is the traditional choice for patients with HCC, without cirrhosis and with good remaining liver function ⁶. Despite nearly 70% 5-year survival, recurrence rates with surgery are high ⁷. Repeated hepatectomies to lengthen survival are not often appropriate owing to multiple-site tumor recurrence or patient background of liver cirrhosis ^{8,9}. Many locoregional therapies have been developed including ablative treatments such as percutaneous ethanol injection (PEI), radiofrequency ablation (RFA), or microwave ablation (MWA), and trans-arterial therapies such as transcatheter arterial chemoembolization (TACE) or transarterial chemotherapy infusion (TACI). Locoregional therapies are minimally invasive and therefore are cheaper and faster to recover from, as compared to resection. Such approaches may be appropriate for patients with unresectable, small or multiple carcinomas or those with severe cirrhosis. However, there may be a greater risk of recurrence because of incomplete destruction of cancer cells at the treatment margin, as seen with RFA ¹⁰.

Selection of treatment strategy is determined by liver function, tumor stage and patient performance status ⁷, but much uncertainty still remains surrounding the comparative efficacy of different treatment approaches. A recent review of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

international guidelines for HCC found similarities but also some discrepancy in treatment allocation recommendations because of regional classification differences, secondary to a lack of solid or high-level evidence ¹¹. A recent review of therapies also revealed that there was no consensus on whether surgery or ablation was better for small tumors ⁷. Some discrepancy in prevalence and treatment outcomes may remain in different regions because of local biology, available resources or expertise and access to care ¹¹. However, if we ever hope to achieve standardized and evidence-based therapy for HCC, the unanswered question surrounding relative treatment efficacy of RES compared to ablative locoregional therapies must be resolved.

Traditional meta-analysis is limited by existing head-to-head treatment comparisons within included studies. It is therefore not possible to gauge the relative benefit of two treatments that have never been directly compared in studies. Real-life treatment-decisions are hindered by gaps in existing evidence, but network meta-analysis enables integration of direct and indirect comparisons to provide estimates for relative comparisons across many treatments ¹². In order to investigate comparative effectiveness among RES and common locoregional ablative therapies, we performed a systematic review and network meta-analysis.

Search Strategy

We conducted a systematic review and report findings in accordance with PRISMA for Network Meta-Analyses (PRISMA-NMA) ¹³ (PRISMA NMA Checklist).

The following databases were searched: PubMed, Embase, Web of science and Scopus, up to December 2015, using these keywords: resection, surgery, hepatectomy, radiofrequency ablation, transarterial chemoembolization, microwave thermal ablation, ethanol injection, liver, cancer, tumor (Additional file 1: Text S1). No language restrictions were used. Bibliographies from other relevant review articles were cross-examined for potential missed studies. Disagreement was resolved by a third reviewer. Citations were downloaded into reference management software and duplicate citations were electronically or manually removed.

We systematically included the studies using the following criteria: 1) original data from prospective or retrospective cohort studies and randomized clinical trials (RCTs) in humans; 2) reporting at least two treatments, including resection or any local ablative therapy (RES, RFA, MWA, PEI, or TACE+RFA (TR)); 3) mean lesion size ≤ 5 cm; 4) evaluating overall survival rate not less than one year after first or recurrent treatments. Conference abstracts and case reports were excluded, as were older publications from studies with multiple publications.

Data Extraction and Study Quality

Two investigators independently extracted and cross-checked the data from the eligible studies: author, year, study design, country, disease type, inclusion criteria, treatment style, study size, gender, age, tumor size, follow-up duration, treatment complications and survival outcomes. If in disagreement, a third reviewer adjudicated. The level of evidence was appraised using the Grading of Recommendations

Assessment, Development and Evaluation (GRADE) guidance¹⁴, which was classified into four levels of high, moderate, low, and very low. The quality score was downgraded according to 5 domains, including risk of bias, inconsistency, indirectness, imprecision, and publication bias while scores were upgraded according to large effect, appropriate control for plausible confounding, and dose-response gradient.

Data Analysis

Network meta-analysis was used if a ring or open evidence loop was available. When possible, pair-wise direct head-to-head comparisons were conducted to calculate the pooled odds ratio (OR) and its 95% confidence interval (CI). Between-study heterogeneity was evaluated using the tau-squared statistic (τ^2)¹⁵. A node-splitting analysis was applied to check the consistency between direct evidence (existing real reported comparisons) and indirect evidence (estimated treatment comparisons) for their agreement on a specific node¹⁶. Bayesian network meta-analysis with Markov Chain Monte Carlo (MCMC), through a consistency model, was utilized to estimate the pooled ORs and its 95% credible interval (CrI) for the direct and indirect comparisons¹⁶. The inconsistency model was used to check for heterogeneity due to chance imbalance in the distribution of effect modifiers. Consistency in every closed loop was checked by the loop-specific approach in order to estimate whether treatment survival effects were disturbed by variance in the distribution of potential confounding factors among the studies. In order to compare

and rank survival rates of different treatments we examined all studies first and then separately assessed smaller (<3cm) and larger (3-5cm) tumors. Random-effect meta-regression models were used, with and without adjustment for study design (cohort or RCT) and subgroup analyses were also conducted for RCTs in order to examine treatment effectiveness. We appraised the ranking probabilities for all therapies for each intervention and the treatment hierarchy was ordered by the surface under the cumulative ranking curve (SUCRA)¹⁷. Sensitivity analysis was conducted to remove each study, in turn, and estimate the treatment effect in the remaining studies. Funnel plots were utilized to check the possible presence of publication bias or small-study bias¹⁸. In this study, we used Bayesian MCMC simulations by WinBUGS 1.4 and graphically presented the results using Stata 13.

Results

Study Characteristics

After screening, 62 relevant studies in 61 articles were identified, of which 18 were randomized controlled trials and 44 were cohort studies¹⁹⁻⁸⁰. We excluded 61571 duplicate or non-relevant citations (Figure 1). The summary characteristics of these studies are shown in Additional file 1: Table S1. Overall, 23893 patients of mean age from 46 to 73.5 years, with approximately 29236 tumors, were assigned to receive RES, RFA, MWA, TR and PEI, and the mean follow-up ranged from 1.5 to 5.7 years. In addition, the numbers of connected studies to the lines (black) and sample size of each treatment (red) were shown in Figure 2 and 3, respectively.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Network Meta-Analysis Results

Ten possible treatment comparisons among the five interventions were examined in the included studies. Comparable survival estimates were made for each treatment (per 1000 patients) and the survival OR among each of the treatment comparisons, according to follow-up duration, are presented in Additional file 1: Table S2, along with estimation of the quality of evidence using GRADE criteria.

Across the range of treatment comparisons and follow-up durations, evidence was graded between low and high quality. Evidence was often graded as low quality owing to publication bias and graded as high quality owing to a larger number of participants in direct comparisons.

Survival probabilities (estimated using Meanrank) and ranks for the five treatments in patients with tumors <3cm, 3-5cm or ≤5cm (with and without adjustment for study design) are graphically displayed in Figures 2-5, and numerical details are given in Additional file 1: Table S3-S4. RES was consistently associated with greater survival (rank 1) compared to MWA, RFA, TR and PEI for the 5-year survival estimates. The ranks were similar for 1 and 3-year survival with RES or TR being ranked as 1 or 2 in most analyses. After adjustment for study design, and in the full sample of available studies (n=40), the treatments were ranked as follows for 5-year survival: 1) RES, 2) TR, 3) RFA, 4) MWA, and 5) PEI (Table S4).

Efficacy comparisons from network meta-regression for all treatments are summarized in Table 1 and 2, according to follow-up duration and initial tumor size.

Compared to RES, the 5-year survival in all studies (trials and observational studies) for all tumors ≤ 5 cm, was 0.47 (95%CrI 0.22 to 0.87) for PEI, 0.79 (95%CrI 0.24 to 1.92) for TR, 0.56 (95%CrI 0.23 to 1.14) for MWA and 0.56 (95%CrI 0.27 to 0.99) for RFA (Table 2). When examining the comparisons across all treatments, the only significant difference for tumors < 3 cm was for 5-year survival, and a significantly worse survival was observed for PEI compared to RES 0.46 (95%CrI 0.18 to 0.95). For tumors between 3 and 5 cm, no significant differences were observed at 5-year survival, but significantly worse 3-year survival was observed with PEI, MWA and RFA compared to RES (Table 2). Despite smaller number of studies in analyses of only RCTs, the pairwise comparisons showed similar results. However, all relative rankings should be interpreted with caution because most network meta-regression comparisons did not suggest a statistically significant difference between treatments. Detailed results of each comparison for survival rates are shown in Additional file 1: Table S5-S10.

Loop-specific methods detected no inconsistency between the pairwise and network meta-analysis for most closed loops in the network (Additional file 1: Figure S1). However, inconsistency was observed between direct and indirect comparisons for the following loops: lesions < 3 cm: RES-RFA-TR, PEI-RES-RFA, MWA-RES-RFA; lesions 3-5cm: MWA-RES-RFA, RES-RFA-TR; and lesions ≤ 5 cm: RES-RFA-TR). In addition, tests for inconsistency were carried out (Additional file 1: Table S11-S13), which indicated a close relationship of between-trial heterogeneity and inconsistency between “direct” and “indirect” evidence.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Sensitivity Analysis and publication bias

No significant change was observed when any one study was deleted. Funnel plots indicated that the included studies in each group were distributed symmetrically around the vertical line ($x=0$), suggesting that no obvious evidence of publication bias or small-sample effect existed in this network (Additional file 1: Figure S2).

Discussion

There are many techniques for attaining a large ablated zone and complete necrosis of HCC and this comprehensive review addresses two of the more common treatments, namely resection and ablation. In this network meta-analysis, of the five examined therapies, the pooled data showed RES ranked best in full sample analysis with or without adjustment for study design. In both smaller ($<3\text{cm}$) and larger tumors ($3\text{-}5\text{cm}$) RES remained the highest ranking treatment. However, most of the individual treatment comparisons were not statistically significant and thus, RES may not be superior to all other therapies. Our evidence indicates locoregional therapies and particularly RES or TR (TACE+RFA) are associated with longer survival.

Our observation of better survival outcomes with TR may be through the advantage of dual mechanisms. With TR, TACE induces hypoxic injury on cancer cells through occlusion of blood vessels and is followed by local ablation. This combination therapy may result in a larger ablated zone⁸¹, reduce the possibility of micrometastasis and recurrence, and thus, result in better survival outcomes than RFA alone.

While being more invasive, and despite risk of complications, RES was associated with better survival outcomes after 1 year, 3 years and 5 years. This may be due to removal of larger sections of liver than can be targeted with locoregional therapies, thus removing a larger area of potentially cancerous cells. Additionally, rat models indicate that the liver has the potential to quickly restore its original size after partial hepatectomy. This may be mediated via interactions of lipopolysaccharide (LPS), tumor necrosis factor (TNF) α , interleukin (IL)-6, and transforming growth factor β (TGF β)⁸². However, evidence from rat models and human studies indicates that resection success is associated with resection size and regeneration is stunted with larger resections⁸³⁻⁸⁵. The safe limit for remnant liver volume in normal liver is approximately 30% of total liver volume, but this is estimated to rise to 40-50% in those with liver disease^{83 86}. Liver resection is recognised as the most efficient treatment for HCC but is only applicable for less than 30% of all patients (Morise 2014). However, developments in preoperative imaging techniques, laproscopic surgery and newly developing combinations with chemotherapy may extend its application to more advanced tumors⁸⁶. Furthermore, the consistent associations observed with all studies and only in RCTs indicates that patient selection bias in the observational studies does not wholly explain the better survival outcomes with RES.

Overall, we found PEI was associated with shorter survival than the other four therapies, a finding which is supported in previous studies^{20 29}. One study reported RFA was superior to PEI in achieving short- and long-term survival outcomes, although PEI and RFA showed similar 5-year survival in lesions <3 cm⁵¹. The

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

possible reason why PEI is less effective than RFA may be because lesions often have a thick capsule and therefore ethanol may not distribute through tissues.

There are several limitations in this study. Firstly, a major limitation is in the inclusion of non-randomised studies, in which selection bias is likely to confound observations. Selection of treatment is likely to be based on individual or tumor characteristics, and thus these factors will bias and confound observations of survival. Secondly, this study included both RCTs and observational studies, in which study designs and type of data collection may not be comparable. However, findings were consistent among both study designs. Thirdly, all included studies did not report our primary outcome of interest (5-year survival) and this was a particular limitation among randomised studies. Fourthly, for many individual comparisons, there were either no direct comparisons or comparisons from only a small number of studies. The lack of evidence may increase the risk of bias, which could enlarge or undervalue effect size, and may explain the small inconsistency seen between direct and estimated comparisons. Thus, we should be cautious in interpreting treatment rankings for the different survival times and for different size lesions. While adverse events from treatments may differ (not evaluated in detail in this review), by examining overall survival outcomes in our review, we have taken account of both long-term potential benefits and harms from treatments. The focus of these findings should therefore be on the overall observation that RES or TR may be superior in terms of survival, rather than focusing on specific OR values for individual treatment comparisons.

In conclusion, the findings of the current bayesian network meta-analysis indicate that RES or TR may be among the most effective therapeutic approaches for HCC for 5-year survival in both smaller ($< 3\text{cm}$) and larger ($3\text{-}5\text{cm}$) lesions. However, evidence was of variable quality, and the majority of evidence came from non randomised studies, which are prone to selection bias and knowledge gaps still exist. For not, at the individual level, selection of strategies should depend on patient and clinical characteristics. To facilitate generation of evidence-based recommendations for HCC therapy, and to standardize treatment approaches, further head-to-head comparisons, especially of resection and ablative therapies, are required from high-quality RCTs, with long follow-up for survival outcomes.

Conflict of interests

The authors have declared that no competing interests regarding the publication of this paper.

Data sharing statement

Because this is a meta-analysis, it is not available for Patient Consent. All data in this network meta-analysis have been provided in either the main manuscript or additional file.

Acknowledgments

This study was supported by the opening foundation of the State Key Laboratory

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center
for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital of
Medical College, Zhejiang University, grant NO. 2015KF06; This study was
supported by the Foundation of Zhejiang Health Committee (2017KY346).

For peer review only

References

1. Collaborators GMaCoD. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;385:117-71
2. Lozano R, Naghavi M, Foreman K, *et al.* Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095-128
3. Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: Global Burden of Disease Study. *Lancet* 1997;349:1269-76
4. El-Serag HB. Hepatocellular carcinoma. *The New England journal of medicine* 2011;365:1118-27
5. Ferlay J, Shin HR, Bray F, *et al.* Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *International journal of cancer* 2010;127:2893-917
6. Forner A, Llovet JM, Bruix J. Hepatocellular carcinoma. *Lancet* 2012;379:1245-55
7. Raza A, Sood GK. Hepatocellular carcinoma review: current treatment, and evidence-based medicine. *World journal of gastroenterology* 2014;20:4115-27
8. Kishi Y, Hasegawa K, Sugawara Y, *et al.* Hepatocellular carcinoma: current management and future development-improved outcomes with surgical resection. *International journal of hepatology* 2011;2011:728103
9. Mazzaferro V, Bhoori S, Sposito C, *et al.* Milan criteria in liver transplantation

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

for hepatocellular carcinoma: an evidence-based analysis of 15 years of experience. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society* 2011;17 Suppl 2:S44-57

10. Shan, Xue-Mei, Ding, *et al.* Radiofrequency ablation of hepatocellular carcinoma sized > 3 and ≤ 5 cm: Is ablative margin of more than 1 cm justified? *World journal of gastroenterology* 2011;19:7389-98

11. Yu SJ. A concise review of updated guidelines regarding the management of hepatocellular carcinoma around the world: 2010-2016. *Clinical & Molecular Hepatology* 2016;22:7-17

12. Chaimani A, Higgins JP, Mavridis D, *et al.* Graphical tools for network meta-analysis in STATA. *PloS one* 2013;8:e76654

13. Hutton B, Salanti G, Caldwell DM, *et al.* The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Annals of internal medicine* 2015;162:777-84

14. Puhan MA, Schunemann HJ, Murad MH, *et al.* A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis. *Bmj* 2014;349:g5630

15. Higgins JP, Green S. Cochrane Handbook for Systematic Reviews of Interventions. *Naunyn-Schmiedebergs Archiv für experimentelle Pathologie und Pharmakologie* 2009;2011:S38

16. Dias S, Welton NJ, Caldwell DM, *et al.* Checking consistency in mixed treatment comparison meta-analysis. *Statistics in medicine* 2010;29:932-44
17. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology* 2011;64:163-71
18. Jin ZC, Zhou XH, He J. Statistical methods for dealing with publication bias in meta-analysis. *Statistics in medicine* 2015;34:343-60
19. Zhang Z, Wu M, Chen H, *et al.* [Percutaneous radiofrequency ablation combined with transcatheter arterial chemoembolization for hepatocellular carcinoma]. *Zhonghua wai ke za zhi [Chinese journal of surgery]* 2002;40:826-9
20. Lencioni RA, Allgaier HP, Cioni D, *et al.* Small hepatocellular carcinoma in cirrhosis: randomized comparison of radio-frequency thermal ablation versus percutaneous ethanol injection. *Radiology* 2003;228:235-40
21. Lin SM, Lin CJ, Lin CC, *et al.* Radiofrequency ablation improves prognosis compared with ethanol injection for hepatocellular carcinoma ≤ 4 cm. *Gastroenterology* 2004;127:1714-23
22. Vivarelli M, Guglielmi A, Ruzzenente A, *et al.* Surgical resection versus percutaneous radiofrequency ablation in the treatment of hepatocellular carcinoma on cirrhotic liver. *Annals of surgery* 2004;240:102-7
23. Cho CM, Tak WY, Kweon YO, *et al.* [The comparative results of radiofrequency ablation versus surgical resection for the treatment of

hepatocellular carcinoma]. *The Korean journal of hepatology* 2005;11:59-71

24. Hong SN, Lee SY, Choi MS, *et al.* Comparing the outcomes of radiofrequency ablation and surgery in patients with a single small hepatocellular carcinoma and well-preserved hepatic function. *Journal of clinical gastroenterology* 2005;39:247-52

25. Huang GT, Lee PH, Tsang YM, *et al.* Percutaneous ethanol injection versus surgical resection for the treatment of small hepatocellular carcinoma: a prospective study. *Annals of surgery* 2005;242:36-42

26. Lin SM, Lin CJ, Lin CC, *et al.* Randomised controlled trial comparing percutaneous radiofrequency thermal ablation, percutaneous ethanol injection, and percutaneous acetic acid injection to treat hepatocellular carcinoma of 3 cm or less. *Gut* 2005;54:1151-6

27. Lu MD, Xu HX, Xie XY, *et al.* Percutaneous microwave and radiofrequency ablation for hepatocellular carcinoma: a retrospective comparative study. *Journal of gastroenterology* 2005;40:1054-60

28. Montorsi M, Santambrogio R, Bianchi P, *et al.* Survival and recurrences after hepatic resection or radiofrequency for hepatocellular carcinoma in cirrhotic patients: a multivariate analysis. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2005;9:62-7; discussion 67-8

29. Shiina S, Teratani T, Obi S, *et al.* A randomized controlled trial of radiofrequency ablation with ethanol injection for small hepatocellular

- carcinoma. *Gastroenterology* 2005;129:122-30
30. Chen MS, Li JQ, Zheng Y, *et al.* A prospective randomized trial comparing percutaneous local ablative therapy and partial hepatectomy for small hepatocellular carcinoma. *Annals of surgery* 2006;243:321-8
31. Lu MD, Kuang M, Liang LJ, *et al.* [Surgical resection versus percutaneous thermal ablation for early-stage hepatocellular carcinoma: a randomized clinical trial]. *Zhonghua yi xue za zhi* 2006;86:801-5
32. Cho YB, Lee KU, Suh KS, *et al.* Hepatic resection compared to percutaneous ethanol injection for small hepatocellular carcinoma using propensity score matching. *Journal of gastroenterology and hepatology* 2007;22:1643-9
33. Gao W, Chen MH, Yan K, *et al.* Therapeutic effect of radiofrequency ablation in unsuitable operative small hepatocellular carcinoma. *Chinese Journal of Medical Imaging Technology* 2007
34. Lupo L, Panzera P, Giannelli G, *et al.* Single hepatocellular carcinoma ranging from 3 to 5 cm: radiofrequency ablation or resection? *HPB : the official journal of the International Hepato Pancreato Biliary Association* 2007;9:429-34
35. Zhou T, Qiu YD, Kong WT. Comparing the effect of radiofrequency ablation and surgical resection for the treatment of small hepatocellular. *Journal of Hepatobiliary Surgery* 2007
36. Abu-Hilal M, Primrose JN, Casaril A, *et al.* Surgical resection versus radiofrequency ablation in the treatment of small unifocal hepatocellular

carcinoma. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2008;12:1521-6

37. Brunello F, Veltri A, Carucci P, *et al.* Radiofrequency ablation versus ethanol injection for early hepatocellular carcinoma: A randomized controlled trial. *Scandinavian journal of gastroenterology* 2008;43:727-35

38. Guglielmi A, Ruzzenente A, Valdegamberi A, *et al.* Radiofrequency ablation versus surgical resection for the treatment of hepatocellular carcinoma in cirrhosis. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2008;12:192-8

39. Hiraoka A, Horiike N, Yamashita Y, *et al.* Efficacy of radiofrequency ablation therapy compared to surgical resection in 164 patients in Japan with single hepatocellular carcinoma smaller than 3 cm, along with report of complications. *Hepato-gastroenterology* 2008;55:2171-4

40. Ohmoto K, Yoshioka N, Tomiyama Y, *et al.* Comparison of therapeutic effects between radiofrequency ablation and percutaneous microwave coagulation therapy for small hepatocellular carcinomas. *Journal of gastroenterology and hepatology* 2009;24:223-7

41. Sakaguchi H, Seki S, Tsuji K, *et al.* Endoscopic thermal ablation therapies for hepatocellular carcinoma: a multi-center study. *Hepatology research : the official journal of the Japan Society of Hepatology* 2009;39:47-52

42. Santambrogio R, Opocher E, Zuin M, *et al.* Surgical resection versus laparoscopic radiofrequency ablation in patients with hepatocellular carcinoma

- and Child-Pugh class a liver cirrhosis. *Annals of surgical oncology* 2009;16:3289-98
43. Shibata T, Isoda H, Hirokawa Y, *et al.* Small hepatocellular carcinoma: is radiofrequency ablation combined with transcatheter arterial chemoembolization more effective than radiofrequency ablation alone for treatment? *Radiology* 2009;252:905-13
44. Ueno S, Sakoda M, Kubo F, *et al.* Surgical resection versus radiofrequency ablation for small hepatocellular carcinomas within the Milan criteria. *Journal of hepato-biliary-pancreatic surgery* 2009;16:359-66
45. Xiang-Yang BU, Wang Y, Zhong GE, *et al.* Comparison of radiofrequency ablation and surgical resection for small primary liver carcinoma. *Chinese Archives of General Surgery* 2009
46. Guo WX, Zhai B, Lai EC, *et al.* Percutaneous radiofrequency ablation versus partial hepatectomy for multicentric small hepatocellular carcinomas: a nonrandomized comparative study. *World journal of surgery* 2010;34:2671-6
47. Huang J, Yan L, Cheng Z, *et al.* A randomized trial comparing radiofrequency ablation and surgical resection for HCC conforming to the Milan criteria. *Annals of surgery* 2010;252:903-12
48. Kagawa T, Koizumi J, Kojima S, *et al.* Transcatheter arterial chemoembolization plus radiofrequency ablation therapy for early stage hepatocellular carcinoma: comparison with surgical resection. *Cancer* 2010;116:3638-44

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

49. Morimoto M, Numata K, Kondou M, *et al.* Midterm outcomes in patients with intermediate-sized hepatocellular carcinoma: a randomized controlled trial for determining the efficacy of radiofrequency ablation combined with transcatheter arterial chemoembolization. *Cancer* 2010;116:5452-60

50. Azab M, Zaki S, El-Shetey AG, *et al.* Radiofrequency ablation combined with percutaneous ethanol injection in patients with hepatocellular carcinoma. *Arab journal of gastroenterology : the official publication of the Pan-Arab Association of Gastroenterology* 2011;12:113-8

51. Giorgio A, Di Sarno A, De Stefano G, *et al.* Percutaneous radiofrequency ablation of hepatocellular carcinoma compared to percutaneous ethanol injection in treatment of cirrhotic patients: an Italian randomized controlled trial. *Anticancer research* 2011;31:2291-5

52. Hung HH, Chiou YY, Hsia CY, *et al.* Survival rates are comparable after radiofrequency ablation or surgery in patients with small hepatocellular carcinomas. *Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association* 2011;9:79-86

53. Nishikawa H, Inuzuka T, Takeda H, *et al.* Comparison of percutaneous radiofrequency thermal ablation and surgical resection for small hepatocellular carcinoma. *BMC gastroenterology* 2011;11:143

54. Yun WK, Choi MS, Choi D, *et al.* Superior long-term outcomes after surgery in child-pugh class a patients with single small hepatocellular carcinoma

- compared to radiofrequency ablation. *Hepatology international* 2011;5:722-9
55. Zhang J, Liu HC, Zhou L. The effectiveness of radiofrequency ablation for the treatment of liver cancer. *Journal of Hepatobiliary Surgery* 2011;19:30-33
56. Zhao M, Wang JP, Li W, *et al.* [Comparison of safety and efficacy for transcatheter arterial chemoembolization alone and plus radiofrequency ablation in the treatment of single branch portal vein tumor thrombus of hepatocellular carcinoma and their prognosis factors]. *Zhonghua yi xue za zhi* 2011;91:1167-72
57. Feng K, Yan J, Li X, *et al.* A randomized controlled trial of radiofrequency ablation and surgical resection in the treatment of small hepatocellular carcinoma. *Journal of hepatology* 2012;57:794-802
58. Peng ZW, Lin XJ, Zhang YJ, *et al.* Radiofrequency ablation versus hepatic resection for the treatment of hepatocellular carcinomas 2 cm or smaller: a retrospective comparative study. *Radiology* 2012;262:1022-33
59. Peng ZW, Zhang YJ, Liang HH, *et al.* Recurrent hepatocellular carcinoma treated with sequential transcatheter arterial chemoembolization and RF ablation versus RF ablation alone: a prospective randomized trial. *Radiology* 2012;262:689-700
60. Signoriello S, Annunziata A, Lama N, *et al.* Survival after locoregional treatments for hepatocellular carcinoma: a cohort study in real-world patients. *TheScientificWorldJournal* 2012;2012:564706
61. Wang JH, Wang CC, Hung CH, *et al.* Survival comparison between surgical

resection and radiofrequency ablation for patients in BCLC very early/early stage hepatocellular carcinoma. *Journal of hepatology* 2012;56:412-8

62. Desiderio J, Trastulli S, Pasquale R, *et al.* Could radiofrequency ablation replace liver resection for small hepatocellular carcinoma in patients with compensated cirrhosis? A 5-year follow-up. *Langenbeck's archives of surgery / Deutsche Gesellschaft fur Chirurgie* 2013;398:55-62

63. Ding J, Jing X, Liu J, *et al.* Comparison of two different thermal techniques for the treatment of hepatocellular carcinoma. *European journal of radiology* 2013;82:1379-84

64. Guo WX, Sun JX, Cheng YQ, *et al.* Percutaneous radiofrequency ablation versus partial hepatectomy for small centrally located hepatocellular carcinoma. *World journal of surgery* 2013;37:602-7

65. Hasegawa K, Kokudo N, Makuuchi M, *et al.* Comparison of resection and ablation for hepatocellular carcinoma: a cohort study based on a Japanese nationwide survey. *Journal of hepatology* 2013;58:724-9

66. Iida H, Aihara T, Ikuta S, *et al.* A comparative study of therapeutic effect between laparoscopic microwave coagulation and laparoscopic radiofrequency ablation. *Hepato-gastroenterology* 2013;60:662-5

67. Imai K, Beppu T, Chikamoto A, *et al.* Comparison between hepatic resection and radiofrequency ablation as first-line treatment for solitary small-sized hepatocellular carcinoma of 3 cm or less. *Hepatology research : the official journal of the Japan Society of Hepatology* 2013;43:853-64

68. Kim JW, Shin SS, Kim JK, *et al.* Radiofrequency ablation combined with transcatheter arterial chemoembolization for the treatment of single hepatocellular carcinoma of 2 to 5 cm in diameter: comparison with surgical resection. *Korean journal of radiology* 2013;14:626-35
69. Lai EC, Tang CN. Radiofrequency ablation versus hepatic resection for hepatocellular carcinoma within the Milan criteria--a comparative study. *International journal of surgery* 2013;11:77-80
70. Lin ZZ, Shau WY, Hsu C, *et al.* Radiofrequency ablation is superior to ethanol injection in early-stage hepatocellular carcinoma irrespective of tumor size. *PloS one* 2013;8:e80276
71. Peng ZW, Zhang YJ, Chen MS, *et al.* Radiofrequency ablation with or without transcatheter arterial chemoembolization in the treatment of hepatocellular carcinoma: a prospective randomized trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2013;31:426-32
72. Tohme S, Geller DA, Cardinal JS, *et al.* Radiofrequency ablation compared to resection in early-stage hepatocellular carcinoma. *HPB : the official journal of the International Hepato Pancreato Biliary Association* 2013;15:210-7
73. Wong KM, Yeh ML, Chuang SC, *et al.* Survival comparison between surgical resection and percutaneous radiofrequency ablation for patients in Barcelona Clinic Liver Cancer early stage hepatocellular carcinoma. *Indian journal of gastroenterology : official journal of the Indian Society of Gastroenterology* 2013;32:253-7

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

74. Zhang L, Wang N, Shen Q, *et al.* Therapeutic efficacy of percutaneous radiofrequency ablation versus microwave ablation for hepatocellular carcinoma. *PloS one* 2013;8:e76119

75. Abdelaziz A, Elbaz T, Shousha HI, *et al.* Efficacy and survival analysis of percutaneous radiofrequency versus microwave ablation for hepatocellular carcinoma: an Egyptian multidisciplinary clinic experience. *Surgical endoscopy* 2014;28:3429-34

76. Shi J, Sun Q, Wang Y, *et al.* Comparison of microwave ablation and surgical resection for treatment of hepatocellular carcinomas conforming to Milan criteria. *Journal of gastroenterology and hepatology* 2014;29:1500-7

77. Yang HJ, Lee JH, Lee DH, *et al.* Small single-nodule hepatocellular carcinoma: comparison of transarterial chemoembolization, radiofrequency ablation, and hepatic resection by using inverse probability weighting. *Radiology* 2014;271:909-18

78. Zhang T, Li K, Luo H, *et al.* [Long-term outcomes of percutaneous microwave ablation versus repeat hepatectomy for treatment of late recurrent small hepatocellular carcinoma: a retrospective study]. *Zhonghua yi xue za zhi* 2014;94:2570-2

79. Pompili M, De Matthaeis N, Saviano A, *et al.* Single hepatocellular carcinoma smaller than 2 cm: are ethanol injection and radiofrequency ablation equally effective? *Anticancer research* 2015;35:325-32

80. Xu J, Zhao Y. Comparison of percutaneous microwave ablation and

- laparoscopic resection in the prognosis of liver cancer. *International journal of clinical and experimental pathology* 2015;8:11665-9
81. Goldberg SN, Giron GD, Lukyanov AN, *et al.* Percutaneous tumor ablation: increased necrosis with combined radio-frequency ablation and intravenous liposomal doxorubicin in a rat breast tumor model. *Radiology* 2002;222:797-804
82. Taub R. Liver regeneration: from myth to mechanism. *Nature reviews Molecular cell biology* 2004;5:836-47
83. Martins PN, Theruvath TP, Neuhaus P. Rodent models of partial hepatectomies. *Liver international : official journal of the International Association for the Study of the Liver* 2008;28:3-11
84. Kele PG, De BM, Ej VDJ, *et al.* Early hepatic regeneration index and completeness of regeneration at 6 months after partial hepatectomy. *British Journal of Surgery* 2012;99:1113–19
85. Preziosi ME, Monga SP. Update on the Mechanisms of Liver Regeneration. *Seminars in Liver Disease* 2017;37:141
86. Morise Z, Kawabe N, Tomishige H, *et al.* Recent advances in liver resection for hepatocellular carcinoma. *Frontiers in Surgery* 2014;1:21

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

File legends:

Figure 1 Flow chart of search.

Figure 2 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in RCTs.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments.

- i Lesions < 3 cm.
- ii Lesions 3-5 cm.
- iii Lesions ≤ 5 cm.

Figure 3 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in all studies.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments.

- i Lesions < 3 cm.
- ii Lesions 3-5 cm.
- iii Lesions ≤ 5 cm.

Figure 4 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in RCTs

A Lesions < 3 cm

B Lesions 3-5 cm

C Lesions ≤ 5 cm (full sample).

Figure 5 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in all studies.

A Lesions < 3 cm

B Lesions 3-5 cm

C Lesions ≤ 5 cm (full sample).

Table 1 Odds ratios (95% credible interval) according to network meta-analyses for the survival for all pairwise comparisons in randomized controlled trials.

3cm for 1-year survival					
PEI					
10.75 (0.01-29.11)	TR				
0.08 (0-0.42)	1.42 (0-5.94)	MWA			
0.68 (0.28-1.36)	13.24 (0.02-55.15)	154.8 (1.74-590.10)	RFA		
0.68 (0.19-1.76)	15.61 (0.02-54.78)	161.8 (1.39-581.00)	1.01 (0.40-2.14)	RES	
3cm for 3-year survival					
PEI					
1.29 (0.13-4.99)	TR				
NA	NA	MWA			
0.88 (0.44-1.79)	1.64 (0.20-5.84)	NA	RFA		
0.75 (0.28-1.89)	1.44 (0.14-5.50)	NA	0.86 (0.40-1.68)	RES	
3cm for 5-year survival					
PEI					
NA	TR				
NA	NA	MWA			
0.93 (0.08-3.85)	NA	NA	RFA		
0.49 (0.04-2.02)	NA	NA	0.71 (0.10-2.47)	RES	
3-5cm for 1-year survival					
PEI					
NA	TR				
NA	NA	MWA			
NA	3.40 (0.64-11.93)	NA	RFA		
NA	1.00 (0-5.00)	NA	0.25 (0-1.47)	RES	
3-5cm for 3-year survival					
PEI					
NA	TR				
NA	NA	MWA			
NA	3.98 (0.71-15.22)	NA	RFA		
NA	1.14 (0-6.20)	NA	0.24 (0-1.25)	RES	
3-5cm for 5-year survival					
PEI					
NA	TR				
NA	NA	MWA			
NA	7.64 (0.14-42.49)	NA	RFA		
NA	12.87 (0.02-44.43)	NA	1.05 (0.03-5.33)	RES	

≤5cm for 1-year survival

PEI					
0.26 (0.06-0.69)	TR				
0.24 (0.03-0.81)	1.26 (0.14-4.73)	MWA			
0.65 (0.32-1.14)	3.3 (1.05-8.21)	4.62 (0.85-15.59)	RFA		
0.42 (0.14-0.98)	2.15 (0.49-6.46)	2.75 (0.52-9.18)	0.65 (0.28-1.31)	RES	

≤5cm for 3-year survival

PEI					
0.49 (0.13-1.33)	TR				
1.25 (0.11-5.36)	3.25 (0.24-14.23)	MWA			
0.83 (0.39-1.73)	2.09 (0.81-4.65)	1.71 (0.17-6.61)	RFA		
0.66 (0.23-1.78)	1.69 (0.47-4.87)	1.18 (0.16-4.30)	0.80 (0.36-1.69)	RES	

≤5cm for 5-year survival

PEI					
1.51 (0.02-7.71)	TR				
NA	NA	MWA			
0.90 (0.08-3.65)	3.59 (0.14-18.06)	NA	RFA		
0.49 (0.04-2.03)	2.96 (0.05-14.70)	NA	0.72 (0.11-2.48)	RES	

The reference treatment (1.00) for all comparisons is listed to the right hand side

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection;

Table 2 Odds ratios (95% credible interval) according to network meta-analyses for the survival for all pairwise comparisons in all studies

3cm for 1-year survival					
PEI					
0.56 (0.07-2.13)	TR				
0.55 (0.18-1.29)	1.96 (0.21-7.87)	MWA			
0.69 (0.39-1.13)	2.45 (0.33-8.72)	1.51 (0.60-3.11)	RFA		
0.71 (0.24-1.60)	2.51 (0.26-9.65)	1.55 (0.41-4.10)	1.03 (0.42-2.07)	RES	
3cm for 3-year survival					
PEI					
0.59 (0.15-1.67)	TR				
1.01 (0.45-2.00)	2.35 (0.54-6.80)	MWA			
0.95 (0.59-1.47)	2.21 (0.60-5.76)	1.02 (0.54-1.76)	RFA		
0.80 (0.33-1.68)	1.87 (0.40-5.56)	0.87 (0.31-1.96)	0.85 (0.40-1.62)	RES	
3cm for 5-year survival					
PEI					
1.06 (0.19-3.41)	TR				
0.90 (0.38-1.83)	1.37 (0.23-4.59)	MWA			
0.81 (0.48-1.28)	1.24 (0.25-3.80)	1.00 (0.50-1.77)	RFA		
0.46 (0.18-0.95)	0.72 (0.11-2.48)	0.58 (0.18-1.33)	0.58 (0.24-1.11)	RES	
3-5cm for 1-year survival					
PEI					
0.21 (0.04-0.56)	TR				
0.60 (0.09-1.94)	3.46 (0.57-11.35)	MWA			
0.50 (0.17-1.13)	2.92 (1.14-6.65)	1.25 (0.31-3.46)	RFA		
0.10 (0-0.63)	0.56 (0-3.31)	0.24 (0-1.61)	0.19 (0-1.18)	RES	
3-5cm for 3-year survival					
PEI					
0.30 (0.03-1.06)	TR				
0.90 (0.08-3.36)	3.48 (0.62-11.64)	MWA			
0.57 (0.10-1.83)	2.37 (0.90-5.53)	1.01 (0.25-2.72)	RFA		
0.09 (0-0.44)	0.36 (0.01-1.73)	0.15 (0-0.77)	0.14 (0.01-0.68)	RES	
3-5cm for 5-year survival					
PEI					
6.11 (0-3.02)	TR				
1.88 (0.04-5.54)	13.88 (0.19-50.64)	MWA			
0.79 (0.05-2.64)	7.08 (0.25-26.41)	1.25 (0.18-3.84)	RFA		
1.88 (0.01-3.18)	14.49 (0.05-27.29)	1.79 (0.03-5.39)	0.91 (0.05-4.18)	RES	

≤5cm for 1-year survival

PEI	TR	MWA	RFA	RES
0.30 (0.11-0.63)				
0.91 (0.41-1.79)	3.51 (1.78-8.52)			
0.78 (0.51-1.13)	3.01 (1.33-6.15)	0.95 (0.48-1.67)		
0.61 (0.26-1.25)	2.35 (0.74-5.96)	0.73 (0.28-1.55)	0.78 (0.37-1.49)	

≤5cm for 3-year survival

PEI	TR	MWA	RFA	RES
0.52 (0.25-0.96)				
1.03 (0.56-1.77)	2.16 (0.99-4.16)			
0.92 (0.63-1.32)	1.93 (1.05-3.29)	0.94 (0.58-1.44)		
0.71 (0.37-1.30)	1.50 (0.64-3.08)	0.72 (0.36-1.32)	0.78 (0.44-1.29)	

≤5cm for 5-year survival

PEI	TR	MWA	RFA	RES
0.71 (0.26-1.57)				
0.90 (0.47-1.58)	1.50 (0.52-3.46)			
0.85 (0.57-1.22)	1.42 (0.58-2.96)	1.01 (0.60-1.59)		
0.47 (0.22-0.87)	0.79 (0.24-1.92)	0.56 (0.23-1.14)	0.56 (0.27-0.99)	

The reference treatment (1.00) for all comparisons is listed to the right hand side

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection.




Figure 1 Flow chart of search.
254x190mm (300 x 300 DPI)





Figure 2 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in RCTs. Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments.

i Lesions < 3 cm.
 ii Lesions 3-5 cm.
 iii Lesions ≤ 5 cm.

227x223mm (300 x 300 DPI)



227x227mm (300 x 300 DPI)




Figure 4 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in RCTs
A Lesions < 3 cm
B Lesions 3-5 cm
C Lesions ≤ 5 cm (full sample).

118x117mm (300 x 300 DPI)




Figure 5 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in all studies.
A Lesions < 3 cm
B Lesions 3-5 cm
C Lesions ≤ 5 cm (full sample).

118x117mm (300 x 300 DPI)

Text S1.**Search strategy:**

Pubmed (1950-present)

1. ("TACE" OR "transarterial chemoembolization")
2. ("RFA" OR "radiofrequency ablation" OR "RF ablation" OR "radiofrequency thermal ablation" OR "RTA")
3. (PEI OR "ethanol injection" OR "ethanol ablation" OR "alcohol ablation")
4. ("microwave ablation" OR "microwave thermal ablation" OR MWA)
5. (liver OR hepato*)
6. (neoplas* OR cancer OR tumor OR tumour OR carcinoma OR oncolog*)
7. 1 OR 2 OR 3 OR 4
8. 5 AND 6 AND 7
9. "Ablation Techniques"[Mesh]
10. "Embolization"[Mesh]
11. "Liver Neoplasms"[Mesh]
12. 9 OR 10
13. 12 AND 11
14. 8 OR 13
15. (resection OR surgery OR hepatectomy)
16. (ablation OR injection OR embolization)
17. 5 AND 6 AND 15 AND 16
18. "Hepatectomy"[Mesh]
19. 12 AND 18 AND 11
20. 17 OR 19
21. 14 OR 20

Embase(1980-present)

1. 'TACE':ab,ti
2. 'transarterial chemoembolization':ab,ti
3. 1 OR 2
4. 'rfa':ab,ti

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
5. 'radiofrequency ablation':ab,ti
6. 'rf ablation':ab,ti
7. 'radiofrequency thermal ablation':ab,ti
8. 'rta':ab,ti
9. 4 OR 5 OR 6 OR 7 OR 8
10. 'PEI':ab,ti
11. ' ethanol injection ':ab,ti
12. ' ethanol ablation ':ab,ti
13. ' alcohol ablation ':ab,ti
14. 10 OR 11 OR 12 OR 13
15. ' microwave ablation ':ab,ti
16. ' microwave thermal ablation ':ab,ti
17. ' MWA ':ab,ti
18. 15 OR 16 OR 17
19. ' liver':ab,ti
20. ' hepato*':ab,ti
21. 19 OR 20
22. ' neoplas*':ab,ti
23. ' cancer ':ab,ti
24. ' tumor ':ab,ti
25. ' tumour ':ab,ti
26. ' carcinoma ':ab,ti
27. ' oncolog*':ab,ti
28. 22 OR 23 OR 24 OR 25 OR 26 OR 27
29. 3 OR 9 OR 14 OR 18
30. 21 AND 28 AND 29
31. ' resection':ab,ti
32. ' surgery':ab,ti
33. ' hepatectomy':ab,ti
34. 31 OR 32 OR 33
35. ' ablation':ab,ti

36. 'injection':ab,ti
37. 'embolization':ab,ti
38. 35 OR 36 OR 37
39. 34 AND 38 AND 21 AND 28
40. 30 OR 39

Scoups

1. TITLE-ABS-KEY ("TACE")
2. TITLE-ABS-KEY ("transarterial chemoembolization")
3. 1 OR 2
4. TITLE-ABS-KEY ("RFA")
5. TITLE-ABS-KEY ("radiofrequency ablation")
6. TITLE-ABS-KEY ("RF ablation")
7. TITLE-ABS-KEY ("radiofrequency thermal ablation")
8. TITLE-ABS-KEY ("RTA")
9. 4 OR 5 OR 6 OR 7 OR 8
10. TITLE-ABS-KEY ("PEI")
11. TITLE-ABS-KEY ("ethanol injection")
12. TITLE-ABS-KEY ("ethanol ablation")
13. TITLE-ABS-KEY ("alcohol ablation")
14. 10 OR 11 OR 12 OR 13
15. TITLE-ABS-KEY ("microwave ablation")
16. TITLE-ABS-KEY ("microwave thermal ablation ")
17. TITLE-ABS-KEY ("MWA")
18. 15 OR 16 OR 17
19. TITLE-ABS-KEY ("liver ")
20. TITLE-ABS-KEY ("hepato*")
21. 19 OR 20
22. TITLE-ABS-KEY ("neoplas*")
23. TITLE-ABS-KEY ("cancer")

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
24. TITLE-ABS-KEY ("tumor")
25. TITLE-ABS-KEY ("tumour")
26. TITLE-ABS-KEY ("carcinoma")
27. TITLE-ABS-KEY ("oncolog*")
28. 22 OR 23 OR 24 OR 25 OR 26 OR 27
29. 3 OR 9 OR 14 OR 18
30. 29 AND 21 AND 28
31. TITLE-ABS-KEY ("resection")
32. TITLE-ABS-KEY ("surgery")
33. TITLE-ABS-KEY ("hepatectomy")
34. 31 OR 32 OR 33
35. TITLE-ABS-KEY ("ablation")
36. TITLE-ABS-KEY ("injection")
37. TITLE-ABS-KEY ("embolization")
38. 35 OR 36 OR 37
39. 34 AND 38 AND 21 AND 28
40. 30 OR 39

- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- Web of science
1. TS=(ablation)
2. TS=(embolization)
3. 1 OR 2
4. TS=(hepatectomy)
5. TS=(liver neoplasms)
6. 3 AND 4 AND 5
7. TI=(resection)
8. TI=(surgery)
9. TI=(hepatectomy)
10. 7 OR 8 OR 9
11. TI=(ablation)

12. TI=(injection)
13. TI=(embolization)
14. 11 OR 12 OR 13
15. TI=(liver)
16. TI=(hepato*)
17. 15 OR 16
18. TI=(neoplas*)
19. TI=(cancer)
20. TI=(tumor)
21. TI=(tumour)
22. TI=(carcinoma)
23. TI=(oncolog*)
24. 18 OR 19 OR 20 OR 21 OR 22 OR 23
25. 10 AND 14 AND 17 AND 24
26. 3 AND 5
27. TI=(TACE)
28. TI=("transarterial chemoembolization")
29. 27 OR 28
30. TI=(RFA)
31. TI=("radiofrequency ablation")
32. TI=("RF ablation")
33. TI=("radiofrequency thermal ablation")
34. TI=(RTA)
35. 30 OR 31 OR 32 OR 33 OR 34
36. TI=(PEI)
37. TI=("ethanol injection")
38. TI=("ethanol ablation")
39. TI=("alcohol ablation")
40. 36 OR 37 OR 38 OR 39
41. TI=("microwave ablation")
42. TI=("microwave thermal ablation")

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
43. TI=(MWA)
44. 41 OR 42 OR 43
45. 29 OR 35 OR 40 OR 44
46. 46 AND 17 AND 24
47. 6 OR 25 OR 26 OR 46

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

Table S1.
Summary of the studies included in the network meta-analysis.

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Zhang 2002 ¹⁹	Prospective cohort	China	HCC	0.3-2	RFA	15(15)	13/2	61.8 (38-78)	4.1 (2.4-6.0)	NA	0.80(1y)	0.80(1y)	NA
					TR	15(15)	12/3	57.8 (39-72)	4.6 (2.3-7.1)	NA	1.00(1y)	1.00 (1y)	NA
Lencioni 2003 ²⁰	RCT	Italy	HCC	1.9±0.8	RFA	52(69)	36/16	67±6 (52-78)	2.8±0.6	1.00(1y)	NA	1.00(1y)	15 pain and 10 fever
					PEI	50(73)	30/20	69±7.4 (40-82)	2.8±0.8	0.96(1y)	NA	0.96(1y)	13 pain and 5 fever
Lin 2004 ²¹	RCT	China	HCC	2±0.9	RFA	52(69)	35/17	62±11	2.9±0.8	0.76(3y)	NA	0.35(3y)	1 transient pleural effusion
					PEI	52(67)	34/18	59±10	2.8±0.8	0.66(3y)	NA	0.17(3y)	1 pain
Vivarelli 2004 ²²	Retrospective cohort	Italy	HCC	2.4	RES	79(92)	57/22	65.2±8.2 (43-81)	≤3/3.1-5 (21/58)	0.81(3y)	0.59(3y)	0.65(3y)	NA
					RFA	79(112)	67/12	67.8±8.7 (41-88)	≤3/3.1-5 (22/57)	0.50(3y)	0.25(3y)	0.33(3y)	NA

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Cho 2005 ²³	Retrospective cohort	Korea	HCC	0.1-3	RES	61	48/13	57	3.4±1.0	NA	0.77(3y)	0.77(3y)	2 bleeding, 1 intraabdominal abscess, 1 wound infection
					RFA	99	76/23	58	3.1±0.8	NA	0.80(3y)	0.80(3y)	1 chest wall metastasis, 1 cholecystitis, 1 iatrogenic burn, 1 ileus, 1 hepatic infarction
Huang 2005 ²⁵	RCT	China	HCC	1-4.9	RES	38(42)	27/11	59±11.4	≤2/2.1-3 (24/14)	0.82	NA	0.82	NA
					PEI	38(46)	19/19	63±10.9	≤2/2.1-3 (21/17)	0.45	NA	0.45	NA
Hong 2005 ²⁴	Retrospective cohort	Korea	HCC	2.9(0.4-4.6)	RES	93	69/24	49.2±9.9	2.5±0.8	0.84(3y)	NA	0.84(3y)	NA
					RFA	55	41/14	59.1±9.6	2.4±0.6	0.73(3y)	NA	0.73(3y)	NA
Lin 2005 ²⁶	RCT	China	HCC	2.3±1	RFA	62(78)	40/22	61±10	2.5±1	0.74(3y)	NA	0.74(3y)	2 haemothorax, 1 gastric bleeding and perforation
					PEI	62(76)	39/23	60±8	2.3±0.8	0.60(3y)	NA	0.60(3y)	1 pain
Lu 2005 ²⁷	Retrospective cohort	China	HCC	2.1±1.1	RFA	53(72)	43/10	54.5±11.7 (24-74)	2.6±1.2 (1.0-6.1)	0.38(3y)	NA	0.38(3y)	2 skin burn, 1 puncture wound infection
					MWA	49(98)	44/5	50.1±13.7 (24-74)	2.5±1.2 (0.9-7.2)	0.51(3y)	NA	0.51(3y)	2 puncture wounds, 2 subcapsular hematoma
Montorsi 2005 ²⁸	Prospective cohort	Italy	HCC	2.1	RES	40	33/7	67±9	<5cm	NA	NA	0.73(3y)	NA
					RFA	58	43/15	67±6		NA	NA	0.60(3y)	NA
Shiina 2005 ²⁹	RCT	Japan	HCC	3.1(0.6-4.3)	RFA	118(184)	79/39	≤65/>65 (44/74)	≤2/>2 (45/73)	NA	NA	0.61(3y)	1 transient jaundice, 1 skin burn, 1 hepatic infarction, 3 neoplastic seeding

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					PEI	114(188)	87/27	≤65/>65 (41/73)	≤2/>2 (57/57)	NA	NA	0.45(3y)	1 abscess2 neoplastic seeding
Chen 2006 ³⁰	RCT	China	HCC	2.4±1	RES	90	75/15	49.4±10.9	≤3/3.1-5 (42/48)	0.53	NA	0.53	2 liver failure, 2 gastrointestinal bleeding, 27 ascites
					RFA	71	56/15	51.9±11.2	≤3/3.1-5 (37/34)	0.58	NA	0.58	3 skin burn
Lu 2006 ³¹	RCT	China	Early HCC	1.8	RES	54(56)	37/17	49±14	3.2±1.0	NA	NA	0.86 (3y)	3 wound infection, 1 gastrointestinal bleeding
					RFA	51(57)	42/9	55±13	2.7±1.0	NA	NA	0.87 (3y)	1 peritoneal bleeding, 1 neoplastic seeding
Cho 2007 ³²	Retrospect ive cohort	Korea	HCC	5.7	RES	130(145)	103/27	56.3±8.8	≤2/2.1-3 (43/87)	0.66	NA	0.66	NA
					PEI	249(275)	181/68	57.7±9.7	≤2/2.1-3 (169/80)	0.49	NA	0.49	NA
Gao 2007 ³³	Retrospect ive cohort	China	HCC	4.6	RES	34(37)	28/6	51.5 (38-67)	2.58±0.41	0.76	NA	0.76	12 fever, 5 ascites
					RFA	53(84)	41/12	57.1 (31-81)	2.45±0.37	0.62	NA	0.62	2 bleeding, 1 fistula, 1 wound infection, 6 fever, 9 ascites
Lupo 2007 ³⁴	Retrospect ive cohort	Italy	HCC	2.6	RES	42	33/9	67(28-80)	4.0(3-5)	NA	0.43	0.43	2 urine infection, 1 bilioma, 1 pleural effusion, 1 renal failure, 1 intra-abdominal bleeding
					RFA	60	47/13	68(42-85)	3.65(3-5)	NA	0.32	0.32	2 liver failure, 1 hepatic abscess, 2 pleural effusion, 1 cutaneous metastasis
Zhou 2007 ³⁵	Retrospect ive cohort	China	HCC	0.5-5.9	RES	40(42)	35/5	53±13	≤2/2.1-5 (7/33)	NA	NA	0.75	NA

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	47(54)	37/10	57±14	≤2/2.1-5 (8/39)	NA	NA	0.19	NA
Abu-Hilal 2008 ³⁶	Retrospective cohort	Italy and China	Early HCC	3.6	RES	34	26/8	67	3.8(1.3-5)	NA	0.56	0.56	3 hepatic failure
					RFA	34	27/7	65	3(2-5)	NA	0.56	0.56	1 artero-portal fistula
Brunello 2008 ³⁷	RCT	Italy	Early HCC	2.2	RFA	70(89)	49/20	70.3±8.1	1.27±0.54	0.60(3y)	NA	0.60(3y)	1 haemoperitoneum 1 right haemothorax
					PEI	69(88)	43/27	69.0±7.7	1.27±0.57	0.58(3y)	NA	0.58(3y)	1 haemoperitoneum 1 death
Guglielmi 2008 ³⁸	Retrospective cohort	Italy	HCC	2.3	RES	91(113)	73/18	≤65/>65 (47/44)	≤3/3.1-6 (31/60)	0.55	0.43	0.48	33 postoperative complications
					RFA	109(153)	88/21	≤65/>65 (38/71)	≤3/3.1-6 (32/77)	0.28	0.14	0.20	11 postoperative complications
Hiraoka 2008 ³⁹	Retrospective cohort	Japan	HCC	2.5	RES	59	44/15	62.4±10.6	2.27±0.55	0.59	NA	0.59	1 death, 2 abscess
					RFA	105	76/29	69.4±9.1	1.98±0.52	0.59	NA	0.59	1 biloma, 2 dermatitis
Bu 2009 ⁴⁵	Retrospective cohort	China	HCC	2.9(0.5-6)	RES	42(46)	36/6	53.93±10.74	≤3/3.1-5 (14/28)	0.57	0.46	0.50	1 postoperative hemorrhage, 3 pleural effusions, 2 subdiaphragmatic effusion
					RFA	46(54)	40/6	55.89±7.37	≤3/3.1-5 (20/26)	0.50	0.31	0.37	4 pleural effusions, 1 postoperative hemorrhage, 1 skin burn

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Ohmoto 2009 ⁴⁰	Retrospect ive cohort	Japan	HCC	2.8±2	RFA	34(37)	25/9	67 (44-78)	1.6 (0.7-2.0)	0.71	NA	0.71	2 pain, 4 fever, 1 bile duct injury, 1 pleural effusion, 1 skin burns, 1 vagovagal reflex
					MWA	49(56)	41/8	64 (38-75)	1.7 (0.8-2.0)	0.37	NA	0.37	11 pain, 17 fever, 9 bile duct injury, 8 pleural effusion, 5 ascites, 4 skin burns, 2 vagovagal reflex, 2 abscess, 2 intrapertitoneal bleeding, 1 hepatic infarction, 1 portal thrombus, 1 biliary peritonitis
Sakaguchi 2009 ⁴¹	Retrospect ive cohort	Japan	HCC	0.1-5	Laparosco pic /thorasc opic RFA	249	169/80	65.6±8.9	2.48±0.89	0.57	NA	0.57	1 frequent premature ventricular contractions, 1 liver decompensation
					Laparosco pic /thorasc opic MWA	142	107/35	64.9±7.8	2.28±0.74	0.63	NA	0.63	1 breath holding and incomplete intestinal obstruction, 2 liver decompensation
Santambrogio 2009 ⁴²	Prospectiv e cohort	Italy	HCC	3.2	RES	78	55/23	68±8	2.87±1.21	0.54	NA	0.54	15 extra-hepatic complications
					Laparosco pic RFA	74	59/15	68±7	2.63±1.07	0.41	NA	0.41	14 extra-hepatic complications
Shibata 2009 ⁴³	RCT	Japan	HCC	2.5±1.2	RFA	43(44)	33/10	69.8±8 (44-87)	1.6±0.5 (0.8-2.6)	0.84(3y)	NA	0.84(3y)	1 pseudoaneurysm

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					TR	46(49)	31/15	67.2±8.9 (45-83)	1.7±0.6 (0.9-3.0)	0.85(3y)	NA	0.85(3y)	1 hepatic infarction
Ueno 2009 ⁴⁴	Retrospective cohort	Japan	HCC	3(0.3-7.9)	RES	123(136)	82/41	67(28-85)	2.7±0.1	0.81	0.72	0.80	NA
					RFA	155(209)	100/55	66(40-79)	2.0±0.1	0.38	0.78	0.63	NA
Guo 2010 ⁴⁶	Retrospective cohort	China	HCC	2.5	RES	73(155)	57/16	50.0 (17.0-68.0)	≤3/3.1-5 (30/43)	0.27	0.47	0.44	1 postoperative hemorrhage, 5 abscess, 3 infected ascites, 1 liver failure, 4 pleural effusion
					RFA	86(211)	63/23	52.5 (26.0-80.0)	≤3/3.1-5 (42/44)	0.33	0.16	0.21	1 postoperative hemorrhage, 1 bile leak, 1 abscess, 1 infected ascites, 3 pleural effusion
Huang 2010 ⁴⁷	RCT	China	HCC	3.87	RES	115(144)	85/30	55.91±12.68	≤3/3.1-5 (45/44)	0.82	0.73	0.76	1 hepatic failure, 13 ascites, 5 effusion, 9 bile leakage, 2 postoperative bleeding, 2 gastrointestinal bleeding
					RFA	115(147)	79/36	56.57±14.30	≤3/3.1-5 (57/27)	0.61	0.52	0.55	1 gastric perforation, 2 hemorrhage, 1 malignant seeding, 1 hepatic infarction
Kagawa 2010 ⁴⁸	Retrospective cohort	Japan	Early HCC	4.2	RES	55(69)	40/15	66.1±8.4	≤2/2.1-5 (9/46)	0.42	NA	0.42	2 deaths, 1 liver failure, 1 pleural effusion, 1 pneumonia, 2 biliary leakage
					TR	62(79)	39/23	67.5±8.4	≤2/2.1-5 (19/43)	0.29	NA	0.29	1 duodenal perforation, 1 hemothorax
Morimoto	RCT	Japan	HCC	2.7	RFA	18(25)	12/6	73 (48-84)	3.7±0.6	NA	0.78(3y)	0.78(3y)	5 pain, 2 pleural effusion

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
2010 ⁴⁹					TR	19(21)	15/4	70 (57-78)	3.6±0.7	NA	0.95(3y)	0.95(3y)	1 pain, 1 pleural effusion
Azab 2011 ⁵⁰	RCT	Egypt	HCC	1.5	RFA	30(33)	75/15	46-77	<5cm	NA	NA	0.90	5 superficial burn, 17 transient pain, 3 portal vein thrombosis, 7 fever, 1 ascites
					PEI	30(32)				NA	NA	0.83	2 portal vein thrombosis, 3 fever, 3 ascites
Giorgio 2011 ⁵¹	RCT	Italy	HCC	1.8	RFA	142	105/37	70±2 (68-74)	2.34±0.45 (1.1-3)	0.70	NA	0.70	1 major complication
					PEI	143	102/41	72±6 (68-79)	2.27±0.48 (1.3-2.9)	0.68	NA	0.68	3 major complication
Hung 2011 ⁵²	Retrospective cohort	China	Early HCC	3.5±2	RES	229	184/45	60.07±12.56	2.88±1.06	0.77	NA	0.77	NA
					RFA	190	121/69	67.42±11.45	2.37±0.92	0.67	NA	0.67	NA
Nishikawa 2011 ⁵³	Retrospective cohort	Japan	HCC	3.3	RES	69	50/19	67.4±9.7	2.68±0.49	0.74	NA	0.74	2 bile leakage, 2 ascites, 1 acute respiratory distress syndrome, 1 gastrointestinal bleeding
					RFA	162	95/67	68.4±8.7	1.99±0.62	0.63	NA	0.63	1 biloma, 1 ascites, 1 intra-abdominal bleeding
Yun 2011 ⁵⁴	Retrospective cohort	Korea	HCC	3.5(0.1-9.1)	RES	215	171/44	51.7±9.7	2.1±0.5	0.94	NA	0.94	NA
					RFA	255	197/58	57.0±9.9	2.1±0.5	0.87	NA	0.87	NA

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Zhang 2011 ⁵⁵	Retrospective cohort	China	HCC	0.5-3.5	RES	103(117)	78/25	56.4±15.2	<5cm	NA	NA	0.35(3y)	12 wound infection, 5 postoperative hemorrhage, 2 hepatic failure, 15 pleural effusions, 6 pleural effusions
					RFA	85(106)	62/23	58.5±12.9	<5cm	NA	NA	0.39(3y)	2 gallbladder cardiac reflex, 4 postoperative hemorrhage, 3 pleural effusions
Feng 2012 ⁵⁷	RCT	China	HCC	3	RES	84(116)	75/9	47 (18-76)	2.6±0.8	0.62(3y)	NA	0.62(3y)	7 pleural effusion, 3 pneumonia, 1 effusion plus infection, 3 wound infection or dehiscence, 1 biliary fistula, 2 abdominal bleeding, 1 pneumothorax or hemothorax
					RFA	84(120)	79/5	51 (24-83)	2.4±0.6	0.55(3y)	NA	0.55(3y)	5 pleural effusion, 1 liver abscess, 2 abdominal bleeding
Peng 2012 ⁵⁸	Retrospective cohort	China	Recurrent HCC	4.9	RES	74	65/9	51.5±12.1 (24-75)	1.1±0.5 (0.8-2.0)	0.62	NA	0.62	1 liver failure, 2 gastrointestinal bleeding, 1 peritoneal bleeding, 1 intestinal obstruction, 1 spontaneous bacterial peritonitis, 1 persistent jaundice, 31 ascites
					RFA	71	63/8	53.1±12.1 (28-74)	1.2±0.6 (0.9-2.0)	0.72	NA	0.72	1 gastrointestinal bleeding, 1 persistent jaundice, 12 ascites

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Peng 2012 ⁵⁹	RCT	China	Recurrent HCC	3.3±1.8	RFA	70(76)	55/15	55.1±9.5 (22-75)	≤3/3.1-5 (46/24)	NA	0.17	0.36	1 persistent jaundice, 1 ascites, 22 fever, 45 pain, 4 vomiting
					TR	69(74)	59/9	57.5±10.0 (19-75)	≤3/3.1-5 (41/28)	NA	0.39	0.46	1 liver failure, 1 ascites, 27 fever, 50 pain, 42 vomiting
Signoriello 2012 ⁶⁰	Retrospective cohort	Italy	HCC	0.1-9	RES	34(44)	30/4	62±7	≤3/3.1-5/>5.1 (13/9/4)	NA	NA	0.29	NA
					RFA	50(74)	40/10	68±7	≤3/3.1-5/>5.1 (24/11/7)	NA	NA	0.15	NA
					PEI	256(349)	188/68	67±8	≤3/3.1-5/>5.1 (143/43/12)	NA	NA	0.20	NA
a. Wang 2012 ⁶¹	Retrospective cohort	China	Early HCC	2.5	RES	52	38/14	≤60 (35)	NA	NA	NA	0.92	NA
					RFA	91	60/31	≤60 (40)		NA	NA	0.73	NA
b. Wang 2012 ⁶²	Retrospective cohort	China	Early HCC	2.5	RES	208	168/40	≤60 (113)	≤2/2.1-5 (6/202)	NA	NA	0.77	NA
					RFA	254	161/93	≤60 (85)	≤2/2.1-5 (60/194)	NA	NA	0.57	NA
Desiderio 2013 ⁶²	Retrospective cohort	Italy	HCC	4.3(2.3-5)	RES	52(94)	37/15	65.6±4.8	≤3	0.46	NA	0.46	2 hepatic failure, 1 biliary fistula, 2 hemoperitoneum, 9 ascites
					RFA	44(81)	35/9	64.4±6.5		0.36	NA	0.36	6 pain, 7 fever
Ding 2013 ⁶³	Retrospective cohort	China	HCC	2.3±1.3	RFA	85(98)	68/17	58.64±8.52 (40-77)	2.38±0.81 (1.0-4.8)	0.82(3y)	NA	0.82(3y)	1 frequent premature ventricular contractions, 1 liver decompensation

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					MWA	113(131)	85/28	59.06±11.68 (30-86)	2.55±0.89 (0.8-5.0)	0.78(3y)	NA	0.78(3y)	1 breath holding and incomplete intestinal obstruction, 2 liver decompensation
Guo 2013 ⁶⁴	Retrospective cohort	China	HCC	2.7	RES	102(129)	94/8	51.5(18-75)	≤3/3.1-5 (75/27)	NA	NA	0.63	5 postoperative hemorrhage, 3 bile leak, 4 abscess, 3 infected ascites, 1 liver failure, 4 pleural effusion
					RFA	94(125)	78/16	56(19-75)	≤3/3.1-5 (62/32)	NA	NA	0.50	1 postoperative hemorrhage, 2 bile leak, 1 abscess, 1 infected ascites, 3 pleural effusion
Hasegawa 2013 ⁶⁵	Retrospective cohort	Japan	HCC	2.2	RES	5361(6461)	3967/1394	66 (48-77)	2.3 (1.2-3)	0.71	NA	0.71	NA
					RFA	5548(7412)	3569/1979	69 (52-80)	2 (1-3)	0.61	NA	0.61	NA
					PEI	2059(2836)	1303/756	69 (52-80)	1.7 (1-3)	0.56	NA	0.56	NA
Iida 2013 ⁶⁶	Retrospective cohort	Japan	HCC	0.1-7.5	Laparoscopic RFA	18(27)	NA	73.5±4.0	2.1±0.5	0.78	NA	0.78	1 abscess
					Laparoscopic MWA	40(56)		70.1±6.6	2.0±0.9	0.78	NA	0.78	1 abscess
Imai 2013 ⁶⁷	Retrospective cohort	Japan	HCC	4.1	RES	101	75/26	63.3±9.7	2.14±0.55	0.87	NA	0.87	NA
					RFA	82	46/36	67.6±8.5	1.87±0.50	0.60	NA	0.60	NA

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Kim 2013 ⁶⁸	Retrospect ive cohort	Korea	Early HCC	0.1-4.2	RES	47	36/11	58.8±10.7	3.66±0.76	NA	0.85(3y)	0.85(3y)	2 pleural effusion, 2 pneumonia, 1 hepatic failure, 1 hepatic abscess, 1 mechanical ileus
					TR	37	31/6	61.7±11.1	3.46±0.75	NA	0.78(3y)	0.78(3y)	1 bile duct dilatation
Lai 2013 ⁶⁹	Retrospect ive cohort	China	HCC	2.9±1.5	RES	80	55/25	60.8±9.9	2.9±1.1	0.71	NA	0.71	NA
					RFA	31	19/12	63.1±12.8	1.8±0.6	0.84	NA	0.84	NA
Lin 2013 ⁷⁰	Retrospect ive cohort	China	Early HCC	3.4	RFA	658	393/265	64.7±10.5	2.4±1.1 (0.8-9.5)	0.60	0.50	0.55	NA
					PEI	378	243/135	63.5±12.1	2.0±0.9 (0.4-7.0)	0.50	0.28	0.40	NA
Peng 2013 ⁷¹	RCT	China	HCC	0.6-5.2	RFA	95(133)	71/24	55.3±13.3	3.39±1.35	NA	0.59(3y)	0.59(3y)	51 pain, 26 fever, 29 vomiting, 4 ascites, 2 pleural effusion, 1 skin burn, 1 abdominal infection, 1 small intestinal obstruction
					TR	94(137)	75/19	53.3±11	3.47±1.44	NA	0.67(3y)	0.67(3y)	57 pain, 33 fever, 40 vomiting, 5 ascites, 3 pleural effusion, 1 skin burn, 1 bile duct stenosis, 1 gastric hemorrhage
Tohme 2013 ⁷²	Retrospect ive cohort	Ameri ca	Early HCC	2.4	RES	50(62)	31/19	66.3±1	3.07±1.17	0.48	NA	0.48	3 pleural effusion, 1 pneumonia, 1 myocardial infarction, 2 biloma, 2 ileus, 1 ascites, 1 hyperbilirubinaemia >6, 1 renal insufficiency, 2 encephalopathy

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	60(75)	38/22	65.6±12	2.36±0.94	0.35	NA	0.35	1 oesophagitis, 3 encephalopathy, 1 cholangitis, 2 ascites, 1 renal insufficiency, 1 pneumonia
Wong 2013 ⁷³	Retrospective cohort	China	Early HCC	0.1-5	RES	46	30/16	55.1±12	2.1±0.6	0.85	NA	0.85	2 fever, 1 increased serum alanine aminotransferase level, 2 atelectasis, 2 biloma
					RFA	36	18/18	63.5±13	1.9±0.6	0.72	NA	0.72	None
Zhang 2013 ⁷⁴	Retrospective cohort	China	HCC	2.2±1	RFA	78(97)	64/14	54±10.5 (30-80)	≤3/3.1-5 (47/31)	0.43	0.39	0.41	1 persistent jaundice, 1 biliary fistula
					MWA	77(105)	67/10	54±9.5 (26-76)	≤3/3.1-5 (36/41)	0.58	0.29	0.39	1 hemothorax and intrahepatic hematoma, 1 peritoneal hemorrhage
Abdelaziz 2014 ⁷⁵	RCT	Egypt	Early HCC	2.3	RFA	45(52)	31/14	56.8±7.3	2.95±1.03	0.68(1y)	NA	0.68(1y)	2 subcapsular hematoma, 1 thigh burn, 2 pleural effusion
					MWA	66(76)	48/18	53.6±5	2.9±0.97	0.96(1y)	NA	0.96(1y)	1 subcapsular hematoma, 1 abdominal wall skin burn
Shi 2014 ⁷⁶	Retrospective cohort	China	HCC	3.8	RES	107(126)	87/20	54.5±9.9	≤3/3.1-5 (37/54)	0.73	0.57	0.60	NA
					MWA	117(143)	93/24	56.6±9.2	≤3/3.1-5 (40/56)	0.65	0.52	0.52	NA

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Yang 2014 ⁷⁷	Retrospect ive cohort	Korea	HCC	0.1-7	RES	52	38/14	55.7±10.6	≤2/2.1-5 (21/31)	0.94	NA	0.94	2 pneumonia, 1 wound infection, 1 biliary anastomotic leak, 1 portal vein thrombosis, 1 nausea, 1 delirium, 4 ascites
					RFA	79	59/20	57.2±9.2	≤2/2.1-5 (36/43)	0.86	NA	0.86	1 vomiting, 1 ascites, 6 abdominal pain, 2 nausea, 1 sinus bradycardia
Zhang 2014 ⁷⁸	Retrospect ive cohort	China	Recurrent HCC	2.7	RES	27(29)	25/2	47±13	3.2±1.0	NA	NA	0.63	NA
					MWA	39(46)	37/2	52±13	2.7±1.1	NA	NA	0.62	NA
Pompili 2015 ⁷⁹	Retrospect ive cohort	Italy	Early HCC	2.8	RFA	136	75/61	68 (41-85)	1.8 (1-2)	0.63	NA	0.63	2 ascites, 1 pleural effusion, 1 hemobilia
					PEI	108	90/18	68.5 (34-86)	1.95 (0.8-2)	0.65	NA	0.65	1 hemobilia, 1 portal vein thrombosis
Xu 2015 ⁸⁰	RCT	China	HCC	0.1-3	Laparoscopic RES	45	34/11	58.3±3.1 (26-78)	3.6±0.7 (1-5)	NA	0.38(3y)	0.38(3y)	3 bile leakage, 3 pleural effusion, 2 postoperative hemorrhage
					MWA	45	32/13	57.9±3.4 (27-76)	3.8±0.9 (2-5)	NA	0.33(3y)	0.33(3y)	1 bile leakage, 1 pleural effusion, 1 postoperative hemorrhage

HCC: hepatocellular carcinoma;

BCLC: Barcelona Clinic Liver Cancer;
 RES: resection;
 RFA: radiofrequency ablation;
 MWA: microwave ablation;
 TR: transcatheter arterial chemoembolization and radiofrequency ablation;
 PEI: percutaneous ethanol injection;
 RCT: randomized controlled trial;
 NA: not available.

Table S2.
Quality assessment of included studies using GRADE framework.

Intervention/Comparator	Illustrative comparative risks* (per 1000, 95% CI)			Relative effect of survival time (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Comparator Assumed survival risk	Corresponding survival risk with intervention				
1-year OS rate						
RES/MWA	923	984 (932 to 997)		OR 5.25 (1.15 to 23.97)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
RFA/MWA	947	944 (902 to 968)		OR 0.94 (0.52 to 1.71)	990 (6 studies)	⊕ ⊕ ⊖ ⊖ low
RES/PEI	835	802 (674 to 889)		OR 0.80 (0.41 to 1.58)	519 (3 studies)	⊕ ⊕ ⊖ ⊖ low
RFA/PEI	944	963 (906 to 1000)		OR 1.02 (0.96 to 1.09)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
RES/RFA	932	945 (931 to 956)		OR 1.25 (0.99 to 1.60)	5006 (30 studies)	⊕ ⊕ ⊕ ⊕ high

RES/TR	939	904 (765 to 965)	OR 0.61 (0.21 to 1.79)	201 (2 studies)	⊕ ⊕ ⊖ ⊖ low
2					
3					
REA/TR	938	802 (310 to 978)	OR 0.27 (0.03 to 2.90)	31 (1 study)	⊕ ⊕ ⊖ ⊖ low
4					
5					
6					
3-7 year OS rate					
7					
RES/MWA	712	734 (623 to 822)	OR 1.12 (0.67 to 1.87)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
8					
9					
10					
11					
REA/MWA	736	779 (717 to 828)	OR 1.26 (0.91 to 1.73)	987 (6 studies)	⊕ ⊕ ⊖ ⊖ low
12					
13					
14					
RES/PEI	499	536 (421 to 645)	OR 1.16 (0.73 to 1.83)	519 (3 studies)	⊕ ⊕ ⊖ ⊖ low
15					
16					
17					
REA/PEI	729	748 (657 to 822)	OR 1.10 (0.71 to 1.71)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
18					
19					
20					
RES/RFA	785	851 (823 to 875)	OR 1.57 (1.28 to 1.93)	15906 (30 studies)	⊕ ⊕ ⊕ ⊖ moderate
21					
22					
23					
RES/TR	798	760 (618 to 860)	OR 0.80 (0.41 to 1.55)	201 (2 studies)	⊕ ⊕ ⊖ ⊖ low
24					
25					
26					
27					
REA/TR	737	611 (516 to 704)	OR 0.56 (0.38 to 0.85)	454 (4 studies)	⊕ ⊕ ⊕ ⊖ moderate
28					
29					
30					
5-year OS rate					
31					
RES/MWA	545	607 (492 to 712)	OR 1.29 (0.81 to 2.07)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
32					
33					
34					
REA/MWA	545	609 (442 to 756)	OR 1.30 (0.66 to 2.58)	687 (4 studies)	⊕ ⊕ ⊖ ⊖ low
35					
36					
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					

RES/PEI	293	436 (334 to 545)	OR 1.87 (1.21 to 2.90)	519 (3 studies)	⊕ ⊕ ⊕ ⊖ moderate
RFA/PEI	533	496 (368 to 624)	OR 0.86 (0.51 to 1.45)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
RES/RFA	601	744 (705 to 779)	OR 1.93 (1.59 to 2.34)	15154 (25 studies)	⊕ ⊕ ⊕ ⊖ moderate
RES/TR	290	419 (251 to 607)	OR 1.76 (0.82 to 3.78)	117 (1 study)	⊕ ⊕ ⊖ ⊖ low
RFA/TR	464	356 (222 to 523)	OR 0.64 (0.33 to 1.27)	139 (1 study)	⊕ ⊕ ⊕ ⊖ moderate

The absolute and relative risk of survival with treatments*. GRADE: Grading of Recommendations, Assessment, Development and Evaluation. *The results presented in the Table S1 were built around the assumption of a consistent relative effect. The implications of this effect for populations were considered at different baseline risks. Based on the assumed risks, corresponding risks after an intervention were calculated using the meta-analytic risk ratio.

Table S3.

Ranking treatments of 1-, 3-year and 5-year survival rate of the lesions < 3 cm, 3-5 cm and ≤ 5 cm in RCT.

Treatment	1-year			3-year			5-year		
	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank
< 3cm	12			10			4		
RES		2	3.06		1	1.80		1	1.25
RFA		3	3.21		3	2.56		2	2.08
MWA		1	1.14		NA	NA		NA	NA
TR		4	3.22		2	2.38		NA	NA
PEI		5	4.36		4	3.26		3	2.68

3-5cm	4			4			2		
RES		1	1.17		1	1.19		1	1.69
RFA		3	2.88		3	2.91		3	2.60
MWA		NA	NA		NA	NA		NA	NA
TR		2	1.94		2	1.90		2	1.71
PEI		NA	NA		NA	NA		NA	NA
All tumours (\leq 5cm)	18			14			5		
RES		3	2.78		2	2.43		1	1.68
RFA		4	3.91		3	3.52		3	2.75
MWA		1	1.62		4	3.10		NA	NA
TR		2	1.79		1	1.68		2	2.09
PEI		5	4.90		5	4.27		4	3.48

RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection.

Table S4.
Ranking treatments of 1-, 3-year and 5-year survival rate of the lesions < 3 cm, 3-5 cm and \leq 5 cm in all studies.

Treatment	1-year			3-year			5-year		
	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank
< 3cm	44			42			31		
RES		3	3.02		2	2.49		1	1.35
RFA		4	3.16		3	3.44		2	3.03
MWA		2	2.19		4	3.52		4	3.31
TR		1	2.05		1	1.66		3	3.18

PEI	5	4.58	5	3.89	5	4.13
3-5cm	17		16		11	
RES	1	1.23	1	1.10	1	1.93
RFA	4	3.52	3	3.43	3	3.18
MWA	3	3.46	4	3.72	4	3.43
TR	2	1.97	2	2.10	2	1.94
PEI	5	4.82	5	4.66	5	4.53
All tumours (\leq 5cm)	62		57		40	
RES	2	2.34	2	2.18	1	1.32
RFA	3	3.27	3	3.48	3	3.36
MWA	4	3.78	4	3.98	4	3.51
TR	1	1.10	1	1.27	2	2.45
PEI	5	4.52	5	4.10	5	4.36

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection.

Table S5.

Survival rates (1-year, 3-year and 5-year) for small lesion (<3cm) treatment comparisons estimated by direct and network meta-analysis in RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	1.01 (0.40-2.14)	0.98 (0.77-1.26)
MWA vs RES	161.8 (1.39-581.0)	NA
TR vs RES	15.61 (0.02-54.78)	NA
PEI vs RES	0.68 (0.19-1.76)	1.03 (0.54-1.94)

MWA vs RFA	154.8 (1.74-590.1)	1.42 (0.63-3.19)
TR vs RFA	13.24 (0.02-55.15)	1.00 (0.56-1.80)
PEI vs RFA	0.68 (0.28-1.36)	0.97 (0.78-1.19)
TR vs MWA	1.42 (0-5.94)	NA
PEI vs MWA	0.08 (0-0.42)	NA
PEI vs TR	10.75 (0.01-29.11)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.86 (0.40-1.68)	0.92 (0.71-1.19)
MWA vs RES	NA	NA
TR vs RES	1.44 (0.14-5.50)	NA
PEI vs RES	0.75 (0.28-1.89)	1.21 (0.59-2.15)
MWA vs RFA	NA	NA
TR vs RFA	1.64 (0.20-5.84)	1.01 (0.55-1.87)
PEI vs RFA	0.88 (0.44-1.79)	0.91 (0.71-1.17)
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	1.29 (0.13-4.99)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.71 (0.10-2.47)	0.93 (0.62-1.37)
MWA vs RES	NA	NA
TR vs RES	NA	NA
PEI vs RES	0.49 (0.04-2.02)	0.55 (0.26-1.15)
MWA vs RFA	NA	NA
TR vs RFA	NA	NA
PEI vs RFA	0.93 (0.08-3.85)	0.97 (0.66-1.40)
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA

Table S6.
Survival rates (1-year, 3-year and 5-year) for lesion (3-5cm) treatment comparisons estimated by direct and network meta-analysis in

RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.25 (0-1.47)	0.89 (0.45-1.77)
MWA vs RES	NA	NA
TR vs RES	1.00 (0-5.0)	NA
PEI vs RES	NA	NA
MWA vs RFA	NA	NA
TR vs RFA	3.40 (0.64-11.93)	1.10 (0.78-1.55)
PEI vs RFA	NA	NA
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.24 (0-1.25)	0.70 (0.34-1.45)
MWA vs RES	NA	NA
TR vs RES	1.14 (0-6.20)	NA
PEI vs RES	NA	NA
MWA vs RFA	NA	NA
TR vs RFA	3.98 (0.71-15.22)	1.29 (0.87-1.89)
PEI vs RFA	NA	NA
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA
5-year OS rate for treatment vs reference		
RFA vs RES	1.05 (0.03-5.33)	0.71 (0.32-1.57)
MWA vs RES	NA	NA
TR vs RES	12.87 (0.02-44.43)	NA
PEI vs RES	NA	NA

MWA vs RFA	NA	NA
TR vs RFA	7.64 (0.14-42.49)	1.93 (0.53-7.06)
PEI vs RFA	NA	NA
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA

Table S7.
Survival rates (1-year, 3-year and 5-year) for lesion (≤ 5 cm) treatment comparisons estimated by direct and network meta-analysis in RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.65 (0.28-1.31)	0.96 (0.77-1.20)
MWA vs RES	2.75 (0.52-9.18)	0.98 (0.54-1.78)
TR vs RES	2.15 (0.49-6.46)	NA
PEI vs RES	0.42 (0.14-0.98)	1.03 (0.54-1.94)
MWA vs RFA	4.62 (0.85-15.59)	1.42 (0.63-3.19)
TR vs RFA	3.3 (1.05-8.21)	1.09 (0.84-1.43)
PEI vs RFA	0.65 (0.32-1.14)	0.95 (0.80-1.14)
TR vs MWA	1.26 (0.14-4.73)	NA
PEI vs MWA	0.24 (0.03-0.81)	NA
PEI vs TR	0.26 (0.06-0.69)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.80 (0.36-1.69)	0.87 (0.69-1.10)
MWA vs RES	1.18 (0.16-4.30)	0.88 (0.39-1.98)
TR vs RES	1.69 (0.47-4.87)	NA
PEI vs RES	0.66 (0.23-1.78)	1.12 (0.59-2.15)
MWA vs RFA	1.71 (0.17-6.61)	NA
TR vs RFA	2.09 (0.81-4.65)	1.20 (0.90-1.60)

1	PEI vs RFA	0.83 (0.39-1.73)	0.84 (0.66-1.07)
2	TR vs MWA	3.25 (0.24-14.23)	NA
3	PEI vs MWA	1.25 (0.11-5.36)	NA
4	PEI vs TR	0.49 (0.13-1.33)	NA
5	5-year OS rate for treatment vs reference		
6	RFA vs RES	0.72 (0.11-2.48)	0.85 (0.61-1.17)
7	MWA vs RES	NA	NA
8	TR vs RES	2.96 (0.05-14.7)	NA
9	PEI vs RES	0.49 (0.04-2.03)	0.55 (0.26-1.15)
10	MWA vs RFA	NA	NA
11	TR vs RFA	3.59 (0.14-18.06)	1.30 (0.70-2.41)
12	PEI vs RFA	0.90 (0.08-3.65)	0.97 (0.66-1.40)
13	TR vs MWA	NA	NA
14	PEI vs MWA	NA	NA
15	PEI vs TR	1.51 (0.02-7.71)	NA

OR: odds ratio;

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection;

NA: not available.

Table S8.

Survival rates (1-year, 3-year and 5-year) for small lesion (<3cm) treatment comparisons estimated by direct and network meta-analysis in all studies.

Intervention	OR (95% CI)	
	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		

1	RFA vs RES	1.03 (0.42-2.07)	1.00(0.95-1.05)
2	MWA vs RES	1.55 (0.41-4.10)	1.00(0.53-1.89)
3	TR vs RES	2.51 (0.26-9.65)	1.00(0.56-1.80)
4	PEI vs RES	0.71 (0.24-1.60)	1.00 (0.93-1.07)
5	MWA vs RFA	1.51 (0.60-3.11)	1.02 (0.85-1.23)
6	TR vs RFA	2.45 (0.33-8.72)	1.00(0.56-1.80)
7	PEI vs RFA	0.69 (0.39-1.13)	0.99 (0.93-1.06)
8	TR vs MWA	1.96 (0.21-7.87)	NA
9	PEI vs MWA	0.55 (0.18-1.29)	NA
10	PEI vs TR	0.56 (0.07-2.13)	NA
11	3-year OS rate for treatment vs reference		
12	RFA vs RES	0.85 (0.40-1.62)	0.94 (0.90-0.99)
13	MWA vs RES	0.87 (0.31-1.96)	0.96 (0.49-1.87)
14	TR vs RES	1.87 (0.40-5.56)	1.17 (0.67-2.04)
15	PEI vs RES	0.80 (0.33-1.68)	1.00 (0.71-1.40)
16	MWA vs RFA	1.02 (0.54-1.76)	1.00 (0.82-1.22)
17	TR vs RFA	2.21 (0.60-5.76)	1.01 (0.55-1.87)
18	PEI vs RFA	0.95 (0.59-1.47)	0.97 (0.90-1.03)
19	TR vs MWA	2.35 (0.54-6.80)	NA
20	PEI vs MWA	1.01 (0.45-2.00)	NA
21	PEI vs TR	0.59 (0.15-1.67)	NA
22	5-year OS rate for treatment vs reference		
23	RFA vs RES	0.58 (0.24-1.11)	0.86 (0.81-0.90)
24	MWA vs RES	0.58 (0.18-1.33)	0.89 (0.44-1.79)
25	TR vs RES	0.72 (0.11-2.48)	0.69 (0.34-1.42)
26	PEI vs RES	0.46 (0.18-0.95)	0.79 (0.73-0.85)
27	MWA vs RFA	1.00 (0.50-1.77)	1.02 (0.78-1.33)
28	TR vs RFA	1.24 (0.25-3.80)	NA
29	PEI vs RFA	0.81 (0.48-1.28)	0.92 (0.85-0.99)
30	TR vs MWA	1.37 (0.23-4.59)	NA
31	PEI vs MWA	0.90 (0.38-1.83)	NA

PEI vs TR

1.06 (0.19-3.41)

NA

Table S9.

Survival rates (1-year, 3-year and 5-year) for lesion (3-5cm) treatment comparisons estimated by direct and network meta-analysis in all studies.

Intervention	OR (95%CI)	
	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.19 (0-1.18)	0.96 (0.78-1.17)
MWA vs RES	0.24 (0-1.61)	NA
TR vs RES	0.56 (0-3.31)	1.02 (0.55-1.88)
PEI vs RES	0.10 (0-0.63)	NA
MWA vs RFA	1.25 (0.31-3.46)	0.98 (0.49-1.95)
TR vs RFA	2.92 (1.14-6.65)	1.11 (0.80-1.54)
PEI vs RFA	0.50 (0.17-1.13)	0.89 (0.66-1.20)
TR vs MWA	3.46 (0.57-11.35)	NA
PEI vs MWA	0.60 (0.09-1.94)	NA
PEI vs TR	0.21 (0.04-0.56)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.14 (0.01-0.68)	0.78 (0.62-0.98)
MWA vs RES	0.15 (0-0.77)	1.02 (0.57-1.81)
TR vs RES	0.36 (0.01-1.73)	0.92 (0.48-1.75)
PEI vs RES	0.09 (0-0.44)	NA
MWA vs RFA	1.01 (0.25-2.72)	0.60 (0.26-1.36)
TR vs RFA	2.37 (0.90-5.53)	1.29 (0.87-1.89)
PEI vs RFA	0.57 (0.10-1.83)	0.71 (0.50-1.00)
TR vs MWA	3.48 (0.62-11.64)	NA
PEI vs MWA	0.90 (0.08-3.36)	NA
PEI vs TR	0.30 (0.03-1.06)	NA
5-year OS rate for treatment vs reference		

RFA vs RES	0.91 (0.05-4.18)	0.62 (0.45-0.85)
MWA vs RES	1.79 (0.03-5.39)	0.90 (0.48-1.69)
TR vs RES	14.49 (0.05-27.29)	NA
PEI vs RES	1.88 (0.01-3.18)	NA
MWA vs RFA	1.25 (0.18-3.84)	0.57 (0.21-1.51)
TR vs RFA	7.08 (0.25-26.41)	2.36 (0.66-8.37)
PEI vs RFA	0.79 (0.05-2.64)	0.56 (0.37-0.84)
TR vs MWA	13.88 (0.19-50.64)	NA
PEI vs MWA	1.88 (0.04-5.54)	NA
PEI vs TR	6.11 (0-3.02)	NA

Table S10.
Survival rates (1-year, 3-year and 5-year) for lesion (≤ 5cm) treatment comparisons estimated by direct, indirect and network meta-analysis in all studies.

Intervention	OR (95%CI)	
	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.78 (0.37-1.49)	0.99 (0.95-1.04)
MWA vs RES	0.73 (0.28-1.55)	0.95 (0.71-1.27)
TR vs RES	2.35 (0.74-5.96)	1.04 (0.70-1.55)
PEI vs RES	0.61 (0.26-1.25)	1.01 (0.74-1.39)
MWA vs RFA	0.95 (0.48-1.67)	1.01 (0.85-1.21)
TR vs RFA	3.01 (1.33-6.15)	1.10 (0.85-1.43)
PEI vs RFA	0.78 (0.51-1.13)	0.98 (0.93-1.05)
TR vs MWA	3.51 (1.78-8.52)	0.91 (0.70-1.18)
PEI vs MWA	0.91 (0.41-1.79)	NA
PEI vs TR	0.30 (0.11-0.63)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.78 (0.44-1.29)	0.93 (0.89-0.98)
MWA vs RES	0.72 (0.36-1.32)	0.96 (0.69-1.32)

TR vs RES	1.50 (0.64-3.08)	1.06 (0.69-1.61)
PEI vs RES	0.71 (0.37-1.30)	0.93 (0.86-1.00)
MWA vs RFA	0.94 (0.58-1.44)	0.95 (0.78-1.16)
TR vs RFA	1.93 (1.05-3.29)	1.20 (0.90-1.60)
PEI vs RFA	0.92 (0.63-1.32)	0.95 (0.89-1.01)
TR vs MWA	2.16 (0.99-4.16)	NA
PEI vs MWA	1.03 (0.56-1.77)	NA
PEI vs TR	0.52 (0.25-0.96)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.56 (0.27-0.99)	0.84 (0.80-0.89)
MWA vs RES	0.56 (0.23-1.14)	0.90 (0.61-1.31)
TR vs RES	0.79 (0.24-1.92)	0.69 (0.34-1.42)
PEI vs RES	0.47 (0.22-0.87)	0.79 (0.73-0.85)
MWA vs RFA	1.01 (0.60-1.59)	0.97 (0.75-1.25)
TR vs RFA	1.42 (0.58-2.96)	1.30 (0.70-2.41)
PEI vs RFA	0.85 (0.57-1.22)	0.91 (0.84-0.98)
TR vs MWA	1.50 (0.52-3.46)	NA
PEI vs MWA	0.90 (0.47-1.58)	NA
PEI vs TR	0.71 (0.26-1.57)	NA

OR: odds ratio;

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection;

NA: not available.

Table S11.

Posterior summaries from random effects consistency and inconsistency models for small lesion (<3cm) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.55	0.21	(0.15-1.00)	0.38	0.23	(0.02-0.88)
τ	11.06	88.80	(1.00-43.58)	4020	78840	(1.28-2366.00)
resdev	90.04	13.04	(66.16-117.10)	94.65	12.94	(70.06-120.70)
pD	59.96			57.5		
DIC	402.44			404.59		
3-year OS rate for treatment vs reference						
σ	0.59	0.14	(0.34-0.88)	0.6	0.14	(0.36-0.91)
τ	3.74	10.43	(1.29-8.74)	3.29	1.92	(1.21-8.05)
resdev	92.02	14.19	(66.64-122.10)	90.7	13.92	(65.64-120.00)
pD	70.71			71.74		
DIC	517.72			517.43		
5-year OS rate for treatment vs reference						
σ	0.53	0.12	(0.32-0.80)	0.55	0.13	(0.34-0.84)
τ	4.19	2.29	(1.57-9.74)	3.82	2.02	(1.42-8.83)
resdev	63.99	11.47	(43.52-88.24)	63.55	11.37	(43.39-87.90)
pD	54.24			54.99		
DIC	411.73			412.03		

Table S12.
Posterior summaries from random effects consistency and inconsistency models for lesion (3-5cm) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency Model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.28	0.25	(0.01-0.92)	0.38	0.34	(0.02-1.28)
τ	42220	1.30E+06	(1.19-19650.00)	19500.00	720600.00	(0.62-4178.00)

resdev	28.90	6.96	(17.25-44.41)	32.18	7.36	(19.64-48.32)
pD	22.80			24.59		
DIC	152.25			157.31		
3-year OS rate for treatment vs reference						
σ	0.62	0.27	(0.17-1.24)	0.67	0.31	(0.14-1.40)
τ	9.02	65.04	(0.66-35.66)	49.29	1164.00	(0.51-48.58)
resdev	32.36	8.17	(18.39-50.07)	32.62	8.22	(18.52-50.51)
pD	28.02			28.65		
DIC	187.98			188.88		
5-year OS rate for treatment vs reference						
σ	0.80	0.46	(0.14-1.94)	0.60	0.42	(0.04-1.64)
τ	49.88	1159	(0.27-49.16)	5839.00	185600.00	(0.37-748.40)
resdev	22.54	6.73	(11.29-37.43)	22.57	6.519	(11.45-36.90)
pD	20.62			19.84		
DIC	132.23			131.49		

Table S13.

Posterior summaries from random effects consistency and inconsistency models for lesion ($\leq 5\text{cm}$) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency Model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.49	0.13	(0.26-0.77)	0.29	0.14	(0.05-0.58)
τ	5.30	3.72	(1.70-14.33)	83.27	806.8	(2.94-391.70)
resdev	129.2	14.99	(101.40-160)	133.1	14.50	(105.70-162.80)
pD	84.95			78.28		
DIC	606.94			604.11		
3-year OS rate for treatment vs reference						
σ	0.50	0.09	(0.33-0.70)	0.47	0.096	(0.29-0.67)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

τ	4.51	1.83	(2.08-9.02)	5.28	2.59	(2.24-11.80)
resdev	124	15.64	(95.16-156.40)	124.5	15.89	(95.35-157.50)
pD	93.89			93.37		
DIC	723.55			723.53		
5-year OS rate for treatment vs reference						
σ	0.44	0.10	(0.26-0.65)	0.44	0.1	(0.26-0.67)
τ	6.25	3.60	(2.38-14.90)	6.08	4.01	(2.25-14.87)
resdev	86.73	13.53	(62.35-115.40)	85.74	13.55	(61.39-114.40)
pD	67.86			68.84		
DIC	544.41			544.41		

sd: standard deviation;
CI: Credible Interval
 σ : between-trial standard deviation
 τ^2 : between-trial variance
resdev: residual deviance
pD: effective number of parameters
DIC: deviance information criterion

Figure S1.**Results of the consistency test for closed loop at 1-year, 3-year, and 5-year survival rate of the lesions < 3 cm, 3-5 cm and ≤ 5 cm.**

- i Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions < 3 cm
- ii Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions 3-5 cm
- iii Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions ≤ 5 cm

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46





Figure S2.**Assessment of publication bias using funnel plot.**

- i Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions < 3 cm.
- ii Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions 3-5 cm.
- iii Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions ≤ 5 cm



For peer review only

PRISMA NMA Checklist of Items to Include When Reporting A Systematic Review Involving a Network Meta-analysis

Section/Topic	Item #	Checklist Item	Reported on Page #
TITLE			
Title	1	Identify the report as a systematic review <i>incorporating a network meta-analysis (or related form of meta-analysis)</i> .	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: Background: main objectives Methods: data sources; study eligibility criteria, participants, and interventions; study appraisal; and <i>synthesis methods, such as network meta-analysis</i> . Results: number of studies and participants identified; summary estimates with corresponding confidence/credible intervals; <i>treatment rankings may also be discussed. Authors may choose to summarize pairwise comparisons against a chosen treatment included in their analyses for brevity.</i> Discussion/Conclusions: limitations; conclusions and implications of findings. Other: primary source of funding; systematic review registration number with registry name.	5,6
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known, <i>including mention of why a network meta-analysis has been conducted.</i>	7,8
Objectives	4	Provide an explicit statement of questions being addressed, with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	8

METHODS

Protocol and registration	5	Indicate whether a review protocol exists and if and where it can be accessed (e.g., Web address); and, if available, provide registration information, including registration number.	8,9
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale. <i>Clearly describe eligible treatments included in the treatment network, and note whether any have been clustered or merged into the same node (with justification).</i>	9,10
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	9
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	9,10, Figure 1, Additional file 1: Text S1
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	9,10
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	10
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	11
Geometry of the network	S1	Describe methods used to explore the geometry of the treatment network under study and potential biases related to it. This should include how the evidence base has been graphically summarized for presentation, and what characteristics were compiled and used to describe the evidence base to readers.	11
Risk of bias within individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	11,12

Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means). <i>Also describe the use of additional summary measures assessed, such as treatment rankings and surface under the cumulative ranking curve (SUCRA) values, as well as modified approaches used to present summary findings from meta-analyses.</i>	11,12
Planned methods of analysis	14	Describe the methods of handling data and combining results of studies for each network meta-analysis. This should include, but not be limited to: <ul style="list-style-type: none">• <i>Handling of multi-arm trials;</i>• <i>Selection of variance structure;</i>• <i>Selection of prior distributions in Bayesian analyses; and</i>• <i>Assessment of model fit.</i>	11,12
Assessment of Inconsistency	S2	Describe the statistical methods used to evaluate the agreement of direct and indirect evidence in the treatment network(s) studied. Describe efforts taken to address its presence when found.	10,11,12
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	10,11,12
Additional analyses	16	Describe methods of additional analyses if done, indicating which were pre-specified. This may include, but not be limited to, the following: <ul style="list-style-type: none">• Sensitivity or subgroup analyses;• Meta-regression analyses;• <i>Alternative formulations of the treatment network; and</i>• <i>Use of alternative prior distributions for Bayesian analyses (if applicable).</i>	11,12

RESULTS†

Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	11,12
Presentation of network structure	S3	Provide a network graph of the included studies to enable visualization of the geometry of the treatment network.	12,13,Figure2-3
Summary of network geometry	S4	Provide a brief overview of characteristics of the treatment network. This may include commentary on the abundance of trials and randomized patients for the different interventions and pairwise comparisons in the network, gaps of evidence in the treatment network, and potential biases reflected by the network structure.	12,13,
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	11,12, Table1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment.	
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: 1) simple summary data for each intervention group, and 2) effect estimates and confidence intervals. <i>Modified approaches may be needed to deal with information from larger networks.</i>	12,13, Figure2-5
Synthesis of results	21	Present results of each meta-analysis done, including confidence/credible intervals. <i>In larger networks, authors may focus on comparisons versus a particular comparator (e.g. placebo or standard care), with full findings presented in an appendix. League tables and forest plots may be considered to summarize pairwise comparisons.</i> If additional summary measures were explored (such as treatment rankings), these should also be presented.	12,13,Figure4-5, Additional file 1: Table S1-S13
Exploration for inconsistency	S5	Describe results from investigations of inconsistency. This may include such information as measures of model fit to compare consistency and inconsistency models, <i>P</i> values from statistical tests, or summary of inconsistency estimates from different parts of the treatment network.	12,13

Risk of bias across studies	22	Present results of any assessment of risk of bias across studies for the evidence base being studied.	12,13, Additional file 1: Figure S1-S2
Results of additional analyses	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression analyses, <i>alternative network geometries studied, alternative choice of prior distributions for Bayesian analyses</i> , and so forth).	12,13
DISCUSSION			
Summary of evidence	24	Summarize the main findings, including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy-makers).	14-16
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review level (e.g., incomplete retrieval of identified research, reporting bias). <i>Comment on the validity of the assumptions, such as transitivity and consistency. Comment on any concerns regarding network geometry (e.g., avoidance of certain comparisons).</i>	16
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	17
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. This should also include information regarding whether funding has been received from manufacturers of treatments in the network and/or whether some of the authors are content experts with professional conflicts of interest that could affect use of treatments in the network.	17

PICOS = population, intervention, comparators, outcomes, study design.

* Text in italics indicateS wording specific to reporting of network meta-analyses that has been added to guidance from the PRISMA statement.

† Authors may wish to plan for use of appendices to present all relevant information in full detail for items in this section.

For peer review only

BMJ Open

Comparative efficacy of treatment strategies for hepatocellular carcinoma: systematic review and network meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-021269.R2
Article Type:	Research
Date Submitted by the Author:	06-Jun-2018
Complete List of Authors:	<p>Tian, Guo; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China.</p> <p>Yang, Shigui; The State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, School of Medicine, Zhejiang University, The Key Laboratory of Infectious Diseases</p> <p>Yuan, Jinqiu; The Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong, Division of Epidemiology</p> <p>Threapleton, Diane; Chinese University of Hong Kong, School of Public Health & Primary Care</p> <p>Zhao, Qiyu ; Department of Ultrasound Medicine, the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China, Department of Ultrasound Medicine</p> <p>Chen, Fen ; Department of Hepatobiliary Pancreatic Surgery, First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China., Department of Hepatobiliary Pancreatic Surgery</p> <p>Cao, Hongcui ; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University</p> <p>Jiang, Tian'an; First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China., Department of Ultrasonography</p> <p>Li, Lanjuan; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University,</p>
Primary Subject Heading:	Oncology
Secondary Subject Heading:	Palliative care, Radiology and imaging, Surgery
Keywords:	resection, radiofrequency ablation, microwave ablation, transcatheter arterial chemoembolization, percutaneous ethanol injection, hepatocellular carcinoma

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHOLARONE™
Manuscripts

For peer review only

**Comparative efficacy of treatment strategies for hepatocellular carcinoma:
systematic review and network meta-analysis**

Guo Tian MD ^{1*}, Shigui Yang PhD ^{1*}, Jinqiu Yuan PhD ^{2,3}, Diane Threapleton PhD ²,
Qiyu Zhao PhD ⁴, Fen Chen PhD ⁵, Hongcui Cao PhD ¹, Tian'an Jiang PhD ⁴, Lanjuan
Li PhD ¹

¹State Key Laboratory for Diagnosis and Treatment of Infectious Diseases,
Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases,
The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou
310003, China.

²Division of Epidemiology, The Jockey Club School of Public Health and Primary
Care, The Chinese University of Hong Kong, Hong Kong;

³Shenzhen Municipal Key Laboratory for Health Risk Analysis, Shenzhen Research
Institute of The Chinese University of Hong Kong, Shenzhen, Guangdong Province,
China

⁴Department of Ultrasonography, First Affiliated Hospital, Zhejiang University
School of Medicine, Hangzhou 310003, China.

⁵Department of Hepatobiliary Pancreatic Surgery, First Affiliated Hospital, Zhejiang
University School of Medicine, Hangzhou 310003, China.

Correspondence to: Lanjuan Li, State Key Laboratory for Diagnosis and Treatment of
Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Infectious Diseases, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China (ljli@zju.edu.cn); Or Tian'an Jiang, Department of Ultrasound, First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou 310003, China (tiananjiang@zju.edu.cn); Or Hongcui Cao, State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China (hccao@zju.edu.cn).

*These authors contributed equally to this work.

Electronic word count: 2981 (excluding Tables and References)

Number of figures and tables: 7 figures and 15 tables

List of abbreviations in order of appearance: HCC: hepatocellular carcinoma; RES: resection; RFA: radiofrequency ablation; MWA: microwave ablation; TACE: transcatheter arterial chemoembolization; PEI: percutaneous ethanol injection; GRADE: Grading of Recommendations Assessment, Development and Evaluation; OR: odds ratio; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses; TR: TACE plus RFA; OS: overall survival; MCMC: Markov Chain Monte Carlo; CrI: credible interval; SUCRA: surface under the cumulative ranking curve LPS: lipopolysaccharide; TNF α : tumor necrosis factor α ; IL: interleukin; TGF β : transforming growth factor β .

Conflict of interest: The authors have declared that no competing interests regarding the publication of this paper.

Data sharing statement: No additional data are available.

Financial support: This study was supported by the opening foundation of the State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital of Medical College, Zhejiang University, grant NO. 2015KF06; This study was supported by the Foundation of Zhejiang Health Committee (2017KY346).

Author Contributions:

1. Conceived and designed the experiments: Hongcui Cao, Tian'an Jiang, Lanjuan Li

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 2. Performed the experiments: Guo Tian, Shigui Yang, Jinqiu Yuan, Diane Threapleton, Qiyu Zhao, Fen Chen, Tian'an Jiang
- 3. Analyzed the data: Guo Tian, Shigui Yang, Jinqiu Yuan, Qiyu Zhao
- 4. Contributed reagents/materials/analysis tools: Qiyu Zhao, Fen Chen
- 5. Wrote the manuscript: Guo Tian, Shigui Yang, Tian'an Jiang
- 6. Critically revised and approved the final version of manuscript: Diane Threapleton, Hongcui Cao, Tian'an Jiang, Lanjuan Li
- 7. Study supervision: Hongcui Cao, Tian'an Jiang, Lanjuan Li

Abstract

Objective: Hepatocellular carcinoma (HCC) is the 3rd leading cause of cancer death worldwide. We conducted network meta-regression within a bayesian framework to compare and rank different treatment strategies for HCC through direct and indirect evidence from international studies.

Methods and analyses: We pooled the odds ratio (OR) for 1-, 3- and 5-year overall survival, based on lesions of size < 3 cm, 3-5 cm and ≤ 5 cm, using five therapeutic options including resection (RES), radiofrequency ablation (RFA), microwave ablation (MWA), transcatheter arterial chemoembolization (TACE) plus RFA (TR) and percutaneous ethanol injection (PEI).

Results: We identified 74 studies, including 26944 patients. After adjustment for study design, and in the full sample of studies, the treatments were ranked in order of greatest to least benefit as follows for 5-year survival: 1) RES, 2) TR, 3) RFA, 4) MWA, and 5) PEI. The ranks were similar for 1 and 3-year survival, with RES and TR being the highest ranking treatments. In both smaller (<3cm) and larger tumors (3-5cm), RES and TR were also the two highest ranking treatments. There was little evidence of inconsistency between direct and indirect evidence.

Conclusion: The comparison of different treatment strategies for HCC indicated that RES is associated with longer survival. However, many of the between-treatment comparisons were not statistically significant and, for now, selection of strategies for treatment will depend patient and disease characteristics. Additionally, much of the evidence was provided by non randomised studies and knowledge gaps still exist.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

More head-to-head comparisons between both RES and TR, or other approaches, will be necessary to confirm these findings.

Key words: resection; radiofrequency ablation; microwave ablation; transcatheter arterial chemoembolization; percutaneous ethanol injection; hepatocellular carcinoma.

Strengths and limitations of this study:

1. This is a network meta-regression within a bayesian framework to compare and rank different treatment strategies for HCC through direct and indirect evidence from international studies.
2. Strong and reliable methodological and statistical procedures were applied.
3. The individual or tumor characteristics within HCC articles would be a source of heterogeneity..
4. A major limitation is in the inclusion of non-randomised studies, in which selection bias is likely to confound observations. Selection of treatment is likely to be based on individual or tumor characteristics, and thus these factors will bias and confound observations of survival.
5. Other studies did not report the primary outcome of interest (5-year survival) and this was a particular limitation among randomised studies.

Introduction

Cancer was the second leading cause of death in 2013, behind cardiovascular disease, and in 2013 more than 8 million people died from cancer globally ¹⁻³. Hepatocellular carcinoma (HCC) was the 6th most common cancer worldwide and the 3rd leading cause of cancer death, with 5-year overall survival rates under 12% ^{4,5}.

Hepatic resection (RES) was the traditional choice for patients with HCC, without cirrhosis and with good remaining liver function ⁶. Despite nearly 70% 5-year survival, recurrence rates after surgery were high ⁷. Repeated hepatectomies to lengthen survival were not often appropriate owing to multiple-site tumor recurrence or patient background of liver cirrhosis ^{8,9}. Many locoregional therapies have been developed including ablative treatments such as percutaneous ethanol injection (PEI), radiofrequency ablation (RFA), or microwave ablation (MWA), and trans-arterial therapies such as transcatheter arterial chemoembolization (TACE) or transarterial chemotherapy infusion (TACI). Locoregional therapies were minimally invasive and therefore are cheaper and faster to recover, as compared to resection. Such approaches may be appropriate for patients with unresectable, small or multiple carcinomas or those with severe cirrhosis. However, there may be a greater risk of recurrence because of incomplete destruction of cancer cells at the treatment margin, as seen with RFA ¹⁰.

Selection of treatment strategy was determined by liver function, tumor stage and patient performance status ⁷, but much uncertainty still remains surrounding the comparative efficacy of different treatment approaches. A recent review of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

international guidelines for HCC found similarities but also some discrepancy in treatment allocation recommendations because of regional classification differences, secondary to a lack of solid or high-level evidence ¹¹. A recent review of therapies also revealed that there was no consensus on whether surgery or ablation was better for small tumors ⁷. Some discrepancy in prevalence and treatment outcomes may be still in different regions because of local biology, available resources or expertise and access to care ¹¹. However, if we ever hope to achieve standardized and evidence-based therapy for HCC, the unanswered question surrounding relative treatment efficacy of RES compared to ablative locoregional therapies should be resolved.

Traditional meta-analysis is limited by existing head-to-head treatment comparisons within included studies. It is therefore not possible to gauge the relative benefit of two treatments that have never been directly compared in studies. Real-life treatment decisions are hindered by gaps in existing evidence, but network meta-analysis enables integration of direct and indirect comparisons to provide estimates for relative comparisons across many treatments ¹². In this study, we included the latest literature, and focused on the comparison of interventional and surgical treatments, including RES, RFA, MWA, and TACE plus RFA (TR), PEI. In order to investigate comparative effectiveness among RES and common locoregional ablative therapies, we performed a systematic review and network meta-analysis.

Search Strategy

We conducted a systematic review and report findings in accordance with PRISMA for Network Meta-Analyses (PRISMA-NMA)¹³ (Additional file 1: Text S1). The following databases were searched: PubMed, Embase, Web of science and Scopus, up to May 2018, using these keywords: resection, surgery, hepatectomy, radiofrequency ablation, transarterial chemoembolization, microwave thermal ablation, ethanol injection, liver, cancer, tumor (Additional file 1: Text S2). No language restrictions were used. Bibliographies from other relevant review articles were cross-examined for potential missed studies. Disagreement was resolved by a third reviewer. Citations were downloaded into reference management software and duplicate citations were electronically or manually removed.

We systematically included the studies using the following criteria: 1) original data from prospective or retrospective cohort studies and randomized clinical trials (RCTs) in humans; 2) reporting at least two treatments, including resection or any local ablative therapy (RES, RFA, MWA, PEI, or TACE+RFA (TR)); 3) mean lesion size ≤ 5 cm; 4) evaluating overall survival rate not less than one year after first or recurrent treatments. Conference abstracts and case reports were excluded, as were older publications from studies with multiple publications.

Patients and public involvement

The patients or public were not involved in the study.

Data Extraction and Study Quality

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Two investigators independently extracted and cross-checked the data from the eligible studies: author, year, study design, country, disease type, inclusion criteria, treatment style, study size, gender, age, tumor size, follow-up duration, treatment complications and survival outcomes. If in disagreement, a third reviewer adjudicated. The level of evidence was appraised using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) guidance ¹⁴, which was classified into four levels of high, moderate, low, and very low. The quality score was downgraded according to 5 domains, including risk of bias, inconsistency, indirectness, imprecision, and publication bias while scores were upgraded according to large effect, appropriate control for plausible confounding, and dose-response gradient.

Data Analysis

Network meta-analysis was used if a ring or open evidence loop was available to know the number of arms and the sample size of each intervention. When possible, pair-wise direct head-to-head comparisons were conducted to calculate the odds ratio (OR) of 1-, 3- and 5-year survival and their 95% confidence intervals (CI). Between-study heterogeneity was evaluated using the tau-squared statistic (τ^2) ¹⁵. A node-splitting analysis was applied to check the consistency between direct evidence (existing real reported comparisons) and indirect evidence (estimated treatment comparisons) for their agreement on a specific node ¹⁶. Bayesian network meta-analysis with Markov Chain Monte Carlo (MCMC), through a consistency

model, was utilized to estimate the pooled ORs and its 95% credible interval (CrI) for the direct and indirect comparisons¹⁶. The inconsistency model was used to check for heterogeneity due to chance imbalance in the distribution of effect modifiers. Consistency in every closed loop was checked by the loop-specific approach in order to estimate whether treatment survival effects were disturbed by variance in the distribution of potential confounding factors among the studies. In order to compare and rank survival rates of different treatments, we examined all studies first and then separately assessed smaller (<3cm) and larger (3-5cm) tumors. Random-effect meta-regression models were used, with and without adjustment for study design (cohort or RCT) and subgroup analyses were also conducted for RCTs in order to examine treatment effectiveness. We appraised the ranking probabilities for all therapies for each intervention and the treatment hierarchy was ordered by the surface under the cumulative ranking curve (SUCRA)¹⁷. Sensitivity analysis was conducted to remove each study, in turn, and estimate the treatment effect in the remaining studies. Funnel plots were utilized to check the possible presence of publication bias or small-study bias¹⁸. In this study, we used Bayesian MCMC simulations by WinBUGS 1.4 and graphically presented the results using Stata 13.

Results

Study Characteristics

After screening, 74 relevant studies in 73 articles were identified, of which 20 were randomized controlled trials and 54 were cohort studies¹⁹⁻⁹². We excluded

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

136504 duplicate or non-relevant citations (Figure 1). The summary characteristics of these studies are shown in Additional file 1: Table S1. Overall, 32345 patients of mean age from 46 to 73.5 years, with approximately 29236 tumors, were assigned to receive RES, RFA, MWA, TR and PEI, and the mean follow-up ranged from 1.5 to 5.7 years. In addition, the numbers of connected studies to the lines (black) and sample size of each treatment (red) were shown in Figure 2 and 3, respectively.

Network Meta-Analysis Results

Ten possible treatment comparisons among the five interventions were examined in the included studies. Comparable survival estimates were made for each treatment (per 1000 patients) and the survival OR among each of the treatment comparisons, according to follow-up duration, are presented in Additional file 1: Table S2, along with estimation of the quality of evidence using GRADE criteria.

Across the range of treatment comparisons and follow-up durations, evidence was graded between low and high quality. Evidence was often graded as low quality owing to publication bias and graded as high quality owing to a larger number of participants in direct comparisons.

Survival probabilities (estimated using Meanrank) and ranks for the five treatments in patients with tumors <3cm, 3-5cm or ≤5cm (with and without adjustment for study design) are graphically displayed in Figures 2-5, and numerical details are given in Additional file 1: Table S3-S4. RES was consistently associated with greater survival (rank 1) compared to MWA, RFA, TR and PEI for the 5-year

1
2
3 survival estimates. The ranks were similar for 1 and 3-year survival with RES or TR
4
5 being ranked as 1 or 2 in most analyses. After adjustment for study design, and in the
6
7 full sample of available studies (n=74), the treatments were ranked as follows for
8
9 5-year survival: 1) RES, 2) TR, 3) RFA, 4) MWA, and 5) PEI (Table S4).
10
11
12

13 Efficacy comparisons from network meta-regression for all treatments are
14
15 summarized in Table 1 and 2, according to follow-up duration and initial tumor size.
16
17 Compared to RES, the 5-year survival in all studies (trials and observational studies)
18
19 for all tumors ≤ 5 cm, was 0.45 (95%CrI 0.23 to 0.82) for PEI, 0.59 (95%CrI 0.25 to
20
21 1.20) for TR, 0.55 (95%CrI 0.25 to 1.05) for MWA and 0.52 (95%CrI 0.29 to 0.88)
22
23 for RFA (Table 2). When examining the comparisons across all treatments, the only
24
25 significant difference for tumors < 3 cm was for 5-year survival, and a significantly
26
27 worse survival was observed for PEI compared to RES 0.43 (95%CrI 0.17 to 0.89).
28
29 For tumors between 3 and 5 cm, no significant differences were observed at 5-year
30
31 survival, but significantly worse 3-year survival was observed with PEI, MWA and
32
33 RFA compared to RES (Table 2). Despite smaller number of studies in analyses of
34
35 only RCTs, the pairwise comparisons showed similar results. However, all relative
36
37 rankings should be interpreted with caution because most network meta-regression
38
39 comparisons did not suggest a statistically significant difference between treatments.
40
41 Detailed results of each comparison for survival rates were shown in Additional file 1:
42
43 Table S5-S10.
44
45
46
47
48
49
50
51

52 Loop-specific methods detected no inconsistency between the pairwise and
53
54 network meta-analysis for most closed loops in the network (Additional file 1: Figure
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

S1). However, inconsistency was observed between direct and indirect comparisons for the following loops: lesions <3cm: RES-RFA-TR, PEI-RES-RFA, MWA-RES-RFA; lesions 3-5cm: MWA-RES-RFA, RES-RFA-TR; and lesions ≤5cm: RES-RFA-TR). In addition, tests for inconsistency were carried out (Additional file 1: Table S11-S13), which indicated a close relationship of between-trial heterogeneity and inconsistency between “direct” and “indirect” evidence.

Sensitivity Analysis and publication bias

No significant change was observed when any one study was deleted. Funnel plots indicated that the included studies in each group were distributed symmetrically around the vertical line (x=0), suggesting that no obvious evidence of publication bias or small-sample effect existed in this network (Additional file 1: Figure S2).

Discussion

There were many techniques for attaining a large ablated zone and complete necrosis of HCC and this comprehensive review addressed two of the more common treatments, namely resection and ablation. In this network meta-analysis, of the five examined therapies, the pooled data showed RES ranked best in full sample analysis with or without adjustment for study design. In both smaller (<3cm) and larger tumors (3-5cm) RES remained the highest ranking treatment. However, most of the individual treatment comparisons were not statistically significant and thus, RES may not be superior to all other therapies. Our evidence indicated locoregional therapies and particularly RES or TR (TACE+RFA) were associated with longer survival.

Our observation of better survival outcomes with TR may be through the advantage of dual mechanisms. With TR, TACE induced hypoxic injury on cancer cells through occlusion of blood vessels and was followed by local ablation. This combination therapy may result in a larger ablated zone⁹³, reducing the possibility of micrometastasis and recurrence, and thus, resulting in better survival outcomes than RFA alone.

While being more invasive, and despite risk of complications, RES was associated with better survival outcomes after 1 year, 3 years and 5 years. This may be due to removal of larger sections of liver than can be targeted with locoregional therapies, thus removing a larger area of potentially cancerous cells. Additionally, rat models indicated that the liver has the potential to quickly restore its original size after partial hepatectomy. This may be mediated via interactions of lipopolysaccharide (LPS), tumor necrosis factor (TNF) α , interleukin (IL)-6, and transforming growth factor β (TGF β)⁹⁴. However, evidence from rat models and human studies indicated that resection success was associated with resection size and regeneration was stunted with larger resections⁹⁵⁻⁹⁷. The safe limit for remnant liver volume in normal liver was approximately 30% of total liver volume, but this was estimated to rise to 40-50% in those with liver disease^{95 98}. Liver resection was recognised as the most efficient treatment for HCC but was only applicable for less than 30% of all patients. However, developments in preoperative imaging techniques, laproscopic surgery and newly developing combinations with chemotherapy may extend its application to more advanced tumors⁹⁸. Furthermore, the consistent associations observed with all

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

studies and only in RCTs indicated that patient selection bias in the observational studies does not wholly explain the better survival outcomes with RES.

Overall, we found PEI was associated with shorter survival than the other four therapies, a finding which is supported in previous studies^{20 29}. One study reported RFA was superior to PEI in achieving short- and long-term survival outcomes, although PEI and RFA showed similar 5-year survival in lesions <3 cm⁵¹. The possible reason why PEI is less effective than RFA may be because lesions often have a thick capsule and therefore ethanol may not distribute through tissues.

There are several limitations in this study. Firstly, a major limitation is in the inclusion of non-randomised studies, in which selection bias is likely to confound observations. Selection of treatment is likely to be based on individual or tumor characteristics, and thus these factors will bias and confound observations of survival. Secondly, this study included both RCTs and observational studies, in which study designs and type of data collection may not be comparable. However, findings were consistent among both study designs. Thirdly, all included studies did not report our primary outcome of interest (5-year survival) and this was a particular limitation among randomised studies. Fourthly, for many individual comparisons, there were either no direct comparisons or comparisons from only a small number of studies. The lack of evidence may increase the risk of bias, which could enlarge or undervalue effect size, and may explain the small inconsistency seen between direct and estimated comparisons. Thus, we should be cautious in interpreting treatment rankings for the different survival times and for different size lesions. While adverse

events from treatments may differ (not evaluated in detail in this review), by examining overall survival outcomes in our review, we have taken account of both long-term potential benefits and harms from treatments. The focus of these findings should therefore be on the overall observation that RES or TR may be superior in terms of survival, rather than focusing on specific OR values for individual treatment comparisons.

In conclusion, the findings of the current bayesian network meta-analysis indicate that RES or TR may be among the most effective therapeutic approaches for HCC for 5-year survival in both smaller ($< 3\text{cm}$) and larger ($3\text{-}5\text{cm}$) lesions. However, evidence was of variable quality, and the majority of evidence came from non randomised studies, which are prone to selection bias and knowledge gaps still exist. For not, at the individual level, selection of strategies should depend on patient and clinical characteristics. To facilitate generation of evidence-based recommendations for HCC therpy, and to standardize treatment approaches, further head-to-head comparisons, especially of resection and ablative therapies, are required from high-quality RCTs, with long follow-up for survival outcomes.

Conflict of interests

The authors have declared that no competing interests regarding the publication of this paper.

Data sharing statement

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

No additional data are available.

Acknowledgments

This study was supported by the opening foundation of the State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital of Medical College, Zhejiang University, grant NO. 2015KF06; This study was supported by the Foundation of Zhejiang Health Committee (2017KY346).

References

1. Collaborators GMaCoD. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;385:117-71
2. Lozano R, Naghavi M, Foreman K, *et al.* Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095-128
3. Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: Global Burden of Disease Study. *Lancet* 1997;349:1269-76
4. El-Serag HB. Hepatocellular carcinoma. *The New England journal of medicine* 2011;365:1118-27
5. Ferlay J, Shin HR, Bray F, *et al.* Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *International journal of cancer* 2010;127:2893-917
6. Forner A, Llovet JM, Bruix J. Hepatocellular carcinoma. *Lancet* 2012;379:1245-55
7. Raza A, Sood GK. Hepatocellular carcinoma review: current treatment, and evidence-based medicine. *World journal of gastroenterology* 2014;20:4115-27
8. Kishi Y, Hasegawa K, Sugawara Y, *et al.* Hepatocellular carcinoma: current management and future development-improved outcomes with surgical resection. *International journal of hepatology* 2011;2011:728103
9. Mazzaferro V, Bhoori S, Sposito C, *et al.* Milan criteria in liver transplantation

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

for hepatocellular carcinoma: an evidence-based analysis of 15 years of experience. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society* 2011;17 Suppl 2:S44-57

10. Shan, Xue-Mei, Ding, *et al.* Radiofrequency ablation of hepatocellular carcinoma sized > 3 and ≤ 5 cm: Is ablative margin of more than 1 cm justified? *World journal of gastroenterology* 2011;19:7389-98

11. Yu SJ. A concise review of updated guidelines regarding the management of hepatocellular carcinoma around the world: 2010-2016. *Clinical & Molecular Hepatology* 2016;22:7-17

12. Chaimani A, Higgins JP, Mavridis D, *et al.* Graphical tools for network meta-analysis in STATA. *PloS one* 2013;8:e76654

13. Hutton B, Salanti G, Caldwell DM, *et al.* The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Annals of internal medicine* 2015;162:777-84

14. Puhan MA, Schunemann HJ, Murad MH, *et al.* A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis. *Bmj* 2014;349:g5630

15. Higgins JP, Green S. Cochrane Handbook for Systematic Reviews of Interventions. *Naunyn-Schmiedebergs Archiv für experimentelle Pathologie und Pharmakologie* 2009;2011:S38

16. Dias S, Welton NJ, Caldwell DM, *et al.* Checking consistency in mixed treatment comparison meta-analysis. *Statistics in medicine* 2010;29:932-44
17. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology* 2011;64:163-71
18. Jin ZC, Zhou XH, He J. Statistical methods for dealing with publication bias in meta-analysis. *Statistics in medicine* 2015;34:343-60
19. Zhang Z, Wu M, Chen H, *et al.* [Percutaneous radiofrequency ablation combined with transcatheter arterial chemoembolization for hepatocellular carcinoma]. *Zhonghua wai ke za zhi [Chinese journal of surgery]* 2002;40:826-9
20. Lencioni RA, Allgaier HP, Cioni D, *et al.* Small hepatocellular carcinoma in cirrhosis: randomized comparison of radio-frequency thermal ablation versus percutaneous ethanol injection. *Radiology* 2003;228:235-40
21. Lin SM, Lin CJ, Lin CC, *et al.* Radiofrequency ablation improves prognosis compared with ethanol injection for hepatocellular carcinoma < or =4 cm. *Gastroenterology* 2004;127:1714-23
22. Vivarelli M, Guglielmi A, Ruzzenente A, *et al.* Surgical resection versus percutaneous radiofrequency ablation in the treatment of hepatocellular carcinoma on cirrhotic liver. *Annals of surgery* 2004;240:102-7
23. Cho CM, Tak WY, Kweon YO, *et al.* [The comparative results of radiofrequency ablation versus surgical resection for the treatment of

hepatocellular carcinoma]. *The Korean journal of hepatology* 2005;11:59-71

24. Hong SN, Lee SY, Choi MS, *et al.* Comparing the outcomes of radiofrequency ablation and surgery in patients with a single small hepatocellular carcinoma and well-preserved hepatic function. *Journal of clinical gastroenterology* 2005;39:247-52

25. Huang GT, Lee PH, Tsang YM, *et al.* Percutaneous ethanol injection versus surgical resection for the treatment of small hepatocellular carcinoma: a prospective study. *Annals of surgery* 2005;242:36-42

26. Lin SM, Lin CJ, Lin CC, *et al.* Randomised controlled trial comparing percutaneous radiofrequency thermal ablation, percutaneous ethanol injection, and percutaneous acetic acid injection to treat hepatocellular carcinoma of 3 cm or less. *Gut* 2005;54:1151-6

27. Lu MD, Xu HX, Xie XY, *et al.* Percutaneous microwave and radiofrequency ablation for hepatocellular carcinoma: a retrospective comparative study. *Journal of gastroenterology* 2005;40:1054-60

28. Montorsi M, Santambrogio R, Bianchi P, *et al.* Survival and recurrences after hepatic resection or radiofrequency for hepatocellular carcinoma in cirrhotic patients: a multivariate analysis. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2005;9:62-7; discussion 67-8

29. Shiina S, Teratani T, Obi S, *et al.* A randomized controlled trial of radiofrequency ablation with ethanol injection for small hepatocellular

- carcinoma. *Gastroenterology* 2005;129:122-30
30. Chen MS, Li JQ, Zheng Y, *et al.* A prospective randomized trial comparing percutaneous local ablative therapy and partial hepatectomy for small hepatocellular carcinoma. *Annals of surgery* 2006;243:321-8
31. Lu MD, Kuang M, Liang LJ, *et al.* [Surgical resection versus percutaneous thermal ablation for early-stage hepatocellular carcinoma: a randomized clinical trial]. *Zhonghua yi xue za zhi* 2006;86:801-5
32. Cho YB, Lee KU, Suh KS, *et al.* Hepatic resection compared to percutaneous ethanol injection for small hepatocellular carcinoma using propensity score matching. *Journal of gastroenterology and hepatology* 2007;22:1643-9
33. Gao W, Chen MH, Yan K, *et al.* Therapeutic effect of radiofrequency ablation in unsuitable operative small hepatocellular carcinoma. *Chinese Journal of Medical Imaging Technology* 2007
34. Lupo L, Panzera P, Giannelli G, *et al.* Single hepatocellular carcinoma ranging from 3 to 5 cm: radiofrequency ablation or resection? *HPB : the official journal of the International Hepato Pancreato Biliary Association* 2007;9:429-34
35. Zhou T, Qiu YD, Kong WT. Comparing the effect of radiofrequency ablation and surgical resection for the treatment of small hepatocellular. *Journal of Hepatobiliary Surgery* 2007
36. Abu-Hilal M, Primrose JN, Casaril A, *et al.* Surgical resection versus radiofrequency ablation in the treatment of small unifocal hepatocellular

carcinoma. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2008;12:1521-6

37. Brunello F, Veltri A, Carucci P, *et al.* Radiofrequency ablation versus ethanol injection for early hepatocellular carcinoma: A randomized controlled trial. *Scandinavian journal of gastroenterology* 2008;43:727-35

38. Guglielmi A, Ruzzenente A, Valdegamberi A, *et al.* Radiofrequency ablation versus surgical resection for the treatment of hepatocellular carcinoma in cirrhosis. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2008;12:192-8

39. Hiraoka A, Horiike N, Yamashita Y, *et al.* Efficacy of radiofrequency ablation therapy compared to surgical resection in 164 patients in Japan with single hepatocellular carcinoma smaller than 3 cm, along with report of complications. *Hepato-gastroenterology* 2008;55:2171-4

40. Ohmoto K, Yoshioka N, Tomiyama Y, *et al.* Comparison of therapeutic effects between radiofrequency ablation and percutaneous microwave coagulation therapy for small hepatocellular carcinomas. *Journal of gastroenterology and hepatology* 2009;24:223-7

41. Sakaguchi H, Seki S, Tsuji K, *et al.* Endoscopic thermal ablation therapies for hepatocellular carcinoma: a multi-center study. *Hepatology research : the official journal of the Japan Society of Hepatology* 2009;39:47-52

42. Santambrogio R, Opocher E, Zuin M, *et al.* Surgical resection versus laparoscopic radiofrequency ablation in patients with hepatocellular carcinoma

- and Child-Pugh class a liver cirrhosis. *Annals of surgical oncology* 2009;16:3289-98
43. Shibata T, Isoda H, Hirokawa Y, *et al.* Small hepatocellular carcinoma: is radiofrequency ablation combined with transcatheter arterial chemoembolization more effective than radiofrequency ablation alone for treatment? *Radiology* 2009;252:905-13
44. Ueno S, Sakoda M, Kubo F, *et al.* Surgical resection versus radiofrequency ablation for small hepatocellular carcinomas within the Milan criteria. *Journal of hepato-biliary-pancreatic surgery* 2009;16:359-66
45. Xiang-Yang BU, Wang Y, Zhong GE, *et al.* Comparison of radiofrequency ablation and surgical resection for small primary liver carcinoma. *Chinese Archives of General Surgery* 2009
46. Guo WX, Zhai B, Lai EC, *et al.* Percutaneous radiofrequency ablation versus partial hepatectomy for multicentric small hepatocellular carcinomas: a nonrandomized comparative study. *World journal of surgery* 2010;34:2671-6
47. Huang J, Yan L, Cheng Z, *et al.* A randomized trial comparing radiofrequency ablation and surgical resection for HCC conforming to the Milan criteria. *Annals of surgery* 2010;252:903-12
48. Kagawa T, Koizumi J, Kojima S, *et al.* Transcatheter arterial chemoembolization plus radiofrequency ablation therapy for early stage hepatocellular carcinoma: comparison with surgical resection. *Cancer* 2010;116:3638-44

49. Morimoto M, Numata K, Kondou M, *et al.* Midterm outcomes in patients with intermediate-sized hepatocellular carcinoma: a randomized controlled trial for determining the efficacy of radiofrequency ablation combined with transcatheter arterial chemoembolization. *Cancer* 2010;116:5452-60

50. Azab M, Zaki S, El-Shetey AG, *et al.* Radiofrequency ablation combined with percutaneous ethanol injection in patients with hepatocellular carcinoma. *Arab journal of gastroenterology : the official publication of the Pan-Arab Association of Gastroenterology* 2011;12:113-8

51. Giorgio A, Di Sarno A, De Stefano G, *et al.* Percutaneous radiofrequency ablation of hepatocellular carcinoma compared to percutaneous ethanol injection in treatment of cirrhotic patients: an Italian randomized controlled trial. *Anticancer research* 2011;31:2291-5

52. Hung HH, Chiou YY, Hsia CY, *et al.* Survival rates are comparable after radiofrequency ablation or surgery in patients with small hepatocellular carcinomas. *Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association* 2011;9:79-86

53. Nishikawa H, Inuzuka T, Takeda H, *et al.* Comparison of percutaneous radiofrequency thermal ablation and surgical resection for small hepatocellular carcinoma. *BMC gastroenterology* 2011;11:143

54. Yun WK, Choi MS, Choi D, *et al.* Superior long-term outcomes after surgery in child-pugh class a patients with single small hepatocellular carcinoma

- compared to radiofrequency ablation. *Hepatology international* 2011;5:722-9
55. Zhang J, Liu HC, Zhou L. The effectiveness of radiofrequency ablation for the treatment of liver cancer. *Journal of Hepatobiliary Surgery* 2011;19:30-33
56. Zhao M, Wang JP, Li W, *et al.* [Comparison of safety and efficacy for transcatheter arterial chemoembolization alone and plus radiofrequency ablation in the treatment of single branch portal vein tumor thrombus of hepatocellular carcinoma and their prognosis factors]. *Zhonghua yi xue za zhi* 2011;91:1167-72
57. Feng K, Yan J, Li X, *et al.* A randomized controlled trial of radiofrequency ablation and surgical resection in the treatment of small hepatocellular carcinoma. *Journal of hepatology* 2012;57:794-802
58. Peng ZW, Lin XJ, Zhang YJ, *et al.* Radiofrequency ablation versus hepatic resection for the treatment of hepatocellular carcinomas 2 cm or smaller: a retrospective comparative study. *Radiology* 2012;262:1022-33
59. Peng ZW, Zhang YJ, Liang HH, *et al.* Recurrent hepatocellular carcinoma treated with sequential transcatheter arterial chemoembolization and RF ablation versus RF ablation alone: a prospective randomized trial. *Radiology* 2012;262:689-700
60. Signoriello S, Annunziata A, Lama N, *et al.* Survival after locoregional treatments for hepatocellular carcinoma: a cohort study in real-world patients. *TheScientificWorldJournal* 2012;2012:564706
61. Wang JH, Wang CC, Hung CH, *et al.* Survival comparison between surgical

resection and radiofrequency ablation for patients in BCLC very early/early stage hepatocellular carcinoma. *Journal of hepatology* 2012;56:412-8

62. Desiderio J, Trastulli S, Pasquale R, *et al.* Could radiofrequency ablation replace liver resection for small hepatocellular carcinoma in patients with compensated cirrhosis? A 5-year follow-up. *Langenbeck's archives of surgery / Deutsche Gesellschaft fur Chirurgie* 2013;398:55-62

63. Ding J, Jing X, Liu J, *et al.* Comparison of two different thermal techniques for the treatment of hepatocellular carcinoma. *European journal of radiology* 2013;82:1379-84

64. Guo WX, Sun JX, Cheng YQ, *et al.* Percutaneous radiofrequency ablation versus partial hepatectomy for small centrally located hepatocellular carcinoma. *World journal of surgery* 2013;37:602-7

65. Hasegawa K, Kokudo N, Makuuchi M, *et al.* Comparison of resection and ablation for hepatocellular carcinoma: a cohort study based on a Japanese nationwide survey. *Journal of hepatology* 2013;58:724-9

66. Iida H, Aihara T, Ikuta S, *et al.* A comparative study of therapeutic effect between laparoscopic microwave coagulation and laparoscopic radiofrequency ablation. *Hepato-gastroenterology* 2013;60:662-5

67. Imai K, Beppu T, Chikamoto A, *et al.* Comparison between hepatic resection and radiofrequency ablation as first-line treatment for solitary small-sized hepatocellular carcinoma of 3 cm or less. *Hepatology research : the official journal of the Japan Society of Hepatology* 2013;43:853-64

68. Kim JW, Shin SS, Kim JK, *et al.* Radiofrequency ablation combined with transcatheter arterial chemoembolization for the treatment of single hepatocellular carcinoma of 2 to 5 cm in diameter: comparison with surgical resection. *Korean journal of radiology* 2013;14:626-35
69. Lai EC, Tang CN. Radiofrequency ablation versus hepatic resection for hepatocellular carcinoma within the Milan criteria--a comparative study. *International journal of surgery* 2013;11:77-80
70. Lin ZZ, Shau WY, Hsu C, *et al.* Radiofrequency ablation is superior to ethanol injection in early-stage hepatocellular carcinoma irrespective of tumor size. *PloS one* 2013;8:e80276
71. Peng ZW, Zhang YJ, Chen MS, *et al.* Radiofrequency ablation with or without transcatheter arterial chemoembolization in the treatment of hepatocellular carcinoma: a prospective randomized trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2013;31:426-32
72. Tohme S, Geller DA, Cardinal JS, *et al.* Radiofrequency ablation compared to resection in early-stage hepatocellular carcinoma. *HPB : the official journal of the International Hepato Pancreato Biliary Association* 2013;15:210-7
73. Wong KM, Yeh ML, Chuang SC, *et al.* Survival comparison between surgical resection and percutaneous radiofrequency ablation for patients in Barcelona Clinic Liver Cancer early stage hepatocellular carcinoma. *Indian journal of gastroenterology : official journal of the Indian Society of Gastroenterology* 2013;32:253-7

74. Zhang L, Wang N, Shen Q, *et al.* Therapeutic efficacy of percutaneous radiofrequency ablation versus microwave ablation for hepatocellular carcinoma. *PloS one* 2013;8:e76119

75. Abdelaziz A, Elbaz T, Shousha HI, *et al.* Efficacy and survival analysis of percutaneous radiofrequency versus microwave ablation for hepatocellular carcinoma: an Egyptian multidisciplinary clinic experience. *Surgical endoscopy* 2014;28:3429-34

76. Shi J, Sun Q, Wang Y, *et al.* Comparison of microwave ablation and surgical resection for treatment of hepatocellular carcinomas conforming to Milan criteria. *Journal of gastroenterology and hepatology* 2014;29:1500-7

77. Yang HJ, Lee JH, Lee DH, *et al.* Small single-nodule hepatocellular carcinoma: comparison of transarterial chemoembolization, radiofrequency ablation, and hepatic resection by using inverse probability weighting. *Radiology* 2014;271:909-18

78. Zhang T, Li K, Luo H, *et al.* [Long-term outcomes of percutaneous microwave ablation versus repeat hepatectomy for treatment of late recurrent small hepatocellular carcinoma: a retrospective study]. *Zhonghua yi xue za zhi* 2014;94:2570-2

79. Pompili M, De Matthaeis N, Saviano A, *et al.* Single hepatocellular carcinoma smaller than 2 cm: are ethanol injection and radiofrequency ablation equally effective? *Anticancer research* 2015;35:325-32

80. Xu J, Zhao Y. Comparison of percutaneous microwave ablation and

- laparoscopic resection in the prognosis of liver cancer. *International journal of clinical and experimental pathology* 2015;8:11665-9
81. Lee HW, Lee JM, Yoon JH, *et al.* A prospective randomized study comparing radiofrequency ablation and hepatic resection for hepatocellular carcinoma. 2018;94:74-82
82. Li W, Zhou X, Huang Z, *et al.* Short-term and long-term outcomes of laparoscopic hepatectomy, microwave ablation, and open hepatectomy for small hepatocellular carcinoma: a 5-year experience in a single center. *Annals of surgical treatment and research* 2017;47:650-57
83. Liu PH, Hsu CY, Hsia CY, *et al.* Surgical Resection Versus Radiofrequency Ablation for Single Hepatocellular Carcinoma ≤ 2 cm in a Propensity Score Model. *Annals of surgery* 2016;263:538-45
84. Liu H, Wang ZG, Fu SY, *et al.* Randomized clinical trial of chemoembolization plus radiofrequency ablation versus partial hepatectomy for hepatocellular carcinoma within the Milan criteria. *The British journal of surgery* 2016;103:348-56
85. Hof J, Wertenbroek MW, Peeters PM, *et al.* Outcomes after resection and/or radiofrequency ablation for recurrence after treatment of colorectal liver metastases. *The British journal of surgery* 2016;103:1055-62
86. Vogl TJ, Farshid P, Naguib NN, *et al.* Ablation therapy of hepatocellular carcinoma: a comparative study between radiofrequency and microwave ablation. *Abdominal imaging* 2015;40:1829-37

87. Lee YH, Hsu CY, Chu CW, *et al.* Radiofrequency ablation is better than surgical resection in patients with hepatocellular carcinoma within the Milan criteria and preserved liver function: a retrospective study using propensity score analyses. *Journal of clinical gastroenterology* 2015;49:242-9

88. Kang TW, Kim JM, Rhim H, *et al.* Small Hepatocellular Carcinoma: Radiofrequency Ablation versus Nonanatomic Resection--Propensity Score Analyses of Long-term Outcomes. *Radiology* 2015;275:908-19

89. Zhou Z, Lei J, Li B, *et al.* Liver resection and radiofrequency ablation of very early hepatocellular carcinoma cases (single nodule <2 cm): a single-center study. *European journal of gastroenterology & hepatology* 2014;26:339-44

90. Ko S, Jo H, Yun S, *et al.* Comparative analysis of radiofrequency ablation and resection for resectable colorectal liver metastases. *World journal of gastroenterology* 2014;20:525-31

91. Kim JM, Kang TW, Kwon CH, *et al.* Single hepatocellular carcinoma \leq 3 cm in left lateral segment: liver resection or radiofrequency ablation? *World journal of gastroenterology* 2014;20:4059-65

92. Agcaoglu O, Aliyev S, Karabulut K, *et al.* Complementary use of resection and radiofrequency ablation for the treatment of colorectal liver metastases: an analysis of 395 patients. *World journal of surgery* 2013;37:1333-9

93. Goldberg SN, Ginn GD, Lukyanov AN, *et al.* Percutaneous tumor ablation: increased necrosis with combined radio-frequency ablation and intravenous liposomal doxorubicin in a rat breast tumor model. *Radiology*

- 2002;222:797-804
94. Taub R. Liver regeneration: from myth to mechanism. *Nature reviews Molecular cell biology* 2004;5:836-47
95. Martins PN, Theruvath TP, Neuhaus P. Rodent models of partial hepatectomies. *Liver international : official journal of the International Association for the Study of the Liver* 2008;28:3-11
96. Kele PG, De BM, Ej VDJ, *et al.* Early hepatic regeneration index and completeness of regeneration at 6 months after partial hepatectomy. *British Journal of Surgery* 2012;99:1113-19
97. Preziosi ME, Monga SP. Update on the Mechanisms of Liver Regeneration. *Seminars in Liver Disease* 2017;37:141
98. Morise Z, Kawabe N, Tomishige H, *et al.* Recent advances in liver resection for hepatocellular carcinoma. *Frontiers in Surgery* 2014;1:21

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

File legends:

Figure 1 Flow chart of search.

Figure 2 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in RCTs.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments. The number in red indicates the sample size and the number in black indicates the number of studies.

- i Lesions < 3 cm.
- ii Lesions 3-5 cm.
- iii Lesions ≤ 5 cm.

Figure 3 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in all studies.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments. The number in red indicates the sample size and the number in black indicates the number of studies.

- i Lesions < 3 cm.
- ii Lesions 3-5 cm.
- iii Lesions ≤ 5 cm.

Figure 4 Treatment ranks for 1-year, 3-year and 5-year survival rates, according

to lesion size in RCTs

A Lesions < 3 cm

B Lesions 3-5 cm

C Lesions \leq 5 cm (full sample).

Figure 5 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in all studies.

A Lesions < 3 cm

B Lesions 3-5 cm

C Lesions \leq 5 cm (full sample).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Odds ratios (95% credible interval) according to network meta-analyses for the survival for all pairwise comparisons in randomized controlled trials.

3cm for 1-year survival				
PEI				
1.17 (0.11-4.66)	TR			
0.08 (0-0.38)	0.15 (0-0.80)	MWA		
0.67 (0.28-1.35)	1.25 (0.16-4.64)	173.30 (1.90-537.40)	RFA	
0.64(0.18-1.61)	1.08 (0.15-3.78)	152.70 (1.44-505.80)	0.97 (0.42-1.98)	RES
3cm for 3-year survival				
PEI				
1.02 (0.14-3.56)	TR			
NA	NA	MWA		
0.79 (0.45-1.39)	1.54 (0.25-13.43)	NA	RFA	
0.58 (0.29-1.16)	1.17 (0.16-4.17)	NA	0.75 (0.41-1.31)	RES
3cm for 5-year survival				
PEI				
3.93 (0.03-19.61)	TR			
NA	NA	MWA		
0.94 (0.08-3.97)	2.87 (0.04-13.43)	NA	RFA	
0.50 (0.04-2.04)	0.84 (0.03-4.18)	NA	0.72 (0.10-2.47)	RES
3-5cm for 1-year survival				
PEI				
NA	TR			
NA	NA	MWA		
NA	3.40 (0.64-11.93)	NA	RFA	
NA	1.00 (0-5.00)	NA	0.25 (0-1.47)	RES
3-5cm for 3-year survival				
PEI				
NA	TR			
NA	NA	MWA		
NA	3.98 (0.71-15.22)	NA	RFA	
NA	1.14 (0-6.20)	NA	0.24 (0-1.25)	RES
3-5cm for 5-year survival				
PEI				
NA	TR			
NA	NA	MWA		
NA	7.64 (0.14-42.49)	NA	RFA	
NA	12.87 (0.02-44.43)	NA	1.05 (0.03-5.33)	RES

≤5cm for 1-year survival

PEI	TR	MWA	RFA	RES
0.29 (0.09-0.73)				
0.27 (0.05-0.84)	1.09 (0.16-3.50)			
0.65 (0.33-1.13)	2.69 (1.02-6.04)	3.84 (0.81-11.60)		
0.37 (0.13-0.82)	1.50 (0.48-3.67)	2.01 (0.47-5.70)	0.57 (0.27-1.08)	

≤5cm for 3-year survival

PEI	TR	MWA	RFA	RES
0.64 (0.19-1.67)				
1.05 (0.12-4.56)	1.86 (0.21-7.59)			
0.86 (0.39-1.79)	1.56 (0.66-3.25)	1.77 (0.22-6.24)		
0.55 (0.19-1.44)	0.98 (0.35-2.41)	1.00 (0.16-3.30)	0.65 (0.31-1.29)	

≤5cm for 5-year survival

PEI	TR	MWA	RFA	RES
0.53 (0.06-1.90)				
NA	NA			
0.74 (0.16-2.00)	2.29 (0.41-7.61)	NA		
0.41 (0.11-1.02)	1.35 (0.23-4.69)	NA	0.66 (0.20-1.62)	

The reference treatment (1.00) for all comparisons is listed to the right hand side

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection;

Table 2 Odds ratios (95% credible interval) according to network meta-analyses for the survival for all pairwise comparisons in all studies

3cm for 1-year survival					
PEI					
0.69 (0.14-2.13)	TR				
0.49 (0.18-1.10)	1.08 (0.21-7.87)	MWA			
0.68 (0.38-1.09)	1.48 (0.34-4.23)	1.59 (0.69-3.17)	RFA		
0.63 (0.22-1.44)	1.30 (0.28-3.88)	1.49(0.44-3.85)	0.94 (0.39-1.91)	RES	
3cm for 3-year survival					
PEI					
0.90 (0.29-2.17)	TR				
1.01 (0.47-1.95)	1.38 (0.42-3.40)	MWA			
0.96(0.59-1.50)	1.31 (0.47-2.92)	1.02 (0.57-1.70)	RFA		
0.68 (0.30-1.39)	0.90 (0.31-2.10)	0.73 (0.30-1.55)	0.72 (0.37-1.30)	RES	
3cm for 5-year survival					
PEI					
1.07 (0.31-2.72)	TR				
0.86 (0.39-1.65)	1.03 (0.28-2.73)	MWA			
0.82 (0.48-1.29)	0.99 (0.32-2.39)	1.04 (0.50-1.77)	RFA		
0.43 (0.17-0.89)	0.49 (0.16-0.18)	0.55 (0.19-1.25)	0.54 (0.24-1.05)	RES	
3-5cm for 1-year survival					
PEI					
0.20 (0.05-0.54)	TR				
0.55 (0.09-1.76)	3.39 (0.58-10.44)	MWA			
0.49 (0.18-1.12)	2.99 (1.14-6.58)	1.29 (0.32-3.60)	RFA		
0.06 (0-0.31)	0.36 (0.01-2.08)	0.15 (0-1.00)	0.12 (0-0.63)	RES	
3-5cm for 3-year survival					
PEI					
0.28 (0.04-0.96)	TR				
0.61 (0.08-2.26)	2.62 (0.61-7.90)	MWA			
0.55 (0.12-1.69)	2.38 (0.93-5.38)	1.15 (0.39-2.65)	RFA		
0.06 (0-0.28)	0.26 (0.01-1.10)	0.12 (0.01-0.53)	0.11 (0.01-0.40)	RES	
3-5cm for 5-year survival					
PEI					
5.77 (0.01-2.84)	TR				
4.15 (0.04-5.18)	11.97 (0.19-46.76)	MWA			
0.86 (0.06-2.68)	6.16 (0.27-25.58)	1.26 (0.19-4.04)	RFA		
3.02 (0.01-2.40)	14.31 (0.04-21.06)	1.24 (0.02-4.46)	0.69 (0.04-3.16)	RES	

≤5cm for 1-year survival

PEI	TR	MWA	RFA	RES
0.34 (0.11-0.63)				
0.81 (0.38-1.51)	2.69 (0.99-6.00)			
0.77 (0.51-1.10)	2.55 (1.20-4.85)	1.04 (0.55-1.76)		
0.52 (0.24-0.96)	1.72 (0.66-3.70)	0.70 (0.29-1.39)	0.68 (0.35-1.17)	

≤5cm for 3-year survival

PEI	TR	MWA	RFA	RES
0.64 (0.32-1.16)				
0.98 (0.55-1.65)	1.65 (0.80-3.03)			
0.94 (0.64-1.34)	1.57 (0.89-2.57)	0.99 (0.64-1.47)		
0.59 (0.30-1.04)	0.97 (0.48-1.79)	0.62 (0.32-1.09)	0.63 (0.37-1.01)	

≤5cm for 5-year survival

PEI	TR	MWA	RFA	RES
0.84 (0.35-1.74)				
0.87 (0.46-1.51)	1.16 (0.46-2.46)			
0.87 (0.57-1.26)	1.16 (0.54-2.21)	1.06 (0.64-1.61)		
0.45 (0.23-0.82)	0.59 (0.25-1.20)	0.55 (0.25-1.05)	0.52 (0.29-0.88)	

The reference treatment (1.00) for all comparisons is listed to the right hand side

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection.




Figure 1 Flow chart of search.
254x190mm (300 x 300 DPI)




Figure 2 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in RCTs.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments. The number in red indicates the sample size and the number in black indicates the number of studies.

i Lesions < 3 cm.

ii Lesions 3-5 cm.

iii Lesions ≤ 5 cm.

500x500mm (300 x 300 DPI)




Figure 3 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in all studies.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments. The number in red indicates the sample size and the number in black indicates the number of studies.

- i Lesions < 3 cm.
- ii Lesions 3-5 cm.
- iii Lesions ≤ 5 cm.

227x227mm (300 x 300 DPI)




Figure 4 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in RCTs
A Lesions < 3 cm
B Lesions 3-5 cm
C Lesions ≤ 5 cm (full sample).

193x165mm (300 x 300 DPI)




Figure 5 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in all studies.
A Lesions < 3 cm
B Lesions 3-5 cm
C Lesions ≤ 5 cm (full sample).

193x165mm (300 x 300 DPI)

Text S1.

PRISMA NMA Checklist of Items to Include When Reporting A Systematic Review Involving a Network Meta-analysis

Section/Topic	Item #	Checklist Item	Reported on Page #
TITLE			
Title	1	Identify the report as a systematic review <i>incorporating a network meta-analysis (or related form of meta-analysis).</i>	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: Background: main objectives Methods: data sources; study eligibility criteria, participants, and interventions; study appraisal; and <i>synthesis methods, such as network meta-analysis.</i> Results: number of studies and participants identified; summary estimates with corresponding confidence/credible intervals; <i>treatment rankings may also be discussed. Authors may choose to summarize pairwise comparisons against a chosen treatment included in their analyses for brevity.</i> Discussion/Conclusions: limitations; conclusions and implications of findings. Other: primary source of funding; systematic review registration number with registry name.	5,6
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known, <i>including mention of why a network meta-analysis has been conducted.</i>	7,8
Objectives	4	Provide an explicit statement of questions being addressed, with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	8
METHODS			
Protocol and registration	5	Indicate whether a review protocol exists and if and where it can be accessed (e.g., Web address); and, if available, provide registration information, including registration number.	8,9

Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale. <i>Clearly describe eligible treatments included in the treatment network, and note whether any have been clustered or merged into the same node (with justification).</i>	9,10
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	9
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	9,10,Figure1, Additional file 1: Text S2
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	9,10
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	10
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	11
Geometry of the network	S1	Describe methods used to explore the geometry of the treatment network under study and potential biases related to it. This should include how the evidence base has been graphically summarized for presentation, and what characteristics were compiled and used to describe the evidence base to readers.	11
Risk of bias within individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	11,12
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means). <i>Also describe the use of additional summary measures assessed, such as treatment rankings and surface under the cumulative ranking curve (SUCRA) values, as well as modified approaches used to present summary findings from meta-analyses.</i>	11,12
Planned methods of analysis	14	Describe the methods of handling data and combining results of studies for each network meta-analysis. This should include, but not be limited to: <ul style="list-style-type: none">• <i>Handling of multi-arm trials;</i>• <i>Selection of variance structure;</i>• <i>Selection of prior distributions in Bayesian analyses; and</i>	11,12

		<ul style="list-style-type: none"> • <i>Assessment of model fit.</i> 	
Assessment of Inconsistency	S2	Describe the statistical methods used to evaluate the agreement of direct and indirect evidence in the treatment network(s) studied. Describe efforts taken to address its presence when found.	10,11,12
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	10,11,12
Additional analyses	16	Describe methods of additional analyses if done, indicating which were pre-specified. This may include, but not be limited to, the following: <ul style="list-style-type: none"> • Sensitivity or subgroup analyses; • Meta-regression analyses; • <i>Alternative formulations of the treatment network; and</i> • <i>Use of alternative prior distributions for Bayesian analyses (if applicable).</i> 	11,12
RESULTS†			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	12
Presentation of network structure	S3	Provide a network graph of the included studies to enable visualization of the geometry of the treatment network.	12,13,Figure2-3
Summary of network geometry	S4	Provide a brief overview of characteristics of the treatment network. This may include commentary on the abundance of trials and randomized patients for the different interventions and pairwise comparisons in the network, gaps of evidence in the treatment network, and potential biases reflected by the network structure.	12,13,
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	12, Table1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment.	

Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: 1) simple summary data for each intervention group, and 2) effect estimates and confidence intervals. <i>Modified approaches may be needed to deal with information from larger networks.</i>	12,13, Figure2-5
Synthesis of results	21	Present results of each meta-analysis done, including confidence/credible intervals. <i>In larger networks, authors may focus on comparisons versus a particular comparator (e.g. placebo or standard care), with full findings presented in an appendix. League tables and forest plots may be considered to summarize pairwise comparisons.</i> If additional summary measures were explored (such as treatment rankings), these should also be presented.	12,13,Figure4-5, Additional file 1: Table S1-S13
Exploration for inconsistency	S5	Describe results from investigations of inconsistency. This may include such information as measures of model fit to compare consistency and inconsistency models, <i>P</i> values from statistical tests, or summary of inconsistency estimates from different parts of the treatment network.	12,13
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies for the evidence base being studied.	12,13, Additional file 1: Figure S1-S2
Results of additional analyses	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression analyses, <i>alternative network geometries studied, alternative choice of prior distributions for Bayesian analyses, and so forth</i>).	12,13
DISCUSSION			
Summary of evidence	24	Summarize the main findings, including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy-makers).	14-16
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review level (e.g., incomplete retrieval of identified research, reporting bias). <i>Comment on the validity of the assumptions, such as transitivity and consistency. Comment on any concerns regarding network geometry (e.g., avoidance of certain comparisons).</i>	16
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	17
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. This should also include information regarding whether funding has been received from manufacturers of treatments in the network and/or whether some of the	17

authors are content experts with professional conflicts of interest that could affect use of treatments in the network.

PICOS = population, intervention, comparators, outcomes, study design.

* Text in *italics* indicateS wording specific to reporting of network meta-analyses that has been added to guidance from the PRISMA statement.

† Authors may wish to plan for use of appendices to present all relevant information in full detail for items in this section.

Text S2.

Search strategy:

Pubmed (1950-present)

1. ("TACE" OR "transarterial chemoembolization")
2. ("RFA" OR "radiofrequency ablation" OR "RF ablation" OR "radiofrequency thermal ablation" OR "RTA")
3. (PEI OR "ethanol injection" OR "ethanol ablation" OR "alcohol ablation")
4. ("microwave ablation" OR "microwave thermal ablation" OR MWA)
5. (liver OR hepato*)
6. (neoplas* OR cancer OR tumor OR tumour OR carcinoma OR oncolog*)
7. 1 OR 2 OR 3 OR 4
8. 5 AND 6 AND 7
9. "Ablation Techniques"[Mesh]
10. "Embolization"[Mesh]
11. "Liver Neoplasms"[Mesh]
12. 9 OR 10
13. 12 AND 11
14. 8 OR 13
15. (resection OR surgery OR hepatectomy)
16. (ablation OR injection OR embolization)
17. 5 AND 6 AND 15 AND 16
18. "Hepatectomy"[Mesh]
19. 12 AND 18 AND 11
20. 17 OR 19

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
21. 14 OR 20
- Embase(1980-present)
1. 'TACE':ab,ti
2. 'transarterial chemoembolization':ab,ti
3. 1 OR 2
4. 'rfa':ab,ti
5. 'radiofrequency ablation':ab,ti
6. 'rf ablation':ab,ti
7. 'radiofrequency thermal ablation':ab,ti
8. 'rta':ab,ti
9. 4 OR 5 OR 6 OR 7 OR 8
10. 'PEI':ab,ti
11. 'ethanol injection ':ab,ti
12. 'ethanol ablation ':ab,ti
13. 'alcohol ablation ':ab,ti
14. 10 OR 11 OR 12 OR 13
15. 'microwave ablation ':ab,ti
16. 'microwave thermal ablation ':ab,ti
17. 'MWA ':ab,ti
18. 15 OR 16 OR 17
19. 'liver':ab,ti
20. 'hepato*':ab,ti
21. 19 OR 20
22. 'neoplas*':ab,ti
23. 'cancer ':ab,ti
24. 'tumor ':ab,ti
25. 'tumour ':ab,ti
26. 'carcinoma ':ab,ti
27. 'oncolog*':ab,ti

28. 22 OR 23 OR 24 OR 25 OR 26 OR 27
29. 3 OR 9 OR 14 OR 18
30. 21 AND 28 AND 29
31. 'resection':ab,ti
32. 'surgery':ab,ti
33. 'hepatectomy':ab,ti
34. 31 OR 32 OR 33
35. 'ablation':ab,ti
36. 'injection':ab,ti
37. 'embolization':ab,ti
38. 35 OR 36 OR 37
39. 34 AND 38 AND 21 AND 28
40. 30 OR 39

Scoups

1. TITLE-ABS-KEY ("TACE")
2. TITLE-ABS-KEY ("transarterial chemoembolization")
3. 1 OR 2
4. TITLE-ABS-KEY ("RFA")
5. TITLE-ABS-KEY ("radiofrequency ablation")
6. TITLE-ABS-KEY ("RF ablation")
7. TITLE-ABS-KEY ("radiofrequency thermal ablation")
8. TITLE-ABS-KEY ("RTA")
9. 4 OR 5 OR 6 OR 7 OR 8
10. TITLE-ABS-KEY ("PEI")
11. TITLE-ABS-KEY ("ethanol injection")
12. TITLE-ABS-KEY ("ethanol ablation")
13. TITLE-ABS-KEY ("alcohol ablation")
14. 10 OR 11 OR 12 OR 13
15. TITLE-ABS-KEY ("microwave ablation")

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
16. TITLE-ABS-KEY ("microwave thermal ablation ")
17. TITLE-ABS-KEY ("MWA")
18. 15 OR 16 OR 17
19. TITLE-ABS-KEY ("liver ")
20. TITLE-ABS-KEY ("hepato*")
21. 19 OR 20
22. TITLE-ABS-KEY ("neoplas*")
23. TITLE-ABS-KEY ("cancer")
24. TITLE-ABS-KEY ("tumor")
25. TITLE-ABS-KEY ("tumour")
26. TITLE-ABS-KEY ("carcinoma")
27. TITLE-ABS-KEY ("oncolog*")
28. 22 OR 23 OR 24 OR 25 OR 26 OR 27
29. 3 OR 9 OR 14 OR 18
30. 29 AND 21 AND 28
31. TITLE-ABS-KEY ("resection")
32. TITLE-ABS-KEY ("surgery")
33. TITLE-ABS-KEY ("hepatectomy")
34. 31 OR 32 OR 33
35. TITLE-ABS-KEY ("ablation")
36. TITLE-ABS-KEY ("injection")
37. TITLE-ABS-KEY ("embolization")
38. 35 OR 36 OR 37
39. 34 AND 38 AND 21 AND 28
40. 30 OR 39

- Web of science
1. TS=(ablation)
2. TS=(embolization)
3. 1 OR 2

4. TS=(hepatectomy)
5. TS=(liver neoplasms)
6. 3 AND 4 AND 5
7. TI=(resection)
8. TI=(surgery)
9. TI=(hepatectomy)
10. 7 OR 8 OR 9
11. TI=(ablation)
12. TI=(injection)
13. TI=(embolization)
14. 11 OR 12 OR 13
15. TI=(liver)
16. TI=(hepato*)
17. 15 OR 16
18. TI=(neoplas*)
19. TI=(cancer)
20. TI=(tumor)
21. TI=(tumour)
22. TI=(carcinoma)
23. TI=(oncolog*)
24. 18 OR 19 OR 20 OR 21 OR 22 OR 23
25. 10 AND 14 AND 17 AND 24
26. 3 AND 5
27. TI=(TACE)
28. TI=("transarterial chemoembolization")
29. 27 OR 28
30. TI=(RFA)
31. TI=("radiofrequency ablation")
32. TI=("RF ablation")
33. TI=("radiofrequency thermal ablation")
34. TI=(RTA)

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
35. 30 OR 31 OR 32 OR 33 OR 34
36. TI=(PEI)
37. TI=("ethanol injection")
38. TI=("ethanol ablation")
39. TI=("alcohol ablation")
40. 36 OR 37 OR 38 OR 39
41. TI=("microwave ablation")
42. TI=("microwave thermal ablation")
43. TI=(MWA)
44. 41 OR 42 OR 43
45. 29 OR 35 OR 40 OR 44
46. 46 AND 17 AND 24
47. 6 OR 25 OR 26 OR 46

23

24

25

26

Table S1.

Summary of the studies included in the network meta-analysis.

27

28

29

30

31

32

33

34

35

36

37

38

39

40

Study Year	Design style	Countr ,	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Zhang 2002 ¹⁹	Prospectiv e cohort	China	HCC	0.3-2	RFA	15(15)	13/2	61.8 (38-78)	4.1 (2.4-6.0)	NA	0.80(1y)	0.80(1y)	NA
					TR	15(15)	12/3	57.8 (39-72)	4.6 (2.3-7.1)	NA	1.00(1y)	1.00 (1y)	NA
Lencioni 2003 ²⁰	RCT	Italy	HCC	1.9±0.8	RFA	52(69)	36/16	67±6 (52-78)	2.8±0.6	1.00(1y)	NA	1.00(1y)	15 pain and 10 fever
					PEI	50(73)	30/20	69±7.4 (40-82)	2.8±0.8	0.96(1y)	NA	0.96(1y)	13 pain and 5 fever
Lin 2004 ²¹	RCT	China	HCC	2±0.9	RFA	52(69)	35/17	62±11	2.9±0.8	0.76(3y)	NA	0.35(3y)	1 transient pleural effusion

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					PEI	52(67)	34/18	59±10	2.8±0.8	0.66(3y)	NA	0.17(3y)	1 pain
Vivarelli 2004 ²²	Retrospective cohort	Italy	HCC	2.4	RES	79(92)	57/22	65.2±8.2 (43-81)	≤3/3.1-5 (21/58)	0.81(3y)	0.59(3y)	0.65(3y)	NA
					RFA	79(112)	67/12	67.8±8.7 (41-88)	≤3/3.1-5 (22/57)	0.50(3y)	0.25(3y)	0.33(3y)	NA
Cho 2005 ²³	Retrospective cohort	Korea	HCC	0.1-3	RES	61	48/13	57	3.4±1.0	NA	0.77(3y)	0.77(3y)	2 bleeding, 1 intraabdominal abscess, 1 wound infection
					RFA	99	76/23	58	3.1±0.8	NA	0.80(3y)	0.80(3y)	1 chest wall metastasis, 1 cholecystitis, 1 iatrogenic burn, 1 ileus, 1 hepatic infarction
Huang 2005 ²⁵	RCT	China	HCC	1-4.9	RES	38(42)	27/11	59±11.4	≤2/2.1-3 (24/14)	0.82	NA	0.82	NA
					PEI	38(46)	19/19	63±10.9	≤2/2.1-3 (21/17)	0.45	NA	0.45	NA
Hong 2005 ²⁴	Retrospective cohort	Korea	HCC	2.9(0.4-4.6)	RES	93	69/24	49.2±9.9	2.5±0.8	0.84(3y)	NA	0.84(3y)	NA
					RFA	55	41/14	59.1±9.6	2.4±0.6	0.73(3y)	NA	0.73(3y)	NA
Lin 2005 ²⁶	RCT	China	HCC	2.3±1	RFA	62(78)	40/22	61±10	2.5±1	0.74(3y)	NA	0.74(3y)	2 haemothorax, 1 gastric bleeding and perforation
					PEI	62(76)	39/23	60±8	2.3±0.8	0.60(3y)	NA	0.60(3y)	1 pain
Lu 2005 ²⁷	Retrospective cohort	China	HCC	2.1±1.1	RFA	53(72)	43/10	54.5±11.7 (24-74)	2.6±1.2 (1.0-6.1)	0.38(3y)	NA	0.38(3y)	2 skin burn, 1 puncture wound infection
					MWA	49(98)	44/5	50.1±13.7 (24-74)	2.5±1.2 (0.9-7.2)	0.51(3y)	NA	0.51(3y)	2 puncture wounds, 2 subcapsular hematoma

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Montorsi 2005 ²⁸	Prospective cohort	Italy	HCC	2.1	RES	40	33/7	67±9	<5cm	NA	NA	0.73(3y)	NA
					RFA	58	43/15	67±6		NA	NA	0.60(3y)	NA
Shiina 2005 ²⁹	RCT	Japan	HCC	3.1(0.6-4.3)	RFA	118(184)	79/39	≤65/>65 (44/74)	≤2/>2 (45/73)	NA	NA	0.61(3y)	1 transient jaundice, 1 skin burn, 1 hepatic infarction, 3 neoplastic seeding
					PEI	114(188)	87/27	≤65/>65 (41/73)	≤2/>2 (57/57)	NA	NA	0.45(3y)	1 abscess2 neoplastic seeding
Chen 2006 ³⁰	RCT	China	HCC	2.4±1	RES	90	75/15	49.4±10.9	≤3/3.1-5 (42/48)	0.53	NA	0.53	2 liver failure, 2 gastrointestinal bleeding, 27 ascites
					RFA	71	56/15	51.9±11.2	≤3/3.1-5 (37/34)	0.58	NA	0.58	3 skin burn
Lu 2006 ³¹	RCT	China	Early HCC	1.8	RES	54(56)	37/17	49±14	3.2±1.0	NA	NA	0.86 (3y)	3 wound infection, 1 gastrointestinal bleeding
					RFA	51(57)	42/9	55±13	2.7±1.0	NA	NA	0.87 (3y)	1 peritoneal bleeding, 1 neoplastic seeding
Cho 2007 ³²	Retrospective cohort	Korea	HCC	5.7	RES	130(145)	103/27	56.3±8.8	≤2/2.1-3 (43/87)	0.66	NA	0.66	NA
					PEI	249(275)	181/68	57.7±9.7	≤2/2.1-3 (169/80)	0.49	NA	0.49	NA
Gao 2007 ³³	Retrospective cohort	China	HCC	4.6	RES	34(37)	28/6	51.5 (38-67)	2.58±0.41	0.76	NA	0.76	12 fever, 5 ascites
					RFA	53(84)	41/12	57.1 (31-81)	2.45±0.37	0.62	NA	0.62	2 bleeding, 1 fistula, 1 wound infection, 6 fever, 9 ascites
Lupo 2007 ³⁴	Retrospective cohort	Italy	HCC	2.6	RES	42	33/9	67(28-80)	4.0(3-5)	NA	0.43	0.43	2 urine infection, 1 bilioma, 1 pleural effusion, 1 renal failure, 1 intra-abdominal bleeding

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	60	47/13	68(42-85)	3.65(3-5)	NA	0.32	0.32	2 liver failure, 1 hepatic abscess, 2 pleural effusion, 1 cutaneous metastasis
Zhou 2007 ³⁵	Retrospect ive cohort	China	HCC	0.5-5.9	RES	40(42)	35/5	53±13	≤2/2.1-5 (7/33)	NA	NA	0.75	NA
					RFA	47(54)	37/10	57±14	≤2/2.1-5 (8/39)	NA	NA	0.19	NA
Abu-Hilal 2008 ³⁶	Retrospect ive cohort	Italy and China	Early HCC	3.6	RES	34	26/8	67	3.8(1.3-5)	NA	0.56	0.56	3 hepatic failure
					RFA	34	27/7	65	3(2-5)	NA	0.56	0.56	1 artero-portal fistula
Brunello 2008 ³⁷	RCT	Italy	Early HCC	2.2	RFA	70(89)	49/20	70.3±8.1	1.27±0.54	0.60(3y)	NA	0.60(3y)	1 haemoperitoneum 1 right haemothorax
					PEI	69(88)	43/27	69.0±7.7	1.27±0.57	0.58(3y)	NA	0.58(3y)	1 haemoperitoneum 1 death
Guglielmi 2008 ³⁸	Retrospect ive cohort	Italy	HCC	2.3	RES	91(113)	73/18	≤65/>65 (47/44)	≤3/3.1-6 (31/60)	0.55	0.43	0.48	33 postoperative complications
					RFA	109(153)	88/21	≤65/>65 (38/71)	≤3/3.1-6 (32/77)	0.28	0.14	0.20	11 postoperative complications
Hiraoka 2008 ³⁹	Retrospect ive cohort	Japan	HCC	2.5	RES	59	44/15	62.4±10.6	2.27±0.55	0.59	NA	0.59	1 death, 2 abscess
					RFA	105	76/29	69.4±9.1	1.98±0.52	0.59	NA	0.59	1 biloma, 2 dermatitis
Bu 2009 ⁴⁵	Retrospect ive cohort	China	HCC	2.9(0.5-6)	RES	42(46)	36/6	53.93±10.74	≤3/3.1-5 (14/28)	0.57	0.46	0.50	1 postoperative hemorrhage, 3 pleural effusions, 2 subdiaphragmatic effusion

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	46(54)	40/6	55.89±7.37	≤3/3.1-5 (20/26)	0.50	0.31	0.37	4 pleural effusions, 1 postoperative hemorrhage, 1 skin burn
Ohmoto 2009 ⁴⁰	Retrospect ive cohort	Japan	HCC	2.8±2	RFA	34(37)	25/9	67 (44-78)	1.6 (0.7-2.0)	0.71	NA	0.71	2 pain, 4 fever, 1 bile duct injury, 1 pleural effusion, 1 skin burns, 1 vagovagal reflex
					MWA	49(56)	41/8	64 (38-75)	1.7 (0.8-2.0)	0.37	NA	0.37	11 pain, 17 fever, 9 bile duct injury, 8 pleural effusion, 5 ascites, 4 skin burns, 2 vagovagal reflex, 2 abscess, 2 intraperitoneal bleeding, 1 hepatic infarction, 1 portal thrombus, 1 biliary peritonitis
Sakaguchi 2009 ⁴¹	Retrospect ive cohort	Japan	HCC	0.1-5	Laparosco pic /thorasc opic RFA	249	169/80	65.6±8.9	2.48±0.89	0.57	NA	0.57	1 frequent premature ventricular contractions, 1 liver decompensation
					Laparosco pic /thorasc opic MWA	142	107/35	64.9±7.8	2.28±0.74	0.63	NA	0.63	1 breath holding and incomplete intestinal obstruction, 2 liver decompensation
Santambrogio 2009 ⁴²	Prospectiv e cohort	Italy	HCC	3.2	RES	78	55/23	68±8	2.87±1.21	0.54	NA	0.54	15 extra-hepatic complications

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					Laparoscopic RFA	74	59/15	68±7	2.63±1.07	0.41	NA	0.41	14 extra-hepatic complications
Shibata 2009 ⁴³	RCT	Japan	HCC	2.5±1.2	RFA	43(44)	33/10	69.8±8 (44-87)	1.6±0.5 (0.8-2.6)	0.84(3y)	NA	0.84(3y)	1 pseudoaneurysm
					TR	46(49)	31/15	67.2±8.9 (45-83)	1.7±0.6 (0.9-3.0)	0.85(3y)	NA	0.85(3y)	1 hepatic infarction
Ueno 2009 ⁴⁴	Retrospective cohort	Japan	HCC	3(0.3-7.9)	RES	123(136)	82/41	67(28-85)	2.7±0.1	0.81	0.72	0.80	NA
					RFA	155(209)	100/55	66(40-79)	2.0±0.1	0.38	0.78	0.63	NA
Guo 2010 ⁴⁶	Retrospective cohort	China	HCC	2.5	RES	73(155)	57/16	50.0 (17.0-68.0)	≤3/3.1-5 (30/43)	0.27	0.47	0.44	1 postoperative hemorrhage, 5 abscess, 3 infected ascites, 1 liver failure, 4 pleural effusion
					RFA	86(211)	63/23	52.5 (26.0-80.0)	≤3/3.1-5 (42/44)	0.33	0.16	0.21	1 postoperative hemorrhage, 1 bile leak, 1 abscess, 1 infected ascites, 3 pleural effusion
Huang 2010 ⁴⁷	RCT	China	HCC	3.87	RES	115(144)	85/30	55.91±12.68	≤3/3.1-5 (45/44)	0.82	0.73	0.76	1 hepatic failure, 13 ascites, 5 effusion, 9 bile leakage, 2 postoperative bleeding, 2 gastrointestinal bleeding
					RFA	115(147)	79/36	56.57±14.30	≤3/3.1-5 (57/27)	0.61	0.52	0.55	1 gastric perforation, 2 hemorrhage, 1 malignant seeding, 1 hepatic infarction
Kagawa 2010 ⁴⁸	Retrospective cohort	Japan	Early HCC	4.2	RES	55(69)	40/15	66.1±8.4	≤2/2.1-5 (9/46)	0.42	NA	0.42	2 deaths, 1 liver failure, 1 pleural effusion, 1 pneumonia, 2 biliary leakage

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					TR	62(79)	39/23	67.5 ±8.4	≤2/2.1-5 (19/43)	0.29	NA	0.29	1 duodenal perforation, 1 hemothorax
Morimoto 2010 ⁴⁹	RCT	Japan	HCC	2.7	RFA	18(25)	12/6	73 (48-84)	3.7±0.6	NA	0.78(3y)	0.78(3y)	5 pain, 2 pleural effusion
					TR	19(21)	15/4	70 (57-78)	3.6±0.7	NA	0.95(3y)	0.95(3y)	1 pain, 1 pleural effusion
Azab 2011 ⁵⁰	RCT	Egypt	HCC	1.5	RFA	30(33)	75/15	46-77	<5cm	NA	NA	0.90	5 superficial burn, 17 transient pain, 3 portal vein thrombosis, 7 fever, 1 ascites
					PEI	30(32)				NA	NA	0.83	2 portal vein thrombosis, 3 fever, 3 ascites
Giorgio 2011 ⁵¹	RCT	Italy	HCC	1.8	RFA	142	105/37	70±2 (68-74)	2.34±0.45 (1.1-3)	0.70	NA	0.70	1 major complication
					PEI	143	102/41	72±6 (68-79)	2.27±0.48 (1.3-2.9)	0.68	NA	0.68	3 major complication
Hung 2011 ⁵²	Retrospect ive cohort	China	Early HCC	3.5 ±2	RES	229	184/45	60.07 ±12.56	2.88±1.06	0.77	NA	0.77	NA
					RFA	190	121/69	67.42 ±11.45	2.37±0.92	0.67	NA	0.67	NA
Nishikawa 2011 ⁵³	Retrospect ive cohort	Japan	HCC	3.3	RES	69	50/19	67.4 ±9.7	2.68±0.49	0.74	NA	0.74	2 bile leakage, 2 ascites, 1 acute respiratory distress syndrome, 1 gastrointestinal bleeding
					RFA	162	95/67	68.4 ±8.7	1.99±0.62	0.63	NA	0.63	1 biloma, 1 ascites, 1 intra-abdominal bleeding
Yun 2011 ⁵⁴	Retrospect	Korea	HCC	3.5(0.1-9.	RES	215	171/44	51.7 ±9.7	2.1 ±0.5	0.94	NA	0.94	NA

Study Year	Design style	Countr	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
	ive cohort			1)	RFA	255	197/58	57.0±9.9	2.1±0.5	0.87	NA	0.87	NA
Zhang 2011 ⁵⁵	Retrospect ive cohort	China	HCC	0.5-3.5	RES	103(117)	78/25	56.4±15.2	<5cm	NA	NA	0.35(3y)	12 wound infection, 5 postoperative hemorrhage, 2 hepatic failure, 15 pleural effusions, 6 pleural effusions
					RFA	85(106)	62/23	58.5±12.9	<5cm	NA	NA	0.39(3y)	2 gallbladder cardiac reflex, 4 postoperative hemorrhage, 3 pleural effusions
Feng 2012 ⁵⁷	RCT	China	HCC	3	RES	84(116)	75/9	47 (18-76)	2.6±0.8	0.62(3y)	NA	0.62(3y)	7 pleural effusion, 3 pneumonia, 1 effusion plus infection, 3 wound infection or dehiscence, 1 biliary fistula, 2 abdominal bleeding, 1 pneumothorax or hemothorax
					RFA	84(120)	79/5	51 (24-83)	2.4±0.6	0.55(3y)	NA	0.55(3y)	5 pleural effusion, 1 liver abscess, 2 abdominal bleeding
Peng 2012 ⁵⁸	Retrospect ive cohort	China	Recurr ent HCC	4.9	RES	74	65/9	51.5±12.1 (24-75)	1.1±0.5 (0.8-2.0)	0.62	NA	0.62	1 liver failure, 2 gastrointestinal bleeding, 1 peritoneal bleeding, 1 intestinal obstruction, 1 spontaneous bacterial peritonitis, 1 persistent jaundice, 31 ascites

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	71	63/8	53.1 ±12.1 (28-74)	1.2 ±0.6 (0.9-2.0)	0.72	NA	0.72	1 gastrointestinal bleeding, 1 persistent jaundice, 12 ascites
Peng 2012 ⁵⁹	RCT	China	Recurrent HCC	3.3 ±1.8	RFA	70(76)	55/15	55.1 ±9.5 (22-75)	≤3/3.1-5 (46/24)	NA	0.17	0.36	1 persistent jaundice, 1 ascites, 22 fever, 45 pain, 4 vomiting
					TR	69(74)	59/9	57.5 ±10.0 (19-75)	≤3/3.1-5 (41/28)	NA	0.39	0.46	1 liver failure, 1 ascites, 27 fever, 50 pain, 42 vomiting
Signoriello 2012 ⁶⁰	Retrospective cohort	Italy	HCC	0.1-9	RES	34(44)	30/4	62 ±7	≤3/3.1-5/>5.1 (13/9/4)	NA	NA	0.29	NA
					RFA	50(74)	40/10	68 ±7	≤3/3.1-5/>5.1 (24/11/7)	NA	NA	0.15	NA
					PEI	256(349)	188/68	67 ±8	≤3/3.1-5/>5.1 (143/43/12)	NA	NA	0.20	NA
a. Wang 2012 ⁶¹	Retrospective cohort	China	Early HCC	2.5	RES	52	38/14	≤60 (35)	NA	NA	NA	0.92	NA
					RFA	91	60/31	≤60 (40)		NA	NA	0.73	NA
b. Wang 2012 ⁶²	Retrospective cohort	China	Early HCC	2.5	RES	208	168/40	≤60 (113)	≤2/2.1-5 (6/202)	NA	NA	0.77	NA
					RFA	254	161/93	≤60 (85)	≤2/2.1-5 (60/194)	NA	NA	0.57	NA
Desiderio 2013 ⁶²	Retrospective cohort	Italy	HCC	4.3(2.3-5)	RES	52(94)	37/15	65.6 ±4.8	≤3	0.46	NA	0.46	2 hepatic failure, 1 biliary fistula, 2 hemoperitoneum, 9 ascites
					RFA	44(81)	35/9	64.4 ±6.5		0.36	NA	0.36	6 pain, 7 fever

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Ding 2013 ⁶³	Retrospect ive cohort	China	HCC	2.3±1.3	RFA	85(98)	68/17	58.64±8.52 (40-77)	2.38±0.81 (1.0-4.8)	0.82(3y)	NA	0.82(3y)	1 frequent premature ventricular contractions, 1 liver decompensation
					MWA	113(131)	85/28	59.06±11.68 (30-86)	2.55±0.89 (0.8-5.0)	0.78(3y)	NA	0.78(3y)	1 breath holding and incomplete intestinal obstruction, 2 liver decompensation
Guo 2013 ⁶⁴	Retrospect ive cohort	China	HCC	2.7	RES	102(129)	94/8	51.5(18-75)	≤3/3.1-5 (75/27)	NA	NA	0.63	5 postoperative hemorrhage, 3 bile leak, 4 abscess, 3 infected ascites, 1 liver failure, 4 pleural effusion
					RFA	94(125)	78/16	56(19-75)	≤3/3.1-5 (62/32)	NA	NA	0.50	1 postoperative hemorrhage, 2 bile leak, 1 abscess, 1 infected ascites, 3 pleural effusion
Hasegawa 2013 ⁶⁵	Retrospect ive cohort	Japan	HCC	2.2	RES	5361(646 1)	3967/139 4	66 (48-77)	2.3 (1.2-3)	0.71	NA	0.71	NA
					RFA	5548(741 2)	3569/197 9	69 (52-80)	2 (1-3)	0.61	NA	0.61	NA
					PEI	2059(283 6)	1303/756	69 (52-80)	1.7 (1-3)	0.56	NA	0.56	NA
Iida 2013 ⁶⁶	Retrospect ive cohort	Japan	HCC	0.1-7.5	Laparosco pic RFA	18(27)	NA	73.5±4.0	2.1±0.5	0.78	NA	0.78	1 abscess

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					Laparosco pic MWA	40(56)		70.1 ±6.6	2.0±0.9	0.78	NA	0.78	1 abscess
Imai 2013 ⁶⁷	Retrospect ive cohort	Japan	HCC	4.1	RES	101	75/26	63.3±9.7	2.14±0.55	0.87	NA	0.87	NA
					RFA	82	46/36	67.6±8.5	1.87±0.50	0.60	NA	0.60	NA
Kim 2013 ⁶⁸	Retrospect ive cohort	Korea	Early HCC	0.1-4.2	RES	47	36/11	58.8±10.7	3.66±0.76	NA	0.85(3y)	0.85(3y)	2 pleural effusion, 2 pneumonia, 1 hepatic failure, 1 hepatic abscess, 1 mechanical ileus
					TR	37	31/6	61.7±11.1	3.46±0.75	NA	0.78(3y)	0.78(3y)	1 bile duct dilatation
Lai 2013 ⁶⁹	Retrospect ive cohort	China	HCC	2.9±1.5	RES	80	55/25	60.8±9.9	2.9±1.1	0.71	NA	0.71	NA
					RFA	31	19/12	63.1±12.8	1.8±0.6	0.84	NA	0.84	NA
Lin 2013 ⁷⁰	Retrospect ive cohort	China	Early HCC	3.4	RFA	658	393/265	64.7±10.5	2.4±1.1 (0.8-9.5)	0.60	0.50	0.55	NA
					PEI	378	243/135	63.5±12.1	2.0±0.9 (0.4-7.0)	0.50	0.28	0.40	NA
Peng 2013 ⁷¹	RCT	China	HCC	0.6-5.2	RFA	95(133)	71/24	55.3±13.3	3.39±1.35	NA	0.59(3y)	0.59(3y)	51 pain, 26 fever, 29 vomiting, 4 ascites, 2 pleural effusion, 1 skin burn, 1 abdominal infection, 1 small intestinal obstruction
					TR	94(137)	75/19	53.3±11	3.47±1.44	NA	0.67(3y)	0.67(3y)	57 pain, 33 fever, 40 vomiting, 5 ascites, 3 pleural effusion, 1 skin burn, 1 bile duct stenosis, 1 gastric hemorrhage

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Tohme 2013 ⁷²	Retrospect ive cohort	Ameri ca	Early HCC	2.4	RES	50(62)	31/19	66.3±1	3.07±1.17	0.48	NA	0.48	3 pleural effusion, 1 pneumonia, 1 myocardial infarction, 2 biloma, 2 ileus, 1 ascites, 1 hyperbilirubinaemia >6, 1 renal insufficiency, 2 encephalopathy
					RFA	60(75)	38/22	65.6±12	2.36±0.94	0.35	NA	0.35	1 oesophagitis, 3 encephalopathy, 1 cholangitis, 2 ascites, 1 renal insufficiency, 1 pneumonia
Wong 2013 ⁷³	Retrospect ive cohort	China	Early HCC	0.1-5	RES	46	30/16	55.1±12	2.1±0.6	0.85	NA	0.85	2 fever, 1 increased serum alanine aminotransferase level, 2 atelectasis, 2 biloma
					RFA	36	18/18	63.5±13	1.9±0.6	0.72	NA	0.72	None
Zhang 2013 ⁷⁴	Retrospect ive cohort	China	HCC	2.2±1	RFA	78(97)	64/14	54±10.5 (30-80)	≤3/3.1-5 (47/31)	0.43	0.39	0.41	1 persistent jaundice, 1 biliary fistula
					MWA	77(105)	67/10	54±9.5 (26-76)	≤3/3.1-5 (36/41)	0.58	0.29	0.39	1 hemothorax and intrahepatic hematoma, 1 peritoneal hemorrhage
Abdelaziz 2014 ⁷⁵	RCT	Egypt	Early HCC	2.3	RFA	45(52)	31/14	56.8±7.3	2.95±1.03	0.68(1y)	NA	0.68(1y)	2 subcapsular hematoma, 1 thigh burn, 2 pleural effusion
					MWA	66(76)	48/18	53.6±5	2.9±0.97	0.96(1y)	NA	0.96(1y)	1 subcapsular hematoma, 1 abdominal wall skin burn
Shi 2014 ⁷⁶	Retrospect ive cohort	China	HCC	3.8	RES	107(126)	87/20	54.5±9.9	≤3/3.1-5 (37/54)	0.73	0.57	0.60	NA

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					MWA	117(143)	93/24	56.6±9.2	≤3/3.1-5 (40/56)	0.65	0.52	0.52	NA
Yang 2014 ⁷⁷	Retrospect ive cohort	Korea	HCC	0.1-7	RES	52	38/14	55.7±10.6	≤2/2.1-5 (21/31)	0.94	NA	0.94	2 pneumonia, 1 wound infection, 1 biliary anastomotic leak, 1 portal vein thrombosis, 1 nausea, 1 delirium, 4 ascites
					RFA	79	59/20	57.2±9.2	≤2/2.1-5 (36/43)	0.86	NA	0.86	1 vomiting, 1 ascites, 6 abdominal pain, 2 nausea, 1 sinus bradycardia
Zhang 2014 ⁷⁸	Retrospect ive cohort	China	Recurr ent HCC	2.7	RES	27(29)	25/2	47±13	3.2±1.0	NA	NA	0.63	NA
					MWA	39(46)	37/2	52±13	2.7±1.1	NA	NA	0.62	NA
Pompili 2015 ⁷⁹	Retrospect ive cohort	Italy	Early HCC	2.8	RFA	136	75/61	68 (41-85)	1.8 (1-2)	0.63	NA	0.63	2 ascites, 1 pleural effusion, 1 hemobilia
					PEI	108	90/18	68.5 (34-86)	1.95 (0.8-2)	0.65	NA	0.65	1 hemobilia, 1 portal vein thrombosis
Xu 2015 ⁸⁰	RCT	China	HCC	0.1-3	Laparosco pic RES	45	34/11	58.3±3.1 (26-78)	3.6±0.7 (1-5)	NA	0.38(3y)	0.38(3y)	3 bile leakage, 3 pleural effusion, 2 postoperative hemorrhage
					MWA	45	32/13	57.9±3.4 (27-76)	3.8±0.9 (2-5)	NA	0.33(3y)	0.33(3y)	1 bile leakage, 1 pleural effusion, 1 postoperative hemorrhage

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Agcaoglu O 2013 ⁹²	Prospective cohort	America	HCC	1.7	RES	94	50/44	61.7±1.2	3.7±0.2	NA	0.53	0.53	2 pulmonary,2 biliary,2 wound-related,1 intestinal,1 hemorrhagic,2 cardiac, and 1 renal
					RFA	295	196/99	63.4 ±0.7	3.4±0.1	NA	0.2	0.2	3 bleeding,2 liver abscess,5 pulmonary,3 renal
Zhou Z 2014 ⁸⁹	Retrospective cohort	China	HCC	5	RES	21	15/6	42.2±7.6	1.7±0.3	0.81	NA	0.81	1 intraperitoneal hemorrhage
					RFA	31	20/11	46.7±9.8	1.7±0.4	0.81	NA	0.81	2 pleural effusion;2 fever;1 pneumonia;1 biloma
Kim JM 2014 ⁹¹	Retrospective cohort	Korea	HCC	2.8	RES	66	48/18	58.	2.1(0.8-3.0)	0.89	NA	0.89	NA
					RFA	67	52/15	59	1.8 (1.0-2.9)	0.49	NA	0.49	NA
Ko S 2014 ⁹⁰	Retrospective cohort	China	HCC	5	RES	12	9/3	71.6±4.3	2.9±1.4	NA	NA	0.67	NA
					RFA	17	9/8	57.3±3.6	2.3±1.1	NA	NA	0.35	NA
Kang TW 2015 ⁸⁸	Retrospective cohort	Korea	HCC	5	RES	142	107/35	53(28-74)	2(1.1–3.0)	0.90	NA	0.90	1 intra-abdominal abscess,3 wound problem,1 abdominal bleeding,1 intestinal obstruction

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	438	337/101	58(30-80)	1.9(1.1–3.0)	0.85	NA	0.85	3 tumor seeding,2 biloma,2 hepatic abscess,1 bile duct stricture,1 hepatic infarction
Lee YH 2015 ⁸⁷	Retrospect ive cohort	China	HCC	3.63	RES	330	261/69	61±12	<5	NA	NA	0.76	NA
					RFA	369	244/125	66±11	<5	NA	NA	0.66	NA
Liu PH 2016 ⁸³	Prospectiv e cohort	China	HCC	3.7	RES	109	78/31	60±13	<2	NA	0.81	0.81	NA
					RFA	128	84/44	64±12	<2	NA	0.76	0.76	NA
Hof J 2016 ⁸⁵	Retrospect ive cohort	Nethe rlands	HCC	3.2	RES	261	151/110	63.4	<5	0.69	NA	0.69	NA
					RFA	75	55/20	65.7	<5	NA	0.33(3y)	0.33(3y)	NA
Lee HW 2018 ⁸¹	RCT	Korea	HCC	5	RES	29	23/6	55.6±7.9	<5	NA	0.97(3y)	0.97(3y)	7 pleural effusion
					RFA	34	24/10	56.1±7.4	<5	NA	0.97(3y)	0.97(3y)	3 pain

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Li W 2017 ⁸²	Retrospect ive cohort	China	HCC	5	RES	220(239)	37/183	61.8 (40-73)	2.1 ±0.5	0.75	NA	0.75	64 complications
					MWA	60(61)	14/46	65(45-71)	2.0 ±0.5	0.67	NA	0.67	13 complications
Vogl TJ 2015 ⁸⁶	Retrospect ive cohort	Germ any	HCC	5	RFA	25(32)	19/6	57 ±3.5	3.2(0.8-4.5)	0.72(3y)	NA	0.72(3y)	NA
					MWA	28(36)	23/5	60 ±4.2	3.6(0.9-5)	0.79	NA	0.79(3y)	NA
Liu H 2016 ⁸⁶	RCT	China	HCC	4.7	TR	100(114)	86/14	52(31-80)	2.8(0.6-5)	0.67	NA	0.67	8 pleural effusion,5 biliary fstula,4 abdominal ascites,2 liver dysfunction,2 pneumonia,1 wound infection,1 abdominal infection
					RES	100(109)	94/6	49(30-76)	3(0.6-5)	0.84	NA	0.84	4 pleural effusion,3 liver dysfunction,3 abdominal ascites,1 abdominal bleeding

HCC: hepatocellular carcinoma;

BCLC: Barcelona Clinic Liver Cancer;

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

1 PEI: percutaneous ethanol injection;
2 RCT: randomized controlled trial;
3 NA: not available.
4
5
6

7 **Table S2.**
8 **Quality assessment of included studies using GRADE framework.**
9
10

Intervention/Comparator	Illustrative comparative risks* (per 1000, 95% CI)			Relative effect of survival time (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Comparator	Assumed survival risk	Corresponding survival risk with intervention			
1-year OS rate						
ES/MWA	923	984 (932 to 997)		OR 5.25 (1.15 to 23.97)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
EA/MWA	947	944 (902 to 968)		OR 0.94 (0.52 to 1.71)	990 (6 studies)	⊕ ⊕ ⊖ ⊖ low
ES/PEI	835	802 (674 to 889)		OR 0.80 (0.41 to 1.58)	519 (3 studies)	⊕ ⊕ ⊖ ⊖ low
EA/PEI	944	963 (906 to 1000)		OR 1.02 (0.96 to 1.09)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
ES/RFA	932	945 (931 to 956)		OR 1.25 (0.99 to 1.60)	5006 (30 studies)	⊕ ⊕ ⊕ ⊕ high
ES/TR	939	904 (765 to 965)		OR 0.61 (0.21 to 1.79)	201 (2 studies)	⊕ ⊕ ⊖ ⊖ low
EA/TR	938	802 (310 to 978)		OR 0.27 (0.03 to 2.90)	31 (1 study)	⊕ ⊕ ⊖ ⊖ low

RES/MWA	712	734 (623 to 822)	OR 1.12 (0.67 to 1.87)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
2					
3					
REA/MWA	736	779 (717 to 828)	OR 1.26 (0.91 to 1.73)	987 (6 studies)	⊕ ⊕ ⊖ ⊖ low
4					
5					
6					
RES/PEI	499	536 (421 to 645)	OR 1.16 (0.73 to 1.83)	519 (3 studies)	⊕ ⊕ ⊖ ⊖ low
7					
8					
9					
10					
RFA/PEI	729	748 (657 to 822)	OR 1.10 (0.71 to 1.71)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
11					
12					
13					
RES/RFA	785	851 (823 to 875)	OR 1.57 (1.28 to 1.93)	15906 (30 studies)	⊕ ⊕ ⊕ ⊖ moderate
14					
15					
16					
RES/TR	798	760 (618 to 860)	OR 0.80 (0.41 to 1.55)	201 (2 studies)	⊕ ⊕ ⊖ ⊖ low
17					
18					
19					
20					
RFA/TR	737	611 (516 to 704)	OR 0.56 (0.38 to 0.85)	454 (4 studies)	⊕ ⊕ ⊕ ⊖ moderate
21					
22					
23					
24					
RES/MWA	545	607 (492 to 712)	OR 1.29 (0.81 to 2.07)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
25					
26					
27					
28					
29					
30					
RES/PEI	293	436 (334 to 545)	OR 1.87 (1.21 to 2.90)	519 (3 studies)	⊕ ⊕ ⊕ ⊖ moderate
31					
32					
33					
RFA/PEI	533	496 (368 to 624)	OR 0.86 (0.51 to 1.45)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
34					
35					
36					
RES/RFA	601	744 (705 to 779)	OR 1.93 (1.59 to 2.34)	15154 (25 studies)	⊕ ⊕ ⊕ ⊖ moderate
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					

RES/TR	290	419 (251 to 607)	OR 1.76 (0.82 to 3.78)	117 (1 study)	⊕ ⊕ ⊖ ⊖ low
RFA/TR	464	356 (222 to 523)	OR 0.64 (0.33 to 1.27)	139 (1 study)	⊕ ⊕ ⊕ ⊖ moderate

The absolute and relative risk of survival with treatments*. GRADE: Grading of Recommendations, Assessment, Development and Evaluation. *The results presented in the Table S1 were built around the assumption of a consistent relative effect. The implications of this effect for populations were considered at different baseline risks. Based on the assumed risks, corresponding risks after an intervention were calculated using the meta-analytic risk ratio.

Table S3.
Ranking treatments of 1-, 3-year and 5-year survival rate of the lesions < 3 cm, 3-5 cm and ≤ 5 cm in RCT.

Treatment	1-year			3-year			5-year		
	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank
< 3cm	13			11			5		
RES		2	2.86		1	1.52		1	1.42
RFA		3	3.13		3	2.58		2	2.46
MWA		1	1.04		NA	NA		NA	NA
TR		4	3.59		2	2.35		3	2.89
PEI		5	4.43		4	3.55		4	3.23
3-5cm	4			4			2		
RES		1	1.17		1	1.19		1	1.69
RFA		3	2.88		3	2.91		3	2.60
MWA		NA	NA		NA	NA		NA	NA
TR		2	1.94		2	1.90		2	1.71
PEI		NA	NA		NA	NA		NA	NA
All tumours (≤	20			16			7		

5cm)						
RES	3	2.53	1	1.85	1	1.62
RFA	4	3.94	4	3.62	3	2.87
MWA	1	1.67	3	2.88	NA	NA
TR	2	1.93	2	2.38	2	1.78
PEI	5	4.92	5	4.27	4	3.73

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection.

Table S4.

Ranking treatments of 1-, 3-year and 5-year survival rate of the lesions < 3 cm, 3-5 cm and ≤ 5 cm in all studies.

Treatment	1-year			3-year			5-year		
	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank
< 3cm	50			48			37		
RES		3	1.18		1	1.71		1	1.16
RFA		4	3.17		3	3.38		2	3.02
MWA		1	1.91		4	3.42		3	3.11
TR		2	2.63		2	2.73		4	3.61
PEI		5	4.62		5	3.76		5	4.11
3-5cm	19			18			12		
RES		1	1.12		1	1.04		1	1.93
RFA		4	3.54		4	3.58		3	3.18
MWA		3	3.45		3	3.50		4	3.43
TR		2	2.05		2	2.14		2	1.94
PEI		5	4.84		5	4.74		5	4.53

1	All tumours (≤ 5cm)	72	68	50		
2	RES	2	2.07	1	1.50	1.11
3	RFA	3	3.48	3	3.68	3.34
4	MWA	4	3.57	4	3.84	3.23
5	TR	1	1.19	2	1.82	3.05
6	PEI	5	4.70	5	4.16	4.28

RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection.

Table S5.
Survival rates (1-year, 3-year and 5-year) for small lesion (<3cm) treatment comparisons estimated by direct and network meta-analysis in RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.97 (0.42-1.98)	0.98 (0.77-1.26)
MWA vs RES	152 (1.44-505.80)	NA
TR vs RES	1.08 (0.15-3.78)	0.99(0.67-1.47)
PEI vs RES	0.64 (0.18-1.61)	1.03 (0.54-1.94)
MWA vs RFA	173.30 (1.90-537.40)	1.42 (0.63-3.19)
TR vs RFA	1.25 (0.16-4.64)	1.00 (0.56-1.80)
PEI vs RFA	0.67 (0.28-1.35)	0.97 (0.78-1.19)
TR vs MWA	0.15 (0-0.80)	NA
PEI vs MWA	0.08 (0-0.38)	NA
PEI vs TR	1.17 (0.11-4.66)	NA
3-year OS rate for treatment vs reference		

RFA vs RES	0.75 (0.41-1.31)	0.92 (0.71-1.19)
MWA vs RES	NA	NA
TR vs RES	1.17 (0.16-4.17)	0.80(0.52-1.22)
PEI vs RES	0.58 (0.29-1.16)	1.21 (0.59-2.15)
MWA vs RFA	NA	NA
TR vs RFA	1.54 (0.25-13.43)	1.01 (0.55-1.87)
PEI vs RFA	0.79 (0.45-1.39)	0.91 (0.71-1.17)
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	1.02 (0.14-3.56)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.72 (0.10-2.47)	0.93 (0.62-1.37)
MWA vs RES	NA	NA
TR vs RES	0.84 (0.03-4.18)	0.88(0.69-1.12)
PEI vs RES	0.50 (0.04-2.04)	0.55 (0.26-1.15)
MWA vs RFA	NA	NA
TR vs RFA	2.87 (0.04-13.43)	NA
PEI vs RFA	0.94 (0.08-3.97)	0.97 (0.66-1.40)
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	3.93 (0.03-19.61)	NA

Table S6.

Survival rates (1-year, 3-year and 5-year) for lesion (3-5cm) treatment comparisons estimated by direct and network meta-analysis in RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.25 (0-1.47)	0.89 (0.45-1.77)
MWA vs RES	NA	NA

1	TR vs RES	1.00 (0-5.0)	NA
2	PEI vs RES	NA	NA
3	MWA vs RFA	NA	NA
4	TR vs RFA	3.40 (0.64-11.93)	1.10 (0.78-1.55)
5	PEI vs RFA	NA	NA
6	TR vs MWA	NA	NA
7	PEI vs MWA	NA	NA
8	PEI vs TR	NA	NA
9	3-year OS rate for treatment vs reference		
10	RFA vs RES	0.24 (0-1.25)	0.70 (0.34-1.45)
11	MWA vs RES	NA	NA
12	TR vs RES	1.14 (0-6.20)	NA
13	PEI vs RES	NA	NA
14	MWA vs RFA	NA	NA
15	TR vs RFA	3.98 (0.71-15.22)	1.29 (0.87-1.89)
16	PEI vs RFA	NA	NA
17	TR vs MWA	NA	NA
18	PEI vs MWA	NA	NA
19	PEI vs TR	NA	NA
20	5-year OS rate for treatment vs reference		
21	RFA vs RES	1.05 (0.03-5.33)	0.71 (0.32-1.57)
22	MWA vs RES	NA	NA
23	TR vs RES	12.87 (0.02-44.43)	NA
24	PEI vs RES	NA	NA
25	MWA vs RFA	NA	NA
26	TR vs RFA	7.64 (0.14-42.49)	1.93 (0.53-7.06)
27	PEI vs RFA	NA	NA
28	TR vs MWA	NA	NA
29	PEI vs MWA	NA	NA
30	PEI vs TR	NA	NA
31			
32			
33			
34			
35			
36			
37			
38			
39			
40			
41			
42			
43			
44			
45			
46			

Table S7.

Survival rates (1-year, 3-year and 5-year) for lesion (≤ 5 cm) treatment comparisons estimated by direct and network meta-analysis in RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.57 (0.27-1.08)	0.96 (0.78-1.19)
MWA vs RES	2.01 (0.47-5.70)	0.98 (0.54-1.78)
TR vs RES	1.50 (0.48-3.67)	0.99 (0.67-1.47)
PEI vs RES	0.37 (0.13-0.82)	1.03 (0.54-1.94)
MWA vs RFA	3.84 (0.81-11.60)	1.42 (0.63-3.19)
TR vs RFA	2.69 (1.02-6.04)	1.09 (0.84-1.43)
PEI vs RFA	0.65 (0.33-1.13)	0.95 (0.80-1.14)
TR vs MWA	1.09 (0.16-3.50)	NA
PEI vs MWA	0.27 (0.05-0.84)	NA
PEI vs TR	0.29 (0.09-0.73)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.65 (0.31-1.29)	0.88 (0.71-1.10)
MWA vs RES	1.00 (0.16-3.30)	0.88 (0.39-1.98)
TR vs RES	0.98 (0.35-2.41)	0.80 (0.51-1.22)
PEI vs RES	0.55 (0.19-1.44)	1.12 (0.59-2.15)
MWA vs RFA	1.77 (0.22-6.24)	NA
TR vs RFA	1.56 (0.66-3.25)	1.20 (0.90-1.60)
PEI vs RFA	0.86 (0.39-1.79)	0.84 (0.66-1.07)
TR vs MWA	1.86 (0.21-7.59)	NA
PEI vs MWA	1.05 (0.12-4.56)	NA
PEI vs TR	0.64 (0.19-1.67)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.66 (0.20-1.62)	0.88 (0.65-1.18)
MWA vs RES	NA	NA

TR vs RES	1.35 (0.23-4.69)	0.80 (0.52-1.22)
PEI vs RES	0.41 (0.11-1.02)	0.55 (0.26-1.15)
MWA vs RFA	NA	NA
TR vs RFA	2.29 (0.41-7.61)	1.30 (0.70-2.41)
PEI vs RFA	0.74 (0.16-2.00)	0.97 (0.66-1.40)
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	0.53 (0.06-1.90)	NA

OR: odds ratio;
RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection;
NA: not available.

Table S8.
Survival rates (1-year, 3-year and 5-year) for small lesion (<3cm) treatment comparisons estimated by direct and network meta-analysis in all studies.

Intervention	OR (95%CI)	
	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.94 (0.39-1.91)	1.00(0.95-1.04)
MWA vs RES	1.49 (0.44-3.85)	1.02(0.72-1.43)
TR vs RES	1.30 (0.28-3.88)	1.01(0.74-1.39)
PEI vs RES	0.63 (0.22-1.44)	1.00 (0.93-1.07)
MWA vs RFA	1.59 (0.69-3.17)	1.02 (0.85-1.23)
TR vs RFA	1.48 (0.34-4.23)	1.00(0.56-1.80)
PEI vs RFA	0.68 (0.38-1.09)	0.99 (0.93-1.06)

TR vs MWA	1.08 (0.21-7.87)	NA
PEI vs MWA	0.49 (0.18-1.10)	NA
PEI vs TR	0.69 (0.14-2.13)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.72 (0.37-1.30)	0.94 (0.90-0.99)
MWA vs RES	0.73 (0.30-1.55)	0.95 (0.78-1.18)
TR vs RES	0.90 (0.31-2.10)	1.08 (0.64-1.33)
PEI vs RES	0.68 (0.30-1.39)	1.00 (0.71-1.40)
MWA vs RFA	1.02 (0.57-1.70)	1.00 (0.82-1.22)
TR vs RFA	1.31 (0.47-2.92)	1.01 (0.55-1.87)
PEI vs RFA	0.96 (0.59-1.50)	0.97 (0.90-1.03)
TR vs MWA	1.38 (0.42-3.40)	NA
PEI vs MWA	1.01 (0.47-1.95)	NA
PEI vs TR	0.90 (0.29-2.17)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.54 (0.24-1.05)	0.85 (0.81-0.90)
MWA vs RES	0.55 (0.19-1.25)	0.88 (0.61-1.30)
TR vs RES	0.49 (0.16-0.18)	0.77 (0.53-1.11)
PEI vs RES	0.43 (0.17-0.89)	0.79 (0.73-0.85)
MWA vs RFA	1.04 (0.50-1.77)	1.02 (0.78-1.33)
TR vs RFA	0.99 (0.32-2.39)	NA
PEI vs RFA	0.82 (0.48-1.29)	0.92 (0.85-0.99)
TR vs MWA	1.03 (0.28-2.73)	NA
PEI vs MWA	0.86 (0.39-1.65)	NA
PEI vs TR	1.07 (0.31-2.72)	NA

Table S9.

Survival rates (1-year, 3-year and 5-year) for lesion (3-5cm) treatment comparisons estimated by direct and network meta-analysis in all studies.

Intervention	OR (95%CI)
--------------	------------

	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.12 (0-0.63)	0.96 (0.81-1.14)
MWA vs RES	0.15 (0-1.00)	NA
TR vs RES	0.36 (0.01-2.08)	1.02 (0.55-1.88)
PEI vs RES	0.06 (0-0.31)	NA
MWA vs RFA	1.29 (0.32-3.60)	0.99 (0.60-1.64)
TR vs RFA	2.99 (1.14-6.58)	1.11 (0.80-1.54)
PEI vs RFA	0.49 (0.18-1.12)	0.89 (0.66-1.20)
TR vs MWA	3.39 (0.58-10.44)	NA
PEI vs MWA	0.55 (0.09-1.76)	NA
PEI vs TR	0.20 (0.05-0.54)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.11 (0.01-0.40)	0.72 (0.60-0.88)
MWA vs RES	0.12 (0.01-0.53)	1.02 (0.57-1.81)
TR vs RES	0.26 (0.01-1.10)	0.92 (0.48-1.75)
PEI vs RES	0.06 (0-0.28)	NA
MWA vs RFA	1.15 (0.39-2.65)	0.81 (0.45-1.43)
TR vs RFA	2.38 (0.93-5.38)	1.29 (0.87-1.89)
PEI vs RFA	0.55 (0.12-1.69)	0.71 (0.50-1.00)
TR vs MWA	2.62 (0.61-7.90)	NA
PEI vs MWA	0.61 (0.08-2.26)	NA
PEI vs TR	0.28 (0.04-0.96)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.69 (0.04-3.16)	0.53 (0.40-0.68)
MWA vs RES	1.24 (0.02-4.46)	0.90 (0.48-1.69)
TR vs RES	14.31 (0.04-21.06)	NA
PEI vs RES	3.02 (0.01-2.40)	NA
MWA vs RFA	1.26 (0.19-4.04)	0.57 (0.21-1.51)
TR vs RFA	6.16 (0.27-25.58)	2.36 (0.66-8.37)
PEI vs RFA	0.86 (0.06-2.68)	0.56 (0.37-0.84)

TR vs MWA	11.97 (0.19-46.76)	NA
PEI vs MWA	4.15 (0.04-5.18)	NA
PEI vs TR	5.77 (0.01-2.84)	NA

Table S10.

Survival rates (1-year, 3-year and 5-year) for lesion (≤ 5 cm) treatment comparisons estimated by direct, indirect and network meta-analysis in all studies.

Intervention	OR (95%CI)	
	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.68 (0.35-1.17)	0.99 (0.95-1.04)
MWA vs RES	0.70 (0.29-1.39)	0.97 (0.77-1.23)
TR vs RES	1.72 (0.66-3.70)	1.01 (0.76-1.33)
PEI vs RES	0.52 (0.24-0.96)	1.01 (0.74-1.39)
MWA vs RFA	1.04 (0.55-1.76)	1.01 (0.85-1.20)
TR vs RFA	2.55 (1.20-4.85)	1.10 (0.85-1.43)
PEI vs RFA	0.77 (0.51-1.10)	0.98 (0.93-1.05)
TR vs MWA	2.69 (0.99-6.00)	0.91 (0.70-1.18)
PEI vs MWA	0.81 (0.38-1.51)	NA
PEI vs TR	0.34 (0.11-0.63)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.63 (0.37-1.01)	0.96 (0.94-0.98)
MWA vs RES	0.62 (0.32-1.09)	0.94 (0.72-1.22)
TR vs RES	0.97 (0.48-1.79)	0.92 (0.68-1.24)
PEI vs RES	0.59 (0.30-1.04)	0.93 (0.86-1.00)
MWA vs RFA	0.99 (0.64-1.47)	1.05 (0.86-1.26)
TR vs RFA	1.57 (0.89-2.57)	1.20 (0.90-1.60)
PEI vs RFA	0.94 (0.64-1.34)	0.95 (0.89-1.01)
TR vs MWA	1.65 (0.80-3.03)	NA
PEI vs MWA	0.98 (0.55-1.65)	NA

1	PEI vs TR	0.64 (0.32-1.16)	NA
2	5-year OS rate for treatment vs reference		
3	RFA vs RES	0.52 (0.29-0.88)	0.84 (0.80-0.88)
4	MWA vs RES	0.55 (0.25-1.05)	0.93(0.78-1.12)
5	TR vs RES	0.59 (0.25-1.20)	0.69 (0.34-1.42)
6	PEI vs RES	0.45 (0.23-0.82)	0.79 (0.73-0.85)
7	MWA vs RFA	1.06 (0.64-1.61)	0.97 (0.75-1.25)
8	TR vs RFA	1.16 (0.54-2.21)	1.30 (0.70-2.41)
9	PEI vs RFA	0.87 (0.57-1.26)	0.91 (0.84-0.98)
10	TR vs MWA	1.16(0.46-2.46)	NA
11	PEI vs MWA	0.87 (0.46-1.51)	NA
12	PEI vs TR	0.84 (0.35-1.74)	NA

OR: odds ratio;
RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection;
NA: not available.

Table S11.
Posterior summaries from random effects consistency and inconsistency models for small lesion (<3cm) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.55	0.21	(0.15-1.00)	0.38	0.23	(0.02-0.88)
τ	12.40	65.04	(1.10-45.68)	109.40	620.40	(1.30-940.00)
resdev	90.04	13.04	(66.16-117.10)	94.65	12.94	(70.06-120.70)

pD	66.48			57.5		
DIC	453.18			404.59		
3-year OS rate for treatment vs reference						
σ	0.59	0.14	(0.34-0.88)	0.6	0.14	(0.36-0.91)
τ	3.26	1.62	(1.34-7.33)	3.28	1.90	(1.19-8.10)
resdev	92.02	14.19	(66.64-122.10)	90.7	13.92	(65.64-120.00)
pD	80.45			71.83		
DIC	589.01			517.44		
5-year OS rate for treatment vs reference						
σ	0.53	0.12	(0.32-0.80)	0.55	0.13	(0.34-0.84)
τ	4.06	2.02	(1.66-8.76)	3.80	2.05	(1.40-8.77)
resdev	63.99	11.47	(43.52-88.24)	63.55	11.37	(43.39-87.90)
pD	64.22			55.07		
DIC	488.23			412.10		

Table S12.

Posterior summaries from random effects consistency and inconsistency models for lesion (3-5cm) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency Model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.28	0.25	(0.01-0.92)	0.38	0.34	(0.02-1.28)
τ	3108.00	68630.00	(1.44-4879.00)	19500.00	720600.00	(0.62-4178.00)
resdev	28.90	6.96	(17.25-44.41)	484.70	5117	(0.63-2616)
pD	24.70			24.62		
DIC	166.90			157.30		
3-year OS rate for treatment vs reference						
σ	0.62	0.27	(0.17-1.24)	0.67	0.31	(0.14-1.40)
τ	5.34	12.61	(0.83-21.20)	41.87	585.80	(0.52-77.13)
resdev	32.36	8.17	(18.39-50.07)	32.62	8.22	(18.52-50.51)

pD	30.91			28.63		
DIC	212.30			188.69		
5-year OS rate for treatment vs reference						
σ	0.80	0.46	(0.14-1.94)	0.60	0.42	(0.04-1.64)
τ	337.00	11980	(0.30-20.22)	10100.00	258400.00	(0.37-691.30)
resdev	22.54	6.73	(11.29-37.43)	22.57	6.519	(11.45-36.90)
pD	22.61			19.88		
DIC	146.84			131.53		

Table S13.
Posterior summaries from random effects consistency and inconsistency models for lesion (≤ 5 cm) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency Model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.49	0.13	(0.26-0.77)	0.29	0.14	(0.05-0.58)
τ	6.00	6.24	(1.92-16.85)	116.80	1122.00	(2.96-419.40)
resdev	129.2	14.99	(101.40-160)	133.1	14.50	(105.70-162.80)
pD	95.71			78.20		
DIC	692.39			604.18		
3-year OS rate for treatment vs reference						
σ	0.50	0.09	(0.33-0.70)	0.47	0.096	(0.29-0.67)
τ	4.20	1.45	(2.15-7.71)	5.31	2.59	(2.24-11.80)
resdev	124	15.64	(95.16-156.40)	124.5	15.89	(95.35-157.50)
pD	111.54			93.41		
DIC	856.01			723.74		
5-year OS rate for treatment vs reference						
σ	0.44	0.10	(0.26-0.65)	0.44	0.1	(0.26-0.67)
τ	5.30	2.27	(2.38-14.90)	6.09	3.95	(2.29-14.87)

resdev	86.73	13.53	(62.35-115.40)	85.74	13.55	(61.39-114.40)
pD	84.53			68.81		
DIC	670.73			544.40		

sd: standard deviation;

CI: Credible Interval

σ : between-trial standard deviation

τ^2 : between-trial variance

resdev: residual deviance

pD: effective number of parameters

DIC: deviance information criterion




Figure S1.

Results of the consistency test for closed loop at 1-year, 3-year, and 5-year survival rate of the lesions < 3 cm, 3-5 cm and \leq 5 cm.

- i Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions < 3 cm

- 1 ii Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions 3-5 cm
- 2
- 3 iii Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions ≤ 5 cm
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46


For peer review only



1 **Figure S2.**

2
3 **Assessment of publication bias using funnel plot.**

- 4
5
6 i Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions < 3 cm.
7
8
9 ii Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions 3-5 cm.
10
11 iii Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions \leq 5 cm
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

For peer review only

PRISMA NMA Checklist of Items to Include When Reporting A Systematic Review Involving a Network Meta-analysis

Section/Topic	Item #	Checklist Item	Reported on Page #
TITLE			
Title	1	Identify the report as a systematic review <i>incorporating a network meta-analysis (or related form of meta-analysis)</i> .	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: Background: main objectives Methods: data sources; study eligibility criteria, participants, and interventions; study appraisal; and <i>synthesis methods, such as network meta-analysis</i> . Results: number of studies and participants identified; summary estimates with corresponding confidence/credible intervals; <i>treatment rankings may also be discussed. Authors may choose to summarize pairwise comparisons against a chosen treatment included in their analyses for brevity.</i> Discussion/Conclusions: limitations; conclusions and implications of findings. Other: primary source of funding; systematic review registration number with registry name.	5,6
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known, <i>including mention of why a network meta-analysis has been conducted.</i>	7,8
Objectives	4	Provide an explicit statement of questions being addressed, with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	8

METHODS

Protocol and registration	5	Indicate whether a review protocol exists and if and where it can be accessed (e.g., Web address); and, if available, provide registration information, including registration number.	8,9
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale. <i>Clearly describe eligible treatments included in the treatment network, and note whether any have been clustered or merged into the same node (with justification).</i>	9,10
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	9
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	9,10,Figure1, Additional file 1: Text S1
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	9,10
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	10
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	11
Geometry of the network	S1	Describe methods used to explore the geometry of the treatment network under study and potential biases related to it. This should include how the evidence base has been graphically summarized for presentation, and what characteristics were compiled and used to describe the evidence base to readers.	11
Risk of bias within individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	11,12

Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means). <i>Also describe the use of additional summary measures assessed, such as treatment rankings and surface under the cumulative ranking curve (SUCRA) values, as well as modified approaches used to present summary findings from meta-analyses.</i>	11,12
Planned methods of analysis	14	Describe the methods of handling data and combining results of studies for each network meta-analysis. This should include, but not be limited to: <ul style="list-style-type: none"> • <i>Handling of multi-arm trials;</i> • <i>Selection of variance structure;</i> • <i>Selection of prior distributions in Bayesian analyses; and</i> • <i>Assessment of model fit.</i> 	11,12
Assessment of Inconsistency	S2	Describe the statistical methods used to evaluate the agreement of direct and indirect evidence in the treatment network(s) studied. Describe efforts taken to address its presence when found.	10,11,12
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	10,11,12
Additional analyses	16	Describe methods of additional analyses if done, indicating which were pre-specified. This may include, but not be limited to, the following: <ul style="list-style-type: none"> • Sensitivity or subgroup analyses; • Meta-regression analyses; • <i>Alternative formulations of the treatment network; and</i> • <i>Use of alternative prior distributions for Bayesian analyses (if applicable).</i> 	11,12

RESULTS†

Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	11,12
Presentation of network structure	S3	Provide a network graph of the included studies to enable visualization of the geometry of the treatment network.	12,13,Figure2-3
Summary of network geometry	S4	Provide a brief overview of characteristics of the treatment network. This may include commentary on the abundance of trials and randomized patients for the different interventions and pairwise comparisons in the network, gaps of evidence in the treatment network, and potential biases reflected by the network structure.	12,13,
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	11,12, Table1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment.	
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: 1) simple summary data for each intervention group, and 2) effect estimates and confidence intervals. <i>Modified approaches may be needed to deal with information from larger networks.</i>	12,13, Figure2-5
Synthesis of results	21	Present results of each meta-analysis done, including confidence/credible intervals. <i>In larger networks, authors may focus on comparisons versus a particular comparator (e.g. placebo or standard care), with full findings presented in an appendix. League tables and forest plots may be considered to summarize pairwise comparisons.</i> If additional summary measures were explored (such as treatment rankings), these should also be presented.	12,13,Figure4-5, Additional file 1: Table S1-S13
Exploration for inconsistency	S5	Describe results from investigations of inconsistency. This may include such information as measures of model fit to compare consistency and inconsistency models, <i>P</i> values from statistical tests, or summary of inconsistency estimates from different parts of the treatment network.	12,13

Risk of bias across studies	22	Present results of any assessment of risk of bias across studies for the evidence base being studied.	12,13, Additional file 1: Figure S1-S2
Results of additional analyses	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression analyses, <i>alternative network geometries studied, alternative choice of prior distributions for Bayesian analyses</i> , and so forth).	12,13
DISCUSSION			
Summary of evidence	24	Summarize the main findings, including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy-makers).	14-16
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review level (e.g., incomplete retrieval of identified research, reporting bias). <i>Comment on the validity of the assumptions, such as transitivity and consistency. Comment on any concerns regarding network geometry (e.g., avoidance of certain comparisons).</i>	16
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	17
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. This should also include information regarding whether funding has been received from manufacturers of treatments in the network and/or whether some of the authors are content experts with professional conflicts of interest that could affect use of treatments in the network.	17

PICOS = population, intervention, comparators, outcomes, study design.

* Text in italics indicates wording specific to reporting of network meta-analyses that has been added to guidance from the PRISMA statement.

† Authors may wish to plan for use of appendices to present all relevant information in full detail for items in this section.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

For peer review only

BMJ Open

Comparative efficacy of treatment strategies for hepatocellular carcinoma: systematic review and network meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-021269.R3
Article Type:	Research
Date Submitted by the Author:	28-Jun-2018
Complete List of Authors:	<p>Tian, Guo; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China.</p> <p>Yang, Shigui; The State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, School of Medicine, Zhejiang University, The Key Laboratory of Infectious Diseases</p> <p>Yuan, Jinqiu; The Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong, Division of Epidemiology</p> <p>Threapleton, Diane; Chinese University of Hong Kong, School of Public Health & Primary Care</p> <p>Zhao, Qiyu ; Department of Ultrasound Medicine, the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China, Department of Ultrasound Medicine</p> <p>Chen, Fen ; Department of Hepatobiliary Pancreatic Surgery, First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China., Department of Hepatobiliary Pancreatic Surgery</p> <p>Cao, Hongcui ; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University</p> <p>Jiang, Tian'an; First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China., Department of Ultrasonography</p> <p>Li, Lanjuan; State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University,</p>
Primary Subject Heading:	Oncology
Secondary Subject Heading:	Palliative care, Radiology and imaging, Surgery
Keywords:	resection, radiofrequency ablation, microwave ablation, transcatheter arterial chemoembolization, percutaneous ethanol injection, hepatocellular carcinoma

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHOLARONE™
Manuscripts

For peer review only

**Comparative efficacy of treatment strategies for hepatocellular carcinoma:
systematic review and network meta-analysis**

Guo Tian MD ^{1*}, Shigui Yang PhD ^{1*}, Jinqiu Yuan PhD ^{2,3}, Diane Threapleton PhD ²,
Qiyu Zhao PhD ⁴, Fen Chen PhD ⁵, Hongcui Cao PhD ^{1†}, Tian'an Jiang PhD ⁴, Lanjuan
Li PhD ^{1†}

*Shigui Yang was noted as the co-first author in the original article.

¹State Key Laboratory for Diagnosis and Treatment of Infectious Diseases,
Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases,
The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou
310003, China.

²Division of Epidemiology, The Jockey Club School of Public Health and Primary
Care, The Chinese University of Hong Kong, Hong Kong;

³Shenzhen Municipal Key Laboratory for Health Risk Analysis, Shenzhen Research
Institute of The Chinese University of Hong Kong, Shenzhen, Guangdong Province,
China

⁴Department of Ultrasonography, First Affiliated Hospital, Zhejiang University
School of Medicine, Hangzhou 310003, China.

⁵Department of Hepatobiliary Pancreatic Surgery, First Affiliated Hospital, Zhejiang
University School of Medicine, Hangzhou 310003, China.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Correspondence to: Lanjuan Li, ljli@zju.edu.cn;
Hongcui Cao, hccao@zju.edu.cn.

[†]Lanjuan Li, and Hongcui Cao contributed equally to this paper.

For peer review only

Electronic word count: 3036 (excluding Tables and References)

Number of figures and tables: 7 figures and 15 tables

List of abbreviations in order of appearance: HCC: hepatocellular carcinoma; RES: resection; RFA: radiofrequency ablation; MWA: microwave ablation; TACE: transcatheter arterial chemoembolization; PEI: percutaneous ethanol injection; GRADE: Grading of Recommendations Assessment, Development and Evaluation; OR: odds ratio; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses; TR: TACE plus RFA; OS: overall survival; MCMC: Markov Chain Monte Carlo; CrI: credible interval; SUCRA: surface under the cumulative ranking curve LPS: lipopolysaccharide; TNF α : tumor necrosis factor α ; IL: interleukin; TGF β : transforming growth factor β .

Conflict of interest: The authors have declared that no competing interests regarding the publication of this paper.

Data sharing statement: No additional data are available.

Financial support: This study was supported by the opening foundation of the State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital of Medical College, Zhejiang University, grant NO. 2015KF06; This study was supported by the Foundation of Zhejiang Health Committee (2017KY346).

Author Contributions:

1. Conceived and designed the experiments: Hongcui Cao, Tian'an Jiang, Lanjuan Li

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2. Performed the experiments: Guo Tian, Shigui Yang, Jinqiu Yuan, Diane Threapleton, Qiyu Zhao, Fen Chen, Tian'an Jiang
3. Analyzed the data: Guo Tian, Shigui Yang, Jinqiu Yuan, Qiyu Zhao
4. Contributed reagents/materials/analysis tools: Qiyu Zhao, Fen Chen
5. Wrote the manuscript: Guo Tian, Shigui Yang, Tian'an Jiang
6. Critically revised and approved the final version of manuscript: Diane Threapleton, Hongcui Cao, Tian'an Jiang, Lanjuan Li
7. Study supervision: Hongcui Cao, Tian'an Jiang, Lanjuan Li

Abstract

Objective: Hepatocellular carcinoma (HCC) is the 3rd leading cause of cancer death worldwide. We conducted network meta-regression within a bayesian framework to compare and rank different treatment strategies for HCC through direct and indirect evidence from international studies.

Methods and analyses: We pooled the odds ratio (OR) for 1-, 3- and 5-year overall survival, based on lesions of size < 3 cm, 3-5 cm and \leq 5 cm, using five therapeutic options including resection (RES), radiofrequency ablation (RFA), microwave ablation (MWA), transcatheter arterial chemoembolization (TACE) plus RFA (TR) and percutaneous ethanol injection (PEI).

Results: We identified 74 studies, including 26944 patients. After adjustment for study design, and in the full sample of studies, the treatments were ranked in order of greatest to least benefit as follows for 5-year survival: 1) RES, 2) TR, 3) RFA, 4) MWA, and 5) PEI. The ranks were similar for 1 and 3-year survival, with RES and TR being the highest ranking treatments. In both smaller (<3cm) and larger tumors (3-5cm), RES and TR were also the two highest ranking treatments. There was little evidence of inconsistency between direct and indirect evidence.

Conclusion: The comparison of different treatment strategies for HCC indicated that RES is associated with longer survival. However, many of the between-treatment comparisons were not statistically significant and, for now, selection of strategies for treatment will depend patient and disease characteristics. Additionally, much of the evidence was provided by non randomised studies and knowledge gaps still exist.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

More head-to-head comparisons between both RES and TR, or other approaches, will be necessary to confirm these findings.

Key words: resection; radiofrequency ablation; microwave ablation; transcatheter arterial chemoembolization; percutaneous ethanol injection; hepatocellular carcinoma.

Strengths and limitations of this study:

1. This is a network meta-regression within a bayesian framework to compare and rank different treatment strategies for HCC through direct and indirect evidence from international studies.
2. Strong and reliable methodological and statistical procedures were applied.
3. The individual or tumor characteristics within HCC articles would be a source of heterogeneity..
4. A major limitation is in the inclusion of non-randomised studies, in which selection bias is likely to confound observations. Selection of treatment is likely to be based on individual or tumor characteristics, and thus these factors will bias and confound observations of survival.
5. Other studies did not report the primary outcome of interest (5-year survival) and this was a particular limitation among randomised studies.

Introduction

Cancer was the second leading cause of death in 2013, behind cardiovascular disease, and in 2013 more than 8 million people died from cancer globally ¹⁻³. Hepatocellular carcinoma (HCC) was the 6th most common cancer worldwide and the 3rd leading cause of cancer death, with 5-year overall survival rates under 12% ^{4,5}.

Hepatic resection (RES) was the traditional choice for patients with HCC, without cirrhosis and with good remaining liver function ⁶. Despite nearly 70% 5-year survival, recurrence rates after surgery were high ⁷. Repeated hepatectomies to lengthen survival were not often appropriate owing to multiple-site tumor recurrence or patient background of liver cirrhosis ^{8,9}. Many locoregional therapies have been developed including ablative treatments such as percutaneous ethanol injection (PEI), radiofrequency ablation (RFA), or microwave ablation (MWA), and trans-arterial therapies such as transcatheter arterial chemoembolization (TACE) or transarterial chemotherapy infusion (TACI). Locoregional therapies were minimally invasive and therefore are cheaper and faster to recover, as compared to resection. Such approaches may be appropriate for patients with unresectable, small or multiple carcinomas or those with severe cirrhosis. However, there may be a greater risk of recurrence because of incomplete destruction of cancer cells at the treatment margin, as seen with RFA ¹⁰.

Selection of treatment strategy was determined by liver function, tumor stage and patient performance status ⁷, but much uncertainty still remains surrounding the comparative efficacy of different treatment approaches. A recent review of

international guidelines for HCC found similarities but also some discrepancy in treatment allocation recommendations because of regional classification differences, secondary to a lack of solid or high-level evidence ¹¹. A recent review of therapies also revealed that there was no consensus on whether surgery or ablation was better for small tumors ⁷. Some discrepancy in prevalence and treatment outcomes may be still in different regions because of local biology, available resources or expertise and access to care ¹¹. However, if we ever hope to achieve standardized and evidence-based therapy for HCC, the unanswered question surrounding relative treatment efficacy of RES compared to ablative locoregional therapies should be resolved.

Traditional meta-analysis is limited by existing head-to-head treatment comparisons within included studies. It is therefore not possible to gauge the relative benefit of two treatments that have never been directly compared in studies. Real-life treatment decisions are hindered by gaps in existing evidence, but network meta-analysis enables integration of direct and indirect comparisons to provide estimates for relative comparisons across many treatments ¹². Recent published network meta-analysis focused on advanced HCC by TACE alone or combined treatments^{13 14}, as well as antineoplastic drugs (sorafenib, erlotinib, linifanib, sunitinib and brivanib)¹⁵, and early- or very early-stage HCC via surgery or thermal ablation¹⁶. However, in this study, we included the latest literature, and focused on the comparison of interventional and surgical treatments, including RES, RFA, MWA, and TACE plus RFA (TR), PEI using subgroup analysis of tumor size (smaller: <3cm;

larger: 3-5cm), and study design (cohort or RCT). In order to investigate comparative effectiveness among RES and common locoregional ablative therapies, we performed a strong and reliable bayesian network meta-analysis.

Search Strategy

We conducted a systematic review and report findings in accordance with PRISMA for Network Meta-Analyses (PRISMA-NMA)¹⁷ (Additional file 1: Text S1). The following databases were searched: PubMed, Embase, Web of science and Scopus, up to May 2018, using these keywords: resection, surgery, hepatectomy, radiofrequency ablation, transarterial chemoembolization, microwave thermal ablation, ethanol injection, liver, cancer, tumor (Additional file 1: Text S2). No language restrictions were used. Bibliographies from other relevant review articles were cross-examined for potential missed studies. Disagreement was resolved by a third reviewer. Citations were downloaded into reference management software and duplicate citations were electronically or manually removed.

We systematically included the studies using the following criteria: 1) original data from prospective or retrospective cohort studies and randomized clinical trials (RCTs) in humans; 2) reporting at least two treatments, including resection or any local ablative therapy (RES, RFA, MWA, PEI, or TACE+RFA (TR)); 3) mean lesion size ≤ 5 cm; 4) evaluating overall survival rate not less than one year after first or recurrent treatments. Conference abstracts and case reports were excluded, as were older publications from studies with multiple publications.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Patients and public involvement

The patients or public were not involved in the study.

Data Extraction and Study Quality

Two investigators independently extracted and cross-checked the data from the eligible studies: author, year, study design, country, disease type, inclusion criteria, treatment style, study size, gender, age, tumor size, follow-up duration, treatment complications and survival outcomes. If in disagreement, a third reviewer adjudicated. The level of evidence was appraised using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) guidance ¹⁸, which was classified into four levels of high, moderate, low, and very low. The quality score was downgraded according to 5 domains, including risk of bias, inconsistency, indirectness, imprecision, and publication bias while scores were upgraded according to large effect, appropriate control for plausible confounding, and dose-response gradient.

Data Analysis

Network meta-analysis was used if a ring or open evidence loop was available to know the number of arms and the sample size of each intervention. When possible, pair-wise direct head-to-head comparisons were conducted to calculate the odds ratio (OR) of 1-, 3- and 5-year survival and their 95% confidence intervals (CI).

Between-study heterogeneity was evaluated using the tau-squared statistic (τ^2)¹⁹. A node-splitting analysis was applied to check the consistency between direct evidence (existing real reported comparisons) and indirect evidence (estimated treatment comparisons) for their agreement on a specific node²⁰. Bayesian network meta-analysis with Markov Chain Monte Carlo (MCMC), through a consistency model, was utilized to estimate the pooled ORs and its 95% credible interval (CrI) for the direct and indirect comparisons¹⁶. The inconsistency model was used to check for heterogeneity due to chance imbalance in the distribution of effect modifiers. Consistency in every closed loop was checked by the loop-specific approach in order to estimate whether treatment survival effects were disturbed by variance in the distribution of potential confounding factors among the studies. In order to compare and rank survival rates of different treatments, we examined all studies first and then separately assessed smaller (<3cm) and larger (3-5cm) tumors. Random-effect meta-regression models were used, with and without adjustment for study design (cohort or RCT) and subgroup analyses were also conducted for RCTs in order to examine treatment effectiveness. We appraised the ranking probabilities for all therapies for each intervention and the treatment hierarchy was ordered by the surface under the cumulative ranking curve (SUCRA)²¹. Sensitivity analysis was conducted to remove each study, in turn, and estimate the treatment effect in the remaining studies. Funnel plots were utilized to check the possible presence of publication bias or small-study bias²². In this study, we used Bayesian MCMC simulations by WinBUGS 1.4 and graphically presented the results using Stata 13.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Study Characteristics

After screening, 74 relevant studies in 73 articles were identified, of which 20 were randomized controlled trials and 54 were cohort studies²³⁻⁹⁶. We excluded 136504 duplicate or non-relevant citations (Figure 1). The summary characteristics of these studies are shown in Additional file 1: Table S1. Overall, 32345 patients of mean age from 46 to 73.5 years, with approximately 29236 tumors, were assigned to receive RES, RFA, MWA, TR and PEI, and the mean follow-up ranged from 1.5 to 5.7 years. In addition, the numbers of connected studies to the lines (black) and sample size of each treatment (red) were shown in Figure 2 and 3, respectively.

Network Meta-Analysis Results

Ten possible treatment comparisons among the five interventions were examined in the included studies. Comparable survival estimates were made for each treatment (per 1000 patients) and the survival OR among each of the treatment comparisons, according to follow-up duration, are presented in Additional file 1: Table S2, along with estimation of the quality of evidence using GRADE criteria.

Across the range of treatment comparisons and follow-up durations, evidence was graded between low and high quality. Evidence was often graded as low quality owing to publication bias and graded as high quality owing to a larger number of participants in direct comparisons.

Survival probabilities (estimated using Meanrank) and ranks for the five treatments in patients with tumors $<3\text{cm}$, $3\text{-}5\text{cm}$ or $\leq 5\text{cm}$ (with and without adjustment for study design) are graphically displayed in Figures 2-5, and numerical details are given in Additional file 1: Table S3-S4. RES was consistently associated with greater survival (rank 1) compared to MWA, RFA, TR and PEI for the 5-year survival estimates. The ranks were similar for 1 and 3-year survival with RES or TR being ranked as 1 or 2 in most analyses. After adjustment for study design, and in the full sample of available studies ($n=74$), the treatments were ranked as follows for 5-year survival: 1) RES, 2) TR, 3) RFA, 4) MWA, and 5) PEI (Table S4).

Efficacy comparisons from network meta-regression for all treatments are summarized in Table 1 and 2, according to follow-up duration and initial tumor size. Compared to RES, the 5-year survival in all studies (trials and observational studies) for all tumors $\leq 5\text{cm}$, was 0.45 (95%CrI 0.23 to 0.82) for PEI, 0.59 (95%CrI 0.25 to 1.20) for TR, 0.55 (95%CrI 0.25 to 1.05) for MWA and 0.52 (95%CrI 0.29 to 0.88) for RFA (Table 2). When examining the comparisons across all treatments, the only significant difference for tumors $<3\text{cm}$ was for 5-year survival, and a significantly worse survival was observed for PEI compared to RES 0.43 (95%CrI 0.17 to 0.89). For tumors between 3 and 5 cm, no significant differences were observed at 5-year survival, but significantly worse 3-year survival was observed with PEI, MWA and RFA compared to RES (Table 2). Despite smaller number of studies in analyses of only RCTs, the pairwise comparisons showed similar results. However, all relative rankings should be interpreted with caution because most network meta-regression

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

comparisons did not suggest a statistically significant difference between treatments. Detailed results of each comparison for survival rates were shown in Additional file 1: Table S5-S10.

Loop-specific methods detected no inconsistency between the pairwise and network meta-analysis for most closed loops in the network (Additional file 1: Figure S1). However, inconsistency was observed between direct and indirect comparisons for the following loops: lesions <3cm: RES-RFA-TR, PEI-RES-RFA, MWA-RES-RFA; lesions 3-5cm: MWA-RES-RFA, RES-RFA-TR; and lesions \leq 5cm: RES-RFA-TR). In addition, tests for inconsistency were carried out (Additional file 1: Table S11-S13), which indicated a close relationship of between-trial heterogeneity and inconsistency between “direct” and “indirect” evidence.

Sensitivity Analysis and publication bias

No significant change was observed when any one study was deleted. Funnel plots indicated that the included studies in each group were distributed symmetrically around the vertical line ($x=0$), suggesting that no obvious evidence of publication bias or small-sample effect existed in this network (Additional file 1: Figure S2).

Discussion

There were many techniques for attaining a large ablated zone and complete necrosis of HCC and this comprehensive review addressed two of the more common treatments, namely resection and ablation. In this network meta-analysis, of the five examined therapies, the pooled data showed RES ranked best in full sample analysis

with or without adjustment for study design. In both smaller (<3cm) and larger tumors (3-5cm) RES remained the highest ranking treatment. However, most of the individual treatment comparisons were not statistically significant and thus, RES may not be superior to all other therapies. Our evidence indicated locoregional therapies and particularly RES or TR (TACE+RFA) were associated with longer survival.

Our observation of better survival outcomes with TR may be through the advantage of dual mechanisms. With TR, TACE induced hypoxic injury on cancer cells through occlusion of blood vessels and was followed by local ablation. This combination therapy may result in a larger ablated zone⁹⁷, reducing the possibility of micrometastasis and recurrence, and thus, resulting in better survival outcomes than RFA alone.

While being more invasive, and despite risk of complications, RES was associated with better survival outcomes after 1 year, 3 years and 5 years. This may be due to removal of larger sections of liver than can be targeted with locoregional therapies, thus removing a larger area of potentially cancerous cells. Additionally, rat models indicated that the liver has the potential to quickly restore its original size after partial hepatectomy. This may be mediated via interactions of lipopolysaccharide (LPS), tumor necrosis factor (TNF) α , interleukin (IL)-6, and transforming growth factor β (TGF β)⁹⁸. However, evidence from rat models and human studies indicated that resection success was associated with resection size and regeneration was stunted with larger resections⁹⁹⁻¹⁰¹. The safe limit for remnant liver volume in normal liver was approximately 30% of total liver volume, but this was estimated to rise to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

40-50% in those with liver disease^{99 102}. Liver resection was recognised as the most efficient treatment for HCC but was only applicable for less than 30% of all patients. However, developments in preoperative imaging techniques, laproscopic surgery and newly developing combinations with chemotherapy may extend its application to more advanced tumors¹⁰². Furthermore, the consistent associations observed with all studies and only in RCTs indicated that patient selection bias in the observational studies does not wholly explain the better survival outcomes with RES.

Overall, we found PEI was associated with shorter survival than the other four therapies, a finding which is supported in previous studies^{24 33}. One study reported RFA was superior to PEI in achieving short- and long-term survival outcomes, although PEI and RFA showed similar 5-year survival in lesions <3 cm⁵⁵. The possible reason why PEI is less effective than RFA may be because lesions often have a thick capsule and therefore ethanol may not distribute through tissues.

There are several limitations in this study. Firstly, a major limitation is in the inclusion of non-randomised studies, in which selection bias is likely to confound observations. Selection of treatment is likely to be based on individual or tumor characteristics, and thus these factors will bias and confound observations of survival. Secondly, this study included both RCTs and observational studies, in which study designs and type of data collection may not be comparable. However, findings were consistent among both study designs. Thirdly, all included studies did not report our primary outcome of interest (5-year survival) and this was a particular limitation among randomised studies. Fourthly, for many individual comparisons, there were

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

either no direct comparisons or comparisons from only a small number of studies. The lack of evidence may increase the risk of bias, which could enlarge or undervalue effect size, and may explain the small inconsistency seen between direct and estimated comparisons. Thus, we should be cautious in interpreting treatment rankings for the different survival times and for different size lesions. While adverse events from treatments may differ (not evaluated in detail in this review), by examining overall survival outcomes in our review, we have taken account of both long-term potential benefits and harms from treatments. The focus of these findings should therefore be on the overall observation that RES or TR may be superior in terms of survival, rather than focusing on specific OR values for individual treatment comparisons.

In conclusion, the findings of the current bayesian network meta-analysis indicate that RES or TR may be among the most effective therapeutic approaches for HCC for 5-year survival in both smaller ($< 3\text{cm}$) and larger ($3\text{-}5\text{cm}$) lesions. However, evidence was of variable quality, and the majority of evidence came from non randomised studies, which are prone to selection bias and knowledge gaps still exist. For not, at the individual level, selection of strategies should depend on patient and clinical characteristics. To facilitate generation of evidence-based recommendations for HCC therapy, and to standardize treatment approaches, further head-to-head comparisons, especially of resection and ablative therapies, are required from high-quality RCTs, with long follow-up for survival outcomes.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conflict of interests

The authors have declared that no competing interests regarding the publication of this paper.

Data sharing statement

No additional data are available.

Acknowledgments

This study was supported by the opening foundation of the State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital of Medical College , Zhejiang University, grant NO. 2015KF06; This study was supported by the Foundation of Zhejiang Health Committee (2017KY346).

References

1. Collaborators GMaCoD. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;385:117-71
2. Lozano R, Naghavi M, Foreman K, *et al.* Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095-128
3. Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: Global Burden of Disease Study. *Lancet* 1997;349:1269-76
4. El-Serag HB. Hepatocellular carcinoma. *The New England journal of medicine* 2011;365:1118-27
5. Ferlay J, Shin HR, Bray F, *et al.* Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *International journal of cancer* 2010;127:2893-917
6. Forner A, Llovet JM, Bruix J. Hepatocellular carcinoma. *Lancet* 2012;379:1245-55
7. Raza A, Sood GK. Hepatocellular carcinoma review: current treatment, and evidence-based medicine. *World journal of gastroenterology* 2014;20:4115-27
8. Kishi Y, Hasegawa K, Sugawara Y, *et al.* Hepatocellular carcinoma: current management and future development-improved outcomes with surgical resection. *International journal of hepatology* 2011;2011:728103
9. Mazzaferro V, Bhoori S, Sposito C, *et al.* Milan criteria in liver transplantation

- for hepatocellular carcinoma: an evidence-based analysis of 15 years of experience. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society* 2011;17 Suppl 2:S44-57
10. Shan, Xue-Mei, Ding, *et al.* Radiofrequency ablation of hepatocellular carcinoma sized > 3 and ≤ 5 cm: Is ablative margin of more than 1 cm justified? *World journal of gastroenterology* 2011;19:7389-98
11. Yu SJ. A concise review of updated guidelines regarding the management of hepatocellular carcinoma around the world: 2010-2016. *Clinical & Molecular Hepatology* 2016;22:7-17
12. Chaimani A, Higgins JP, Mavridis D, *et al.* Graphical tools for network meta-analysis in STATA. *PloS one* 2013;8:e76654
13. Xie H, Yu H, Tian S, *et al.* What is the best combination treatment with transarterial chemoembolization of unresectable hepatocellular carcinoma? a systematic review and network meta-analysis. *Oncotarget* 2017;8:100508
14. Katsanos K, Kitrou P, Spiliopoulos S, *et al.* Comparative effectiveness of different transarterial embolization therapies alone or in combination with local ablative or adjuvant systemic treatments for unresectable hepatocellular carcinoma: A network meta-analysis of randomized controlled trials. *PloS one* 2017;12:e0184597
15. Cucchetti A, Piscaglia F, Pinna AD, *et al.* Efficacy and Safety of Systemic Therapies for Advanced Hepatocellular Carcinoma: A Network Meta-Analysis

- of Phase III Trials. *Liver Cancer* 2017;6:337-48
16. Management of people with early or very early stage hepatocellular carcinoma: an attempted network meta-analysis. International Liver Congress / Meeting of the European-Association-For-The-Study-Of-The-Liver; 2017.
 17. Hutton B, Salanti G, Caldwell DM, *et al.* The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Annals of internal medicine* 2015;162:777-84
 18. Puhan MA, Schunemann HJ, Murad MH, *et al.* A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis. *Bmj* 2014;349:g5630
 19. Higgins JP, Green S. Cochrane Handbook for Systematic Reviews of Interventions. *Naunyn-Schmiedebergs Archiv für experimentelle Pathologie und Pharmakologie* 2009;2011:S38
 20. Dias S, Welton NJ, Caldwell DM, *et al.* Checking consistency in mixed treatment comparison meta-analysis. *Statistics in medicine* 2010;29:932-44
 21. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology* 2011;64:163-71
 22. Jin ZC, Zhou XH, He J. Statistical methods for dealing with publication bias in meta-analysis. *Statistics in medicine* 2015;34:343-60
 23. Zhang Z, Wu M, Chen H, *et al.* [Percutaneous radiofrequency ablation

combined with transcatheter arterial chemoembolization for hepatocellular carcinoma]. *Zhonghua wai ke za zhi [Chinese journal of surgery]* 2002;40:826-9

24. Lencioni RA, Allgaier HP, Cioni D, *et al.* Small hepatocellular carcinoma in cirrhosis: randomized comparison of radio-frequency thermal ablation versus percutaneous ethanol injection. *Radiology* 2003;228:235-40

25. Lin SM, Lin CJ, Lin CC, *et al.* Radiofrequency ablation improves prognosis compared with ethanol injection for hepatocellular carcinoma \leq 4 cm. *Gastroenterology* 2004;127:1714-23

26. Vivarelli M, Guglielmi A, Ruzzenente A, *et al.* Surgical resection versus percutaneous radiofrequency ablation in the treatment of hepatocellular carcinoma on cirrhotic liver. *Annals of surgery* 2004;240:102-7

27. Cho CM, Tak WY, Kweon YO, *et al.* [The comparative results of radiofrequency ablation versus surgical resection for the treatment of hepatocellular carcinoma]. *The Korean journal of hepatology* 2005;11:59-71

28. Hong SN, Lee SY, Choi MS, *et al.* Comparing the outcomes of radiofrequency ablation and surgery in patients with a single small hepatocellular carcinoma and well-preserved hepatic function. *Journal of clinical gastroenterology* 2005;39:247-52

29. Huang GT, Lee PH, Tsang YM, *et al.* Percutaneous ethanol injection versus surgical resection for the treatment of small hepatocellular carcinoma: a prospective study. *Annals of surgery* 2005;242:36-42

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
30. Lin SM, Lin CJ, Lin CC, *et al.* Randomised controlled trial comparing percutaneous radiofrequency thermal ablation, percutaneous ethanol injection, and percutaneous acetic acid injection to treat hepatocellular carcinoma of 3 cm or less. *Gut* 2005;54:1151-6
31. Lu MD, Xu HX, Xie XY, *et al.* Percutaneous microwave and radiofrequency ablation for hepatocellular carcinoma: a retrospective comparative study. *Journal of gastroenterology* 2005;40:1054-60
32. Montorsi M, Santambrogio R, Bianchi P, *et al.* Survival and recurrences after hepatic resection or radiofrequency for hepatocellular carcinoma in cirrhotic patients: a multivariate analysis. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2005;9:62-7; discussion 67-8
33. Shiina S, Teratani T, Obi S, *et al.* A randomized controlled trial of radiofrequency ablation with ethanol injection for small hepatocellular carcinoma. *Gastroenterology* 2005;129:122-30
34. Chen MS, Li JQ, Zheng Y, *et al.* A prospective randomized trial comparing percutaneous local ablative therapy and partial hepatectomy for small hepatocellular carcinoma. *Annals of surgery* 2006;243:321-8
35. Lu MD, Kuang M, Liang LJ, *et al.* [Surgical resection versus percutaneous thermal ablation for early-stage hepatocellular carcinoma: a randomized clinical trial]. *Zhonghua yi xue za zhi* 2006;86:801-5
36. Cho YB, Lee KU, Suh KS, *et al.* Hepatic resection compared to percutaneous

ethanol injection for small hepatocellular carcinoma using propensity score matching. *Journal of gastroenterology and hepatology* 2007;22:1643-9

37. Gao W, Chen MH, Yan K, *et al.* Therapeutic effect of radiofrequency ablation in unsuitable operative small hepatocellular carcinoma. *Chinese Journal of Medical Imaging Technology* 2007

38. Lupo L, Panzera P, Giannelli G, *et al.* Single hepatocellular carcinoma ranging from 3 to 5 cm: radiofrequency ablation or resection? *HPB : the official journal of the International Hepato Pancreato Biliary Association* 2007;9:429-34

39. Zhou T, Qiu YD, Kong WT. Comparing the effect of radiofrequency ablation and surgical resection for the treatment of small hepatocellular. *Journal of Hepatobiliary Surgery* 2007

40. Abu-Hilal M, Primrose JN, Casaril A, *et al.* Surgical resection versus radiofrequency ablation in the treatment of small unifocal hepatocellular carcinoma. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2008;12:1521-6

41. Brunello F, Veltri A, Carucci P, *et al.* Radiofrequency ablation versus ethanol injection for early hepatocellular carcinoma: A randomized controlled trial. *Scandinavian journal of gastroenterology* 2008;43:727-35

42. Guglielmi A, Ruzzenente A, Valdegamberi A, *et al.* Radiofrequency ablation versus surgical resection for the treatment of hepatocellular carcinoma in cirrhosis. *Journal of gastrointestinal surgery : official journal of the Society*

- for *Surgery of the Alimentary Tract* 2008;12:192-8
43. Hiraoka A, Horiike N, Yamashita Y, *et al.* Efficacy of radiofrequency ablation therapy compared to surgical resection in 164 patients in Japan with single hepatocellular carcinoma smaller than 3 cm, along with report of complications. *Hepato-gastroenterology* 2008;55:2171-4
44. Ohmoto K, Yoshioka N, Tomiyama Y, *et al.* Comparison of therapeutic effects between radiofrequency ablation and percutaneous microwave coagulation therapy for small hepatocellular carcinomas. *Journal of gastroenterology and hepatology* 2009;24:223-7
45. Sakaguchi H, Seki S, Tsuji K, *et al.* Endoscopic thermal ablation therapies for hepatocellular carcinoma: a multi-center study. *Hepatology research : the official journal of the Japan Society of Hepatology* 2009;39:47-52
46. Santambrogio R, Opocher E, Zuin M, *et al.* Surgical resection versus laparoscopic radiofrequency ablation in patients with hepatocellular carcinoma and Child-Pugh class a liver cirrhosis. *Annals of surgical oncology* 2009;16:3289-98
47. Shibata T, Isoda H, Hirokawa Y, *et al.* Small hepatocellular carcinoma: is radiofrequency ablation combined with transcatheter arterial chemoembolization more effective than radiofrequency ablation alone for treatment? *Radiology* 2009;252:905-13
48. Ueno S, Sakoda M, Kubo F, *et al.* Surgical resection versus radiofrequency ablation for small hepatocellular carcinomas within the Milan criteria. *Journal*

of hepato-biliary-pancreatic surgery 2009;16:359-66

49. Xiang-Yang BU, Wang Y, Zhong GE, *et al.* Comparison of radiofrequency ablation and surgical resection for small primary liver carcinoma. *Chinese Archives of General Surgery* 2009

50. Guo WX, Zhai B, Lai EC, *et al.* Percutaneous radiofrequency ablation versus partial hepatectomy for multicentric small hepatocellular carcinomas: a nonrandomized comparative study. *World journal of surgery* 2010;34:2671-6

51. Huang J, Yan L, Cheng Z, *et al.* A randomized trial comparing radiofrequency ablation and surgical resection for HCC conforming to the Milan criteria. *Annals of surgery* 2010;252:903-12

52. Kagawa T, Koizumi J, Kojima S, *et al.* Transcatheter arterial chemoembolization plus radiofrequency ablation therapy for early stage hepatocellular carcinoma: comparison with surgical resection. *Cancer* 2010;116:3638-44

53. Morimoto M, Numata K, Kondou M, *et al.* Midterm outcomes in patients with intermediate-sized hepatocellular carcinoma: a randomized controlled trial for determining the efficacy of radiofrequency ablation combined with transcatheter arterial chemoembolization. *Cancer* 2010;116:5452-60

54. Azab M, Zaki S, El-Shetey AG, *et al.* Radiofrequency ablation combined with percutaneous ethanol injection in patients with hepatocellular carcinoma. *Arab journal of gastroenterology : the official publication of the Pan-Arab Association of Gastroenterology* 2011;12:113-8

- 1
2
3
4 55. Giorgio A, Di Sarno A, De Stefano G, *et al.* Percutaneous radiofrequency
5
6 ablation of hepatocellular carcinoma compared to percutaneous ethanol
7
8 injection in treatment of cirrhotic patients: an Italian randomized controlled
9
10 trial. *Anticancer research* 2011;31:2291-5
11
12
13 56. Hung HH, Chiou YY, Hsia CY, *et al.* Survival rates are comparable after
14
15 radiofrequency ablation or surgery in patients with small hepatocellular
16
17 carcinomas. *Clinical gastroenterology and hepatology : the official clinical*
18
19 *practice journal of the American Gastroenterological Association*
20
21 2011;9:79-86
22
23
24 57. Nishikawa H, Inuzuka T, Takeda H, *et al.* Comparison of percutaneous
25
26 radiofrequency thermal ablation and surgical resection for small hepatocellular
27
28 carcinoma. *BMC gastroenterology* 2011;11:143
29
30
31
32 58. Yun WK, Choi MS, Choi D, *et al.* Superior long-term outcomes after surgery
33
34 in child-pugh class a patients with single small hepatocellular carcinoma
35
36 compared to radiofrequency ablation. *Hepatology international* 2011;5:722-9
37
38
39 59. Zhang J, Liu HC, Zhou L. The effectiveness of radiofrequency ablation for the
40
41 treatment of liver cancer. *Journal of Hepatobiliary Surgery* 2011;19:30-33
42
43
44 60. Zhao M, Wang JP, Li W, *et al.* [Comparison of safety and efficacy for
45
46 transcatheter arterial chemoembolization alone and plus radiofrequency
47
48 ablation in the treatment of single branch portal vein tumor thrombus of
49
50 hepatocellular carcinoma and their prognosis factors]. *Zhonghua yi xue za zhi*
51
52 2011;91:1167-72
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

61. Feng K, Yan J, Li X, *et al.* A randomized controlled trial of radiofrequency ablation and surgical resection in the treatment of small hepatocellular carcinoma. *Journal of hepatology* 2012;57:794-802

62. Peng ZW, Lin XJ, Zhang YJ, *et al.* Radiofrequency ablation versus hepatic resection for the treatment of hepatocellular carcinomas 2 cm or smaller: a retrospective comparative study. *Radiology* 2012;262:1022-33

63. Peng ZW, Zhang YJ, Liang HH, *et al.* Recurrent hepatocellular carcinoma treated with sequential transcatheter arterial chemoembolization and RF ablation versus RF ablation alone: a prospective randomized trial. *Radiology* 2012;262:689-700

64. Signoriello S, Annunziata A, Lama N, *et al.* Survival after locoregional treatments for hepatocellular carcinoma: a cohort study in real-world patients. *TheScientificWorldJournal* 2012;2012:564706

65. Wang JH, Wang CC, Hung CH, *et al.* Survival comparison between surgical resection and radiofrequency ablation for patients in BCLC very early/early stage hepatocellular carcinoma. *Journal of hepatology* 2012;56:412-8

66. Desiderio J, Trastulli S, Pasquale R, *et al.* Could radiofrequency ablation replace liver resection for small hepatocellular carcinoma in patients with compensated cirrhosis? A 5-year follow-up. *Langenbeck's archives of surgery / Deutsche Gesellschaft fur Chirurgie* 2013;398:55-62

67. Ding J, Jing X, Liu J, *et al.* Comparison of two different thermal techniques for the treatment of hepatocellular carcinoma. *European journal of radiology*

- 2013;82:1379-84
68. Guo WX, Sun JX, Cheng YQ, *et al.* Percutaneous radiofrequency ablation versus partial hepatectomy for small centrally located hepatocellular carcinoma. *World journal of surgery* 2013;37:602-7
69. Hasegawa K, Kokudo N, Makuuchi M, *et al.* Comparison of resection and ablation for hepatocellular carcinoma: a cohort study based on a Japanese nationwide survey. *Journal of hepatology* 2013;58:724-9
70. Iida H, Aihara T, Ikuta S, *et al.* A comparative study of therapeutic effect between laparoscopic microwave coagulation and laparoscopic radiofrequency ablation. *Hepato-gastroenterology* 2013;60:662-5
71. Imai K, Beppu T, Chikamoto A, *et al.* Comparison between hepatic resection and radiofrequency ablation as first-line treatment for solitary small-sized hepatocellular carcinoma of 3 cm or less. *Hepatology research : the official journal of the Japan Society of Hepatology* 2013;43:853-64
72. Kim JW, Shin SS, Kim JK, *et al.* Radiofrequency ablation combined with transcatheter arterial chemoembolization for the treatment of single hepatocellular carcinoma of 2 to 5 cm in diameter: comparison with surgical resection. *Korean journal of radiology* 2013;14:626-35
73. Lai EC, Tang CN. Radiofrequency ablation versus hepatic resection for hepatocellular carcinoma within the Milan criteria--a comparative study. *International journal of surgery* 2013;11:77-80
74. Lin ZZ, Shau WY, Hsu C, *et al.* Radiofrequency ablation is superior to ethanol

injection in early-stage hepatocellular carcinoma irrespective of tumor size. *PloS one* 2013;8:e80276

75. Peng ZW, Zhang YJ, Chen MS, *et al.* Radiofrequency ablation with or without transcatheter arterial chemoembolization in the treatment of hepatocellular carcinoma: a prospective randomized trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2013;31:426-32

76. Tohme S, Geller DA, Cardinal JS, *et al.* Radiofrequency ablation compared to resection in early-stage hepatocellular carcinoma. *HPB : the official journal of the International Hepato Pancreato Biliary Association* 2013;15:210-7

77. Wong KM, Yeh ML, Chuang SC, *et al.* Survival comparison between surgical resection and percutaneous radiofrequency ablation for patients in Barcelona Clinic Liver Cancer early stage hepatocellular carcinoma. *Indian journal of gastroenterology : official journal of the Indian Society of Gastroenterology* 2013;32:253-7

78. Zhang L, Wang N, Shen Q, *et al.* Therapeutic efficacy of percutaneous radiofrequency ablation versus microwave ablation for hepatocellular carcinoma. *PloS one* 2013;8:e76119

79. Abdelaziz A, Elbaz T, Shousha HI, *et al.* Efficacy and survival analysis of percutaneous radiofrequency versus microwave ablation for hepatocellular carcinoma: an Egyptian multidisciplinary clinic experience. *Surgical endoscopy* 2014;28:3429-34

80. Shi J, Sun Q, Wang Y, *et al.* Comparison of microwave ablation and surgical

- resection for treatment of hepatocellular carcinomas conforming to Milan criteria. *Journal of gastroenterology and hepatology* 2014;29:1500-7
81. Yang HJ, Lee JH, Lee DH, *et al.* Small single-nodule hepatocellular carcinoma: comparison of transarterial chemoembolization, radiofrequency ablation, and hepatic resection by using inverse probability weighting. *Radiology* 2014;271:909-18
82. Zhang T, Li K, Luo H, *et al.* [Long-term outcomes of percutaneous microwave ablation versus repeat hepatectomy for treatment of late recurrent small hepatocellular carcinoma: a retrospective study]. *Zhonghua yi xue za zhi* 2014;94:2570-2
83. Pompili M, De Matthaeis N, Saviano A, *et al.* Single hepatocellular carcinoma smaller than 2 cm: are ethanol injection and radiofrequency ablation equally effective? *Anticancer research* 2015;35:325-32
84. Xu J, Zhao Y. Comparison of percutaneous microwave ablation and laparoscopic resection in the prognosis of liver cancer. *International journal of clinical and experimental pathology* 2015;8:11665-9
85. Lee HW, Lee JM, Yoon JH, *et al.* A prospective randomized study comparing radiofrequency ablation and hepatic resection for hepatocellular carcinoma. 2018;94:74-82
86. Li W, Zhou X, Huang Z, *et al.* Short-term and long-term outcomes of laparoscopic hepatectomy, microwave ablation, and open hepatectomy for small hepatocellular carcinoma: a 5-year experience in a single center. *Annals*

of surgical treatment and research 2017;47:650-57

87. Liu PH, Hsu CY, Hsia CY, *et al.* Surgical Resection Versus Radiofrequency Ablation for Single Hepatocellular Carcinoma \leq 2 cm in a Propensity Score Model. *Annals of surgery* 2016;263:538-45

88. Liu H, Wang ZG, Fu SY, *et al.* Randomized clinical trial of chemoembolization plus radiofrequency ablation versus partial hepatectomy for hepatocellular carcinoma within the Milan criteria. *The British journal of surgery* 2016;103:348-56

89. Hof J, Wertenbroek MW, Peeters PM, *et al.* Outcomes after resection and/or radiofrequency ablation for recurrence after treatment of colorectal liver metastases. *The British journal of surgery* 2016;103:1055-62

90. Vogl TJ, Farshid P, Naguib NN, *et al.* Ablation therapy of hepatocellular carcinoma: a comparative study between radiofrequency and microwave ablation. *Abdominal imaging* 2015;40:1829-37

91. Lee YH, Hsu CY, Chu CW, *et al.* Radiofrequency ablation is better than surgical resection in patients with hepatocellular carcinoma within the Milan criteria and preserved liver function: a retrospective study using propensity score analyses. *Journal of clinical gastroenterology* 2015;49:242-9

92. Kang TW, Kim JM, Rhim H, *et al.* Small Hepatocellular Carcinoma: Radiofrequency Ablation versus Nonanatomic Resection--Propensity Score Analyses of Long-term Outcomes. *Radiology* 2015;275:908-19

93. Zhou Z, Lei J, Li B, *et al.* Liver resection and radiofrequency ablation of very

- early hepatocellular carcinoma cases (single nodule <2 cm): a single-center study. *European journal of gastroenterology & hepatology* 2014;26:339-44
94. Ko S, Jo H, Yun S, *et al.* Comparative analysis of radiofrequency ablation and resection for resectable colorectal liver metastases. *World journal of gastroenterology* 2014;20:525-31
95. Kim JM, Kang TW, Kwon CH, *et al.* Single hepatocellular carcinoma ≤ 3 cm in left lateral segment: liver resection or radiofrequency ablation? *World journal of gastroenterology* 2014;20:4059-65
96. Agcaoglu O, Aliyev S, Karabulut K, *et al.* Complementary use of resection and radiofrequency ablation for the treatment of colorectal liver metastases: an analysis of 395 patients. *World journal of surgery* 2013;37:1333-9
97. Goldberg SN, Giran GD, Lukyanov AN, *et al.* Percutaneous tumor ablation: increased necrosis with combined radio-frequency ablation and intravenous liposomal doxorubicin in a rat breast tumor model. *Radiology* 2002;222:797-804
98. Taub R. Liver regeneration: from myth to mechanism. *Nature reviews Molecular cell biology* 2004;5:836-47
99. Martins PN, Theruvath TP, Neuhaus P. Rodent models of partial hepatectomies. *Liver international : official journal of the International Association for the Study of the Liver* 2008;28:3-11
100. Kele PG, De BM, Ej VDJ, *et al.* Early hepatic regeneration index and completeness of regeneration at 6 months after partial hepatectomy. *British*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Journal of Surgery 2012;99:1113–19

101. Preziosi ME, Monga SP. Update on the Mechanisms of Liver Regeneration.

Seminars in Liver Disease 2017;37:141

102. Morise Z, Kawabe N, Tomishige H, *et al.* Recent advances in liver resection for hepatocellular carcinoma. *Frontiers in Surgery* 2014;1:21

For peer review only

File legends:**Figure 1 Flow chart of search.****Figure 2 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in RCTs.**

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments. The number in red indicates the sample size and the number in black indicates the number of studies.

i Lesions < 3 cm.

ii Lesions 3-5 cm.

iii Lesions \leq 5 cm.

Figure 3 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in all studies.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments. The number in red indicates the sample size and the number in black indicates the number of studies.

i Lesions < 3 cm.

ii Lesions 3-5 cm.

iii Lesions \leq 5 cm.

Figure 4 Treatment ranks for 1-year, 3-year and 5-year survival rates, according

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

to lesion size in RCTs

- A Lesions < 3 cm
- B Lesions 3-5 cm
- C Lesions \leq 5 cm (full sample).

Figure 5 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in all studies.

- A Lesions < 3 cm
- B Lesions 3-5 cm
- C Lesions \leq 5 cm (full sample).

Table 1 Odds ratios (95% credible interval) according to network meta-analyses for the survival for all pairwise comparisons in randomized controlled trials.

3cm for 1-year survival					
PEI					
1.17 (0.11-4.66)	TR				
0.08 (0-0.38)	0.15 (0-0.80)	MWA			
0.67 (0.28-1.35)	1.25 (0.16-4.64)	173.30 (1.90-537.40)	RFA		
0.64(0.18-1.61)	1.08 (0.15-3.78)	152.70 (1.44-505.80)	0.97 (0.42-1.98)	RES	
3cm for 3-year survival					
PEI					
1.02 (0.14-3.56)	TR				
NA	NA	MWA			
0.79 (0.45-1.39)	1.54 (0.25-13.43)	NA	RFA		
0.58 (0.29-1.16)	1.17 (0.16-4.17)	NA	0.75 (0.41-1.31)	RES	
3cm for 5-year survival					
PEI					
3.93 (0.03-19.61)	TR				
NA	NA	MWA			
0.94 (0.08-3.97)	2.87 (0.04-13.43)	NA	RFA		
0.50 (0.04-2.04)	0.84 (0.03-4.18)	NA	0.72 (0.10-2.47)	RES	
3-5cm for 1-year survival					
PEI					
NA	TR				
NA	NA	MWA			
NA	3.40 (0.64-11.93)	NA	RFA		
NA	1.00 (0-5.00)	NA	0.25 (0-1.47)	RES	
3-5cm for 3-year survival					
PEI					
NA	TR				
NA	NA	MWA			
NA	3.98 (0.71-15.22)	NA	RFA		
NA	1.14 (0-6.20)	NA	0.24 (0-1.25)	RES	
3-5cm for 5-year survival					
PEI					
NA	TR				
NA	NA	MWA			
NA	7.64 (0.14-42.49)	NA	RFA		
NA	12.87 (0.02-44.43)	NA	1.05 (0.03-5.33)	RES	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

≤5cm for 1-year survival

PEI					
0.29 (0.09-0.73)	TR				
0.27 (0.05-0.84)	1.09 (0.16-3.50)	MWA			
0.65 (0.33-1.13)	2.69 (1.02-6.04)	3.84 (0.81-11.60)	RFA		
0.37 (0.13-0.82)	1.50 (0.48-3.67)	2.01 (0.47-5.70)	0.57 (0.27-1.08)	RES	

≤5cm for 3-year survival

PEI					
0.64 (0.19-1.67)	TR				
1.05 (0.12-4.56)	1.86 (0.21-7.59)	MWA			
0.86 (0.39-1.79)	1.56 (0.66-3.25)	1.77 (0.22-6.24)	RFA		
0.55 (0.19-1.44)	0.98 (0.35-2.41)	1.00 (0.16-3.30)	0.65 (0.31-1.29)	RES	

≤5cm for 5-year survival

PEI					
0.53 (0.06-1.90)	TR				
NA	NA	MWA			
0.74 (0.16-2.00)	2.29 (0.41-7.61)	NA	RFA		
0.41 (0.11-1.02)	1.35 (0.23-4.69)	NA	0.66 (0.20-1.62)	RES	

The reference treatment (1.00) for all comparisons is listed to the right hand side
RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection;

Table 2 Odds ratios (95% credible interval) according to network meta-analyses for the survival for all pairwise comparisons in all studies

3cm for 1-year survival					
PEI	TR	MWA	RFA	RES	
0.69 (0.14-2.13)					
0.49 (0.18-1.10)	1.08 (0.21-7.87)				
0.68 (0.38-1.09)	1.48 (0.34-4.23)	1.59 (0.69-3.17)			
0.63 (0.22-1.44)	1.30 (0.28-3.88)	1.49 (0.44-3.85)	0.94 (0.39-1.91)		
3cm for 3-year survival					
PEI	TR	MWA	RFA	RES	
0.90 (0.29-2.17)					
1.01 (0.47-1.95)	1.38 (0.42-3.40)				
0.96 (0.59-1.50)	1.31 (0.47-2.92)	1.02 (0.57-1.70)			
0.68 (0.30-1.39)	0.90 (0.31-2.10)	0.73 (0.30-1.55)	0.72 (0.37-1.30)		
3cm for 5-year survival					
PEI	TR	MWA	RFA	RES	
1.07 (0.31-2.72)					
0.86 (0.39-1.65)	1.03 (0.28-2.73)				
0.82 (0.48-1.29)	0.99 (0.32-2.39)	1.04 (0.50-1.77)			
0.43 (0.17-0.89)	0.49 (0.16-0.18)	0.55 (0.19-1.25)	0.54 (0.24-1.05)		
3-5cm for 1-year survival					
PEI	TR	MWA	RFA	RES	
0.20 (0.05-0.54)					
0.55 (0.09-1.76)	3.39 (0.58-10.44)				
0.49 (0.18-1.12)	2.99 (1.14-6.58)	1.29 (0.32-3.60)			
0.06 (0-0.31)	0.36 (0.01-2.08)	0.15 (0-1.00)	0.12 (0-0.63)		
3-5cm for 3-year survival					
PEI	TR	MWA	RFA	RES	
0.28 (0.04-0.96)					
0.61 (0.08-2.26)	2.62 (0.61-7.90)				
0.55 (0.12-1.69)	2.38 (0.93-5.38)	1.15 (0.39-2.65)			
0.06 (0-0.28)	0.26 (0.01-1.10)	0.12 (0.01-0.53)	0.11 (0.01-0.40)		
3-5cm for 5-year survival					
PEI	TR	MWA	RFA	RES	
5.77 (0.01-2.84)					
4.15 (0.04-5.18)	11.97 (0.19-46.76)				
0.86 (0.06-2.68)	6.16 (0.27-25.58)	1.26 (0.19-4.04)			
3.02 (0.01-2.40)	14.31 (0.04-21.06)	1.24 (0.02-4.46)	0.69 (0.04-3.16)		

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

≤5cm for 1-year survival

PEI					
0.34 (0.11-0.63)	TR				
0.81 (0.38-1.51)	2.69 (0.99-6.00)	MWA			
0.77 (0.51-1.10)	2.55 (1.20-4.85)	1.04 (0.55-1.76)	RFA		
0.52 (0.24-0.96)	1.72 (0.66-3.70)	0.70 (0.29-1.39)	0.68 (0.35-1.17)	RES	

≤5cm for 3-year survival

PEI					
0.64 (0.32-1.16)	TR				
0.98 (0.55-1.65)	1.65 (0.80-3.03)	MWA			
0.94 (0.64-1.34)	1.57 (0.89-2.57)	0.99 (0.64-1.47)	RFA		
0.59 (0.30-1.04)	0.97 (0.48-1.79)	0.62 (0.32-1.09)	0.63 (0.37-1.01)	RES	

≤5cm for 5-year survival

PEI					
0.84 (0.35-1.74)	TR				
0.87(0.46-1.51)	1.16 (0.46-2.46)	MWA			
0.87 (0.57-1.26)	1.16 (0.54-2.21)	1.06 (0.64-1.61)	RFA		
0.45 (0.23-0.82)	0.59 (0.25-1.20)	0.55 (0.25-1.05)	0.52 (0.29-0.88)	RES	

The reference treatment (1.00) for all comparisons is listed to the right hand side

RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection.




Figure 1 Flow chart of search.

254x190mm (300 x 300 DPI)




Figure 2 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in RCTs. Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments. The number in red indicates the sample size and the number in black indicates the number of studies.

- i Lesions < 3 cm.
- ii Lesions 3-5 cm.
- iii Lesions ≤ 5 cm.

500x500mm (300 x 300 DPI)




Figure 3 Networks of treatment comparisons for 1-year (A), 3-year (B), and 5-year (C) survival rates in all studies.

Circle size is proportional to the number of included patients and line width indicates the number of studies comparing the connected treatments. The number in red indicates the sample size and the number in black indicates the number of studies.

i Lesions < 3 cm.

ii Lesions 3-5 cm.

iii Lesions ≤ 5 cm.

227x227mm (300 x 300 DPI)




Figure 4 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in RCTs
A Lesions < 3 cm
B Lesions 3-5 cm
C Lesions ≤ 5 cm (full sample).

193x165mm (300 x 300 DPI)




Figure 5 Treatment ranks for 1-year, 3-year and 5-year survival rates, according to lesion size in all studies.
A Lesions < 3 cm
B Lesions 3-5 cm
C Lesions ≤ 5 cm (full sample).

193x165mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Text S1.
PRISMA NMA Checklist of Items to Include When Reporting A Systematic Review Involving a Network Meta-analysis

Section/Topic	Item #	Checklist Item	Reported on Page #
TITLE			
Title	1	Identify the report as a systematic review <i>incorporating a network meta-analysis (or related form of meta-analysis).</i>	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: Background: main objectives Methods: data sources; study eligibility criteria, participants, and interventions; study appraisal; and <i>synthesis methods, such as network meta-analysis.</i> Results: number of studies and participants identified; summary estimates with corresponding confidence/credible intervals; <i>treatment rankings may also be discussed. Authors may choose to summarize pairwise comparisons against a chosen treatment included in their analyses for brevity.</i> Discussion/Conclusions: limitations; conclusions and implications of findings. Other: primary source of funding; systematic review registration number with registry name.	5,6
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known, <i>including mention of why a network meta-analysis has been conducted.</i>	7,8
Objectives	4	Provide an explicit statement of questions being addressed, with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	8
METHODS			
Protocol and registration	5	Indicate whether a review protocol exists and if and where it can be accessed (e.g., Web address); and, if available, provide registration information, including registration number.	8,9

Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale. <i>Clearly describe eligible treatments included in the treatment network, and note whether any have been clustered or merged into the same node (with justification).</i>	9,10
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	9
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	9,10, Figure 1, Additional file 1: Text S2
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	9,10
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	10
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	11
Geometry of the network	S1	Describe methods used to explore the geometry of the treatment network under study and potential biases related to it. This should include how the evidence base has been graphically summarized for presentation, and what characteristics were compiled and used to describe the evidence base to readers.	11
Risk of bias within individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	11,12
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means). <i>Also describe the use of additional summary measures assessed, such as treatment rankings and surface under the cumulative ranking curve (SUCRA) values, as well as modified approaches used to present summary findings from meta-analyses.</i>	11,12
Planned methods of analysis	14	Describe the methods of handling data and combining results of studies for each network meta-analysis. This should include, but not be limited to: <ul style="list-style-type: none"> • <i>Handling of multi-arm trials;</i> • <i>Selection of variance structure;</i> • <i>Selection of prior distributions in Bayesian analyses; and</i> 	11,12

		<ul style="list-style-type: none">Assessment of model fit.	
Assessment of Inconsistency	S2	Describe the statistical methods used to evaluate the agreement of direct and indirect evidence in the treatment network(s) studied. Describe efforts taken to address its presence when found.	10,11,12
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	10,11,12
Additional analyses	16	Describe methods of additional analyses if done, indicating which were pre-specified. This may include, but not be limited to, the following: <ul style="list-style-type: none">Sensitivity or subgroup analyses;Meta-regression analyses;Alternative formulations of the treatment network; andUse of alternative prior distributions for Bayesian analyses (if applicable).	11,12
RESULTS†			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	12
Presentation of network structure	S3	Provide a network graph of the included studies to enable visualization of the geometry of the treatment network.	12,13,Figure2-3
Summary of network geometry	S4	Provide a brief overview of characteristics of the treatment network. This may include commentary on the abundance of trials and randomized patients for the different interventions and pairwise comparisons in the network, gaps of evidence in the treatment network, and potential biases reflected by the network structure.	12,13,
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	12, Table1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment.	

Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: 1) simple summary data for each intervention group, and 2) effect estimates and confidence intervals. <i>Modified approaches may be needed to deal with information from larger networks.</i>	12,13, Figure2-5
Synthesis of results	21	Present results of each meta-analysis done, including confidence/credible intervals. <i>In larger networks, authors may focus on comparisons versus a particular comparator (e.g. placebo or standard care), with full findings presented in an appendix. League tables and forest plots may be considered to summarize pairwise comparisons.</i> If additional summary measures were explored (such as treatment rankings), these should also be presented.	12,13, Figure4-5, Additional file 1: Table S1-S13
Exploration for inconsistency	S5	Describe results from investigations of inconsistency. This may include such information as measures of model fit to compare consistency and inconsistency models, <i>P</i> values from statistical tests, or summary of inconsistency estimates from different parts of the treatment network.	12,13
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies for the evidence base being studied.	12,13, Additional file 1: Figure S1-S2
Results of additional analyses	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression analyses, <i>alternative network geometries studied, alternative choice of prior distributions for Bayesian analyses, and so forth</i>).	12,13
DISCUSSION			
Summary of evidence	24	Summarize the main findings, including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy-makers).	14-16
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review level (e.g., incomplete retrieval of identified research, reporting bias). <i>Comment on the validity of the assumptions, such as transitivity and consistency. Comment on any concerns regarding network geometry (e.g., avoidance of certain comparisons).</i>	16
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	17
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. This should also include information regarding whether funding has been received from manufacturers of treatments in the network and/or whether some of the	17

authors are content experts with professional conflicts of interest that could affect use of treatments in the network.

PICOS = population, intervention, comparators, outcomes, study design.
* Text in *italics* indicateS wording specific to reporting of network meta-analyses that has been added to guidance from the PRISMA statement.
† Authors may wish to plan for use of appendices to present all relevant information in full detail for items in this section.

Text S2.

Search strategy:

Pubmed (1950-present)

1. ("TACE" OR "transarterial chemoembolization")
2. ("RFA" OR "radiofrequency ablation" OR "RF ablation" OR "radiofrequency thermal ablation" OR "RTA")
3. (PEI OR "ethanol injection" OR "ethanol ablation" OR "alcohol ablation")
4. ("microwave ablation" OR "microwave thermal ablation" OR MWA)
5. (liver OR hepato*)
6. (neoplas* OR cancer OR tumor OR tumour OR carcinoma OR oncolog*)
7. 1 OR 2 OR 3 OR 4
8. 5 AND 6 AND 7
9. "Ablation Techniques"[Mesh]
10. "Embolization"[Mesh]
11. "Liver Neoplasms"[Mesh]
12. 9 OR 10
13. 12 AND 11
14. 8 OR 13
15. (resection OR surgery OR hepatectomy)
16. (ablation OR injection OR embolization)
17. 5 AND 6 AND 15 AND 16
18. "Hepatectomy"[Mesh]
19. 12 AND 18 AND 11
20. 17 OR 19

21. 14 OR 20

Embase(1980-present)

1. 'TACE':ab,ti
2. 'transarterial chemoembolization':ab,ti
3. 1 OR 2
4. 'rfa':ab,ti
5. 'radiofrequency ablation':ab,ti
6. 'rf ablation':ab,ti
7. 'radiofrequency thermal ablation':ab,ti
8. 'rta':ab,ti
9. 4 OR 5 OR 6 OR 7 OR 8
10. 'PEI':ab,ti
11. 'ethanol injection ':ab,ti
12. 'ethanol ablation ':ab,ti
13. 'alcohol ablation ':ab,ti
14. 10 OR 11 OR 12 OR 13
15. 'microwave ablation ':ab,ti
16. 'microwave thermal ablation ':ab,ti
17. 'MWA ':ab,ti
18. 15 OR 16 OR 17
19. 'liver':ab,ti
20. 'hepato*':ab,ti
21. 19 OR 20
22. 'neoplas*':ab,ti
23. 'cancer ':ab,ti
24. 'tumor ':ab,ti
25. 'tumour ':ab,ti
26. 'carcinoma ':ab,ti
27. 'oncolog*':ab,ti

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
28. 22 OR 23 OR 24 OR 25 OR 26 OR 27
29. 3 OR 9 OR 14 OR 18
30. 21 AND 28 AND 29
31. ' resection':ab,ti
32. ' surgery':ab,ti
33. ' hepatectomy':ab,ti
34. 31 OR 32 OR 33
35. ' ablation':ab,ti
36. ' injection':ab,ti
37. ' embolization':ab,ti
38. 35 OR 36 OR 37
39. 34 AND 38 AND 21 AND 28
40. 30 OR 39

Scoups

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
1. TITLE-ABS-KEY (“TACE”)
2. TITLE-ABS-KEY (“transarterial chemoembolization”)
3. 1 OR 2
4. TITLE-ABS-KEY ("RFA")
5. TITLE-ABS-KEY (“radiofrequency ablation”)
6. TITLE-ABS-KEY ("RF ablation")
7. TITLE-ABS-KEY ("radiofrequency thermal ablation")
8. TITLE-ABS-KEY ("RTA")
9. 4 OR 5 OR 6 OR 7 OR8
10. TITLE-ABS-KEY (“PEI”)
11. TITLE-ABS-KEY (“ethanol injection”)
12. TITLE-ABS-KEY (“ethanol ablation”)
13. TITLE-ABS-KEY (“alcohol ablation”)
14. 10 OR 11 OR 12 OR 13
15. TITLE-ABS-KEY ("microwave ablation")

16. TITLE-ABS-KEY ("microwave thermal ablation ")
17. TITLE-ABS-KEY ("MWA")
18. 15 OR 16 OR 17
19. TITLE-ABS-KEY ("liver ")
20. TITLE-ABS-KEY ("hepato*")
21. 19 OR 20
22. TITLE-ABS-KEY ("neoplas*")
23. TITLE-ABS-KEY ("cancer")
24. TITLE-ABS-KEY ("tumor")
25. TITLE-ABS-KEY ("tumour")
26. TITLE-ABS-KEY ("carcinoma")
27. TITLE-ABS-KEY ("oncolog*")
28. 22 OR 23 OR 24 OR 25 OR 26 OR 27
29. 3 OR 9 OR 14 OR 18
30. 29 AND 21 AND 28
31. TITLE-ABS-KEY ("resection")
32. TITLE-ABS-KEY ("surgery")
33. TITLE-ABS-KEY ("hepatectomy")
34. 31 OR 32 OR 33
35. TITLE-ABS-KEY ("ablation")
36. TITLE-ABS-KEY ("injection")
37. TITLE-ABS-KEY ("embolization")
38. 35 OR 36 OR 37
39. 34 AND 38 AND 21 AND 28
40. 30 OR 39

Web of science

1. TS=(ablation)
2. TS=(embolization)
3. 1 OR 2

- 4. TS=(hepatectomy)
- 5. TS=(liver neoplasms)
- 6. 3 AND 4 AND 5
- 7. TI=(resection)
- 8. TI=(surgery)
- 9. TI=(hepatectomy)
- 10. 7 OR 8 OR 9
- 11. TI=(ablation)
- 12. TI=(injection)
- 13. TI=(embolization)
- 14. 11 OR 12 OR 13
- 15. TI=(liver)
- 16. TI=(hepato*)
- 17. 15 OR 16
- 18. TI=(neoplas*)
- 19. TI=(cancer)
- 20. TI=(tumor)
- 21. TI=(tumour)
- 22. TI=(carcinoma)
- 23. TI=(oncolog*)
- 24. 18 OR 19 OR 20 OR 21 OR 22 OR 23
- 25. 10 AND 14 AND 17 AND 24
- 26. 3 AND 5
- 27. TI=(TACE)
- 28. TI=("transarterial chemoembolization")
- 29. 27 OR 28
- 30. TI=(RFA)
- 31. TI=("radiofrequency ablation")
- 32. TI=("RF ablation")
- 33. TI=("radiofrequency thermal ablation")
- 34. TI=(RTA)

35. 30 OR 31 OR 32 OR 33 OR 34
36. TI=(PEI)
37. TI=("ethanol injection")
38. TI=("ethanol ablation")
39. TI=("alcohol ablation")
40. 36 OR 37 OR 38 OR 39
41. TI=("microwave ablation")
42. TI=("microwave thermal ablation")
43. TI=(MWA)
44. 41 OR 42 OR 43
45. 29 OR 35 OR 40 OR 44
46. 46 AND 17 AND 24
47. 6 OR 25 OR 26 OR 46

Table S1.
Summary of the studies included in the network meta-analysis.

Study Year	Design style	Countr ,	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Zhang 2002 ²³	Prospectiv e cohort	China	HCC	0.3-2	RFA	15(15)	13/2	61.8 (38-78)	4.1 (2.4-6.0)	NA	0.80(1y)	0.80(1y)	NA
					TR	15(15)	12/3	57.8 (39-72)	4.6 (2.3-7.1)	NA	1.00(1y)	1.00 (1y)	NA
Lencioni 2003 ²⁴	RCT	Italy	HCC	1.9±0.8	RFA	52(69)	36/16	67±6 (52-78)	2.8±0.6	1.00(1y)	NA	1.00(1y)	15 pain and 10 fever
					PEI	50(73)	30/20	69±7.4 (40-82)	2.8±0.8	0.96(1y)	NA	0.96(1y)	13 pain and 5 fever
Lin 2004 ²⁵	RCT	China	HCC	2±0.9	RFA	52(69)	35/17	62±11	2.9±0.8	0.76(3y)	NA	0.35(3y)	1 transient pleural effusion

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					PEI	52(67)	34/18	59±10	2.8±0.8	0.66(3y)	NA	0.17(3y)	1 pain
Vivarelli 2004 ²⁶	Retrospect ive cohort	Italy	HCC	2.4	RES	79(92)	57/22	65.2±8.2 (43-81)	≤3/3.1-5 (21/58)	0.81(3y)	0.59(3y)	0.65(3y)	NA
					RFA	79(112)	67/12	67.8±8.7 (41-88)	≤3/3.1-5 (22/57)	0.50(3y)	0.25(3y)	0.33(3y)	NA
Cho 2005 ²⁷	Retrospect ive cohort	Korea	HCC	0.1-3	RES	61	48/13	57	3.4±1.0	NA	0.77(3y)	0.77(3y)	2 bleeding, 1 intraabdominal abscess, 1 wound infection
					RFA	99	76/23	58	3.1±0.8	NA	0.80(3y)	0.80(3y)	1 chest wall metastasis, 1 cholecystitis, 1 iatrogenic burn, 1 ileus, 1 hepatic infarction
Huang 2005 ²⁹	RCT	China	HCC	1-4.9	RES	38(42)	27/11	59±11.4	≤2/2.1-3 (24/14)	0.82	NA	0.82	NA
					PEI	38(46)	19/19	63±10.9	≤2/2.1-3 (21/17)	0.45	NA	0.45	NA
Hong 2005 ²⁸	Retrospect ive cohort	Korea	HCC	2.9(0.4-4.6)	RES	93	69/24	49.2±9.9	2.5±0.8	0.84(3y)	NA	0.84(3y)	NA
					RFA	55	41/14	59.1±9.6	2.4±0.6	0.73(3y)	NA	0.73(3y)	NA
Lin 2005 ³⁰	RCT	China	HCC	2.3±1	RFA	62(78)	40/22	61±10	2.5±1	0.74(3y)	NA	0.74(3y)	2 haemothorax, 1 gastric bleeding and perforation
					PEI	62(76)	39/23	60±8	2.3±0.8	0.60(3y)	NA	0.60(3y)	1 pain
Lu 2005 ³¹	Retrospect ive cohort	China	HCC	2.1±1.1	RFA	53(72)	43/10	54.5±11.7 (24-74)	2.6±1.2 (1.0-6.1)	0.38(3y)	NA	0.38(3y)	2 skin burn, 1 puncture wound infection
					MWA	49(98)	44/5	50.1±13.7 (24-74)	2.5±1.2 (0.9-7.2)	0.51(3y)	NA	0.51(3y)	2 puncture wounds, 2 subcapsular hematoma

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Montorsi 2005 ³²	Prospective cohort	Italy	HCC	2.1	RES	40	33/7	67±9	<5cm	NA	NA	0.73(3y)	NA
					RFA	58	43/15	67±6		NA	NA	0.60(3y)	NA
Shiina 2005 ³³	RCT	Japan	HCC	3.1(0.6-4.3)	RFA	118(184)	79/39	≤65/>65 (44/74)	≤2/>2 (45/73)	NA	NA	0.61(3y)	1 transient jaundice, 1 skin burn, 1 hepatic infarction, 3 neoplastic seeding
					PEI	114(188)	87/27	≤65/>65 (41/73)	≤2/>2 (57/57)	NA	NA	0.45(3y)	1 abscess2 neoplastic seeding
Chen 2006 ³⁴	RCT	China	HCC	2.4±1	RES	90	75/15	49.4±10.9	≤3/3.1-5 (42/48)	0.53	NA	0.53	2 liver failure, 2 gastrointestinal bleeding, 27 ascites
					RFA	71	56/15	51.9±11.2	≤3/3.1-5 (37/34)	0.58	NA	0.58	3 skin burn
Lu 2006 ³⁵	RCT	China	Early HCC	1.8	RES	54(56)	37/17	49±14	3.2±1.0	NA	NA	0.86 (3y)	3 wound infection, 1 gastrointestinal bleeding
					RFA	51(57)	42/9	55±13	2.7±1.0	NA	NA	0.87 (3y)	1 peritoneal bleeding, 1 neoplastic seeding
Cho 2007 ³⁶	Retrospective cohort	Korea	HCC	5.7	RES	130(145)	103/27	56.3±8.8	≤2/2.1-3 (43/87)	0.66	NA	0.66	NA
					PEI	249(275)	181/68	57.7±9.7	≤2/2.1-3 (169/80)	0.49	NA	0.49	NA
Gao 2007 ³⁷	Retrospective cohort	China	HCC	4.6	RES	34(37)	28/6	51.5 (38-67)	2.58±0.41	0.76	NA	0.76	12 fever, 5 ascites
					RFA	53(84)	41/12	57.1 (31-81)	2.45±0.37	0.62	NA	0.62	2 bleeding, 1 fistula, 1 wound infection, 6 fever, 9 ascites
Lupo 2007 ³⁸	Retrospective cohort	Italy	HCC	2.6	RES	42	33/9	67(28-80)	4.0(3-5)	NA	0.43	0.43	2 urine infection, 1 bilioma, 1 pleural effusion, 1 renal failure, 1 intra-abdominal bleeding

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	60	47/13	68(42-85)	3.65(3-5)	NA	0.32	0.32	2 liver failure, 1 hepatic abscess, 2 pleural effusion, 1 cutaneous metastasis
Zhou 2007 ³⁹	Retrospect ive cohort	China	HCC	0.5-5.9	RES	40(42)	35/5	53±13	≤2/2.1-5 (7/33)	NA	NA	0.75	NA
					RFA	47(54)	37/10	57±14	≤2/2.1-5 (8/39)	NA	NA	0.19	NA
Abu-Hilal 2008 ⁴⁰	Retrospect ive cohort	Italy and China	Early HCC	3.6	RES	34	26/8	67	3.8(1.3-5)	NA	0.56	0.56	3 hepatic failure
					RFA	34	27/7	65	3(2-5)	NA	0.56	0.56	1 artero-portal fistula
Brunello 2008 ⁴¹	RCT	Italy	Early HCC	2.2	RFA	70(89)	49/20	70.3±8.1	1.27±0.54	0.60(3y)	NA	0.60(3y)	1 haemoperitoneum 1 right haemothorax
					PEI	69(88)	43/27	69.0±7.7	1.27±0.57	0.58(3y)	NA	0.58(3y)	1 haemoperitoneum 1 death
Guglielmi 2008 ⁴²	Retrospect ive cohort	Italy	HCC	2.3	RES	91(113)	73/18	≤65/>65 (47/44)	≤3/3.1-6 (31/60)	0.55	0.43	0.48	33 postoperative complications
					RFA	109(153)	88/21	≤65/>65 (38/71)	≤3/3.1-6 (32/77)	0.28	0.14	0.20	11 postoperative complications
Hiraoka 2008 ⁴³	Retrospect ive cohort	Japan	HCC	2.5	RES	59	44/15	62.4±10.6	2.27±0.55	0.59	NA	0.59	1 death, 2 abscess
					RFA	105	76/29	69.4±9.1	1.98±0.52	0.59	NA	0.59	1 biloma, 2 dermatitis
Bu 2009 ⁴⁹	Retrospect ive cohort	China	HCC	2.9(0.5-6)	RES	42(46)	36/6	53.93±10.74	≤3/3.1-5 (14/28)	0.57	0.46	0.50	1 postoperative hemorrhage, 3 pleural effusions, 2 subdiaphragmatic effusion

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	46(54)	40/6	55.89±7.37	≤3/3.1-5 (20/26)	0.50	0.31	0.37	4 pleural effusions, 1 postoperative hemorrhage, 1 skin burn
Ohmoto 2009 ⁴⁴	Retrospect ive cohort	Japan	HCC	2.8±2	RFA	34(37)	25/9	67 (44-78)	1.6 (0.7-2.0)	0.71	NA	0.71	2 pain, 4 fever, 1 bile duct injury, 1 pleural effusion, 1 skin burns, 1 vagovagal reflex
					MWA	49(56)	41/8	64 (38-75)	1.7 (0.8-2.0)	0.37	NA	0.37	11 pain, 17 fever, 9 bile duct injury, 8 pleural effusion, 5 ascites, 4 skin burns, 2 vagovagal reflex, 2 abscess, 2 intraperitoneal bleeding, 1 hepatic infarction, 1 portal thrombus, 1 biliary peritonitis
Sakaguchi 2009 ⁴⁵	Retrospect ive cohort	Japan	HCC	0.1-5	Laparosco pic /thorasc opic RFA	249	169/80	65.6±8.9	2.48±0.89	0.57	NA	0.57	1 frequent premature ventricular contractions, 1 liver decompensation
					Laparosco pic /thorasc opic MWA	142	107/35	64.9±7.8	2.28±0.74	0.63	NA	0.63	1 breath holding and incomplete intestinal obstruction, 2 liver decompensation
Santambrogio 2009 ⁴⁶	Prospectiv e cohort	Italy	HCC	3.2	RES	78	55/23	68±8	2.87±1.21	0.54	NA	0.54	15 extra-hepatic complications

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					Laparosco pic RFA	74	59/15	68±7	2.63±1.07	0.41	NA	0.41	14 extra-hepatic complications
Shibata 2009 ⁴⁷	RCT	Japan	HCC	2.5±1.2	RFA	43(44)	33/10	69.8±8 (44-87)	1.6±0.5 (0.8-2.6)	0.84(3y)	NA	0.84(3y)	1 pseudoaneurysm
					TR	46(49)	31/15	67.2±8.9 (45-83)	1.7±0.6 (0.9-3.0)	0.85(3y)	NA	0.85(3y)	1 hepatic infarction
Ueno 2009 ⁴⁸	Retrospect ive cohort	Japan	HCC	3(0.3-7.9)	RES	123(136)	82/41	67(28-85)	2.7±0.1	0.81	0.72	0.80	NA
					RFA	155(209)	100/55	66(40-79)	2.0±0.1	0.38	0.78	0.63	NA
Guo 2010 ⁵⁰	Retrospect ive cohort	China	HCC	2.5	RES	73(155)	57/16	50.0 (17.0-68.0)	≤3/3.1-5 (30/43)	0.27	0.47	0.44	1 postoperative hemorrhage, 5 abscess, 3 infected ascites, 1 liver failure, 4 pleural effusion
					RFA	86(211)	63/23	52.5 (26.0-80.0)	≤3/3.1-5 (42/44)	0.33	0.16	0.21	1 postoperative hemorrhage, 1 bile leak, 1 abscess, 1 infected ascites, 3 pleural effusion
Huang 2010 ⁵¹	RCT	China	HCC	3.87	RES	115(144)	85/30	55.91±12.68	≤3/3.1-5 (45/44)	0.82	0.73	0.76	1 hepatic failure, 13 ascites, 5 effusion, 9 bile leakage, 2 postoperative bleeding, 2 gastrointestinal bleeding
					RFA	115(147)	79/36	56.57±14.30	≤3/3.1-5 (57/27)	0.61	0.52	0.55	1 gastric perforation, 2 hemorrhage, 1 malignant seeding, 1 hepatic infarction
Kagawa 2010 ⁵²	Retrospect ive cohort	Japan	Early HCC	4.2	RES	55(69)	40/15	66.1±8.4	≤2/2.1-5 (9/46)	0.42	NA	0.42	2 deaths, 1 liver failure, 1 pleural effusion, 1 pneumonia, 2 biliary leakage

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					TR	62(79)	39/23	67.5±8.4	≤2/2.1-5 (19/43)	0.29	NA	0.29	1 duodenal perforation, 1 hemothorax
Morimoto 2010 ⁵³	RCT	Japan	HCC	2.7	RFA	18(25)	12/6	73 (48-84)	3.7±0.6	NA	0.78(3y)	0.78(3y)	5 pain, 2 pleural effusion
					TR	19(21)	15/4	70 (57-78)	3.6±0.7	NA	0.95(3y)	0.95(3y)	1 pain, 1 pleural effusion
Azab 2011 ⁵⁴	RCT	Egypt	HCC	1.5	RFA	30(33)	75/15	46-77	<5cm	NA	NA	0.90	5 superficial burn, 17 transient pain, 3 portal vein thrombosis, 7 fever, 1 ascites
					PEI	30(32)				NA	NA	0.83	2 portal vein thrombosis, 3 fever, 3 ascites
Giorgio 2011 ⁵⁵	RCT	Italy	HCC	1.8	RFA	142	105/37	70±2 (68-74)	2.34±0.45 (1.1-3)	0.70	NA	0.70	1 major complication
					PEI	143	102/41	72±6 (68-79)	2.27±0.48 (1.3-2.9)	0.68	NA	0.68	3 major complication
Hung 2011 ⁵⁶	Retrospect ive cohort	China	Early HCC	3.5±2	RES	229	184/45	60.07±12.56	2.88±1.06	0.77	NA	0.77	NA
					RFA	190	121/69	67.42±11.45	2.37±0.92	0.67	NA	0.67	NA
Nishikawa 2011 ⁵⁷	Retrospect ive cohort	Japan	HCC	3.3	RES	69	50/19	67.4±9.7	2.68±0.49	0.74	NA	0.74	2 bile leakage, 2 ascites, 1 acute respiratory distress syndrome, 1 gastrointestinal bleeding
					RFA	162	95/67	68.4±8.7	1.99±0.62	0.63	NA	0.63	1 biloma, 1 ascites, 1 intra-abdominal bleeding
Yun 2011 ⁵⁸	Retrospect	Korea	HCC	3.5(0.1-9.	RES	215	171/44	51.7±9.7	2.1±0.5	0.94	NA	0.94	NA

Study Year	Design style	Countr	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
	ive cohort			1)	RFA	255	197/58	57.0±9.9	2.1±0.5	0.87	NA	0.87	NA
Zhang 2011 ⁵⁹	Retrospect ive cohort	China	HCC	0.5-3.5	RES	103(117)	78/25	56.4±15.2	<5cm	NA	NA	0.35(3y)	12 wound infection, 5 postoperative hemorrhage, 2 hepatic failure, 15 pleural effusions, 6 pleural effusions
					RFA	85(106)	62/23	58.5±12.9	<5cm	NA	NA	0.39(3y)	2 gallbladder cardiac reflex, 4 postoperative hemorrhage, 3 pleural effusions
Feng 2012 ⁶¹	RCT	China	HCC	3	RES	84(116)	75/9	47 (18-76)	2.6±0.8	0.62(3y)	NA	0.62(3y)	7 pleural effusion, 3 pneumonia, 1 effusion plus infection, 3 wound infection or dehiscence, 1 biliary fistula, 2 abdominal bleeding, 1 pneumothorax or hemothorax
					RFA	84(120)	79/5	51 (24-83)	2.4±0.6	0.55(3y)	NA	0.55(3y)	5 pleural effusion, 1 liver abscess, 2 abdominal bleeding
Peng 2012 ⁶²	Retrospect ive cohort	China	Reccurre nt HCC	4.9	RES	74	65/9	51.5±12.1 (24-75)	1.1±0.5 (0.8-2.0)	0.62	NA	0.62	1 liver failure, 2 gastrointestinal bleeding, 1 peritoneal bleeding, 1 intestinal obstruction, 1 spontaneous bacterial peritonitis, 1 persistent jaundice, 31 ascites

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	71	63/8	53.1 ±12.1 (28-74)	1.2 ±0.6 (0.9-2.0)	0.72	NA	0.72	1 gastrointestinal bleeding, 1 persistent jaundice, 12 ascites
Peng 2012 ⁶³	RCT	China	Recurrent HCC	3.3 ±1.8	RFA	70(76)	55/15	55.1 ±9.5 (22-75)	≤3/3.1-5 (46/24)	NA	0.17	0.36	1 persistent jaundice, 1 ascites, 22 fever, 45 pain, 4 vomiting
					TR	69(74)	59/9	57.5 ±10.0 (19-75)	≤3/3.1-5 (41/28)	NA	0.39	0.46	1 liver failure, 1 ascites, 27 fever, 50 pain, 42 vomiting
Signoriello 2012 ⁶⁴	Retrospective cohort	Italy	HCC	0.1-9	RES	34(44)	30/4	62 ±7	≤3/3.1-5/>5.1 (13/9/4)	NA	NA	0.29	NA
					RFA	50(74)	40/10	68 ±7	≤3/3.1-5/>5.1 (24/11/7)	NA	NA	0.15	NA
					PEI	256(349)	188/68	67 ±8	≤3/3.1-5/>5.1 (143/43/12)	NA	NA	0.20	NA
a. Wang 2012 ⁶⁵	Retrospective cohort	China	Early HCC	2.5	RES	52	38/14	≤60 (35)	NA	NA	NA	0.92	NA
					RFA	91	60/31	≤60 (40)		NA	NA	0.73	NA
b. Wang 2012 ⁶⁵	Retrospective cohort	China	Early HCC	2.5	RES	208	168/40	≤60 (113)	≤2/2.1-5 (6/202)	NA	NA	0.77	NA
					RFA	254	161/93	≤60 (85)	≤2/2.1-5 (60/194)	NA	NA	0.57	NA
Desiderio 2013 ⁶⁶	Retrospective cohort	Italy	HCC	4.3(2.3-5)	RES	52(94)	37/15	65.6 ±4.8	≤3	0.46	NA	0.46	2 hepatic failure, 1 biliary fistula, 2 hemoperitoneum, 9 ascites
					RFA	44(81)	35/9	64.4 ±6.5		0.36	NA	0.36	6 pain, 7 fever

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Ding 2013 ⁶⁷	Retrospect ive cohort	China	HCC	2.3±1.3	RFA	85(98)	68/17	58.64±8.52 (40-77)	2.38±0.81 (1.0-4.8)	0.82(3y)	NA	0.82(3y)	1 frequent premature ventricular contractions, 1 liver decompensation
					MWA	113(131)	85/28	59.06±11.68 (30-86)	2.55±0.89 (0.8-5.0)	0.78(3y)	NA	0.78(3y)	1 breath holding and incomplete intestinal obstruction, 2 liver decompensation
Guo 2013 ⁶⁸	Retrospect ive cohort	China	HCC	2.7	RES	102(129)	94/8	51.5(18-75)	≤3/3.1-5 (75/27)	NA	NA	0.63	5 postoperative hemorrhage, 3 bile leak, 4 abscess, 3 infected ascites, 1 liver failure, 4 pleural effusion
					RFA	94(125)	78/16	56(19-75)	≤3/3.1-5 (62/32)	NA	NA	0.50	1 postoperative hemorrhage, 2 bile leak, 1 abscess, 1 infected ascites, 3 pleural effusion
Hasegawa 2013 ⁶⁹	Retrospect ive cohort	Japan	HCC	2.2	RES	5361(646 1)	3967/139 4	66 (48-77)	2.3 (1.2-3)	0.71	NA	0.71	NA
					RFA	5548(741 2)	3569/197 9	69 (52-80)	2 (1-3)	0.61	NA	0.61	NA
					PEI	2059(283 6)	1303/756	69 (52-80)	1.7 (1-3)	0.56	NA	0.56	NA
Iida 2013 ⁷⁰	Retrospect ive cohort	Japan	HCC	0.1-7.5	Laparosco pic RFA	18(27)	NA	73.5±4.0	2.1±0.5	0.78	NA	0.78	1 abscess

Study Year	Design style	Country	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					Laparoscopic MWA	40(56)		70.1±6.6	2.0±0.9	0.78	NA	0.78	1 abscess
Imai 2013 ⁷¹	Retrospective cohort	Japan	HCC	4.1	RES	101	75/26	63.3±9.7	2.14±0.55	0.87	NA	0.87	NA
					RFA	82	46/36	67.6±8.5	1.87±0.50	0.60	NA	0.60	NA
Kim 2013 ⁷²	Retrospective cohort	Korea	Early HCC	0.1-4.2	RES	47	36/11	58.8±10.7	3.66±0.76	NA	0.85(3y)	0.85(3y)	2 pleural effusion, 2 pneumonia, 1 hepatic failure, 1 hepatic abscess, 1 mechanical ileus
					TR	37	31/6	61.7±11.1	3.46±0.75	NA	0.78(3y)	0.78(3y)	1 bile duct dilatation
Lai 2013 ⁷³	Retrospective cohort	China	HCC	2.9±1.5	RES	80	55/25	60.8±9.9	2.9±1.1	0.71	NA	0.71	NA
					RFA	31	19/12	63.1±12.8	1.8±0.6	0.84	NA	0.84	NA
Lin 2013 ⁷⁴	Retrospective cohort	China	Early HCC	3.4	RFA	658	393/265	64.7±10.5	2.4±1.1 (0.8-9.5)	0.60	0.50	0.55	NA
					PEI	378	243/135	63.5±12.1	2.0±0.9 (0.4-7.0)	0.50	0.28	0.40	NA
Peng 2013 ⁷⁵	RCT	China	HCC	0.6-5.2	RFA	95(133)	71/24	55.3±13.3	3.39±1.35	NA	0.59(3y)	0.59(3y)	51 pain, 26 fever, 29 vomiting, 4 ascites, 2 pleural effusion, 1 skin burn, 1 abdominal infection, 1 small intestinal obstruction
					TR	94(137)	75/19	53.3±11	3.47±1.44	NA	0.67(3y)	0.67(3y)	57 pain, 33 fever, 40 vomiting, 5 ascites, 3 pleural effusion, 1 skin burn, 1 bile duct stenosis, 1 gastric hemorrhage

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Tohme 2013 ⁷⁶	Retrospect ive cohort	Ameri ca	Early HCC	2.4	RES	50(62)	31/19	66.3±1	3.07±1.17	0.48	NA	0.48	3 pleural effusion, 1 pneumonia, 1 myocardial infarction, 2 biloma, 2 ileus, 1 ascites, 1 hyperbilirubinaemia >6, 1 renal insufficiency, 2 encephalopathy
					RFA	60(75)	38/22	65.6±12	2.36±0.94	0.35	NA	0.35	1 oesophagitis, 3 encephalopathy, 1 cholangitis, 2 ascites, 1 renal insufficiency, 1 pneumonia
Wong 2013 ⁷⁷	Retrospect ive cohort	China	Early HCC	0.1-5	RES	46	30/16	55.1±12	2.1±0.6	0.85	NA	0.85	2 fever, 1 increased serum alanine aminotransferase level, 2 atelectasis, 2 biloma
					RFA	36	18/18	63.5±13	1.9±0.6	0.72	NA	0.72	None
Zhang 2013 ⁷⁸	Retrospect ive cohort	China	HCC	2.2±1	RFA	78(97)	64/14	54±10.5 (30-80)	≤3/3.1-5 (47/31)	0.43	0.39	0.41	1 persistent jaundice, 1 biliary fistula
					MWA	77(105)	67/10	54±9.5 (26-76)	≤3/3.1-5 (36/41)	0.58	0.29	0.39	1 hemothorax and intrahepatic hematoma, 1 peritoneal hemorrhage
Abdelaziz 2014 ⁷⁹	RCT	Egypt	Early HCC	2.3	RFA	45(52)	31/14	56.8±7.3	2.95±1.03	0.68(1y)	NA	0.68(1y)	2 subcapsular hematoma, 1 thigh burn, 2 pleural effusion
					MWA	66(76)	48/18	53.6±5	2.9±0.97	0.96(1y)	NA	0.96(1y)	1 subcapsular hematoma, 1 abdominal wall skin burn
Shi 2014 ⁸⁰	Retrospect ive cohort	China	HCC	3.8	RES	107(126)	87/20	54.5±9.9	≤3/3.1-5 (37/54)	0.73	0.57	0.60	NA

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					MWA	117(143)	93/24	56.6±9.2	≤3/3.1-5 (40/56)	0.65	0.52	0.52	NA
Yang 2014 ⁸¹	Retrospect ive cohort	Korea	HCC	0.1-7	RES	52	38/14	55.7±10.6	≤2/2.1-5 (21/31)	0.94	NA	0.94	2 pneumonia, 1 wound infection, 1 biliary anastomotic leak, 1 portal vein thrombosis, 1 nausea, 1 delirium, 4 ascites
					RFA	79	59/20	57.2±9.2	≤2/2.1-5 (36/43)	0.86	NA	0.86	1 vomiting, 1 ascites, 6 abdominal pain, 2 nausea, 1 sinus bradycardia
Zhang 2014 ⁸²	Retrospect ive cohort	China	Recurr ent HCC	2.7	RES	27(29)	25/2	47±13	3.2±1.0	NA	NA	0.63	NA
					MWA	39(46)	37/2	52±13	2.7±1.1	NA	NA	0.62	NA
Pompili 2015 ⁸³	Retrospect ive cohort	Italy	Early HCC	2.8	RFA	136	75/61	68 (41-85)	1.8 (1-2)	0.63	NA	0.63	2 ascites, 1 pleural effusion, 1 hemobilia
					PEI	108	90/18	68.5 (34-86)	1.95 (0.8-2)	0.65	NA	0.65	1 hemobilia, 1 portal vein thrombosis
Xu 2015 ⁸⁴	RCT	China	HCC	0.1-3	Laparosco pic RES	45	34/11	58.3±3.1 (26-78)	3.6±0.7 (1-5)	NA	0.38(3y)	0.38(3y)	3 bile leakage, 3 pleural effusion, 2 postoperative hemorrhage
					MWA	45	32/13	57.9±3.4 (27-76)	3.8±0.9 (2-5)	NA	0.33(3y)	0.33(3y)	1 bile leakage, 1 pleural effusion, 1 postoperative hemorrhage

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Agcaoglu O 2013 ⁹⁶	Prospectiv e cohort	Ameri ca	HCC	1.7	RES	94	50/44	61.7±1.2	3.7±0.2	NA	0.53	0.53	2 pulmonary,2 biliary,2 wound-related,1 intestinal,1 hemorrhagic,2 cardiac , and 1 renal
					RFA	295	196/99	63.4 ±0.7	3.4±0.1	NA	0.2	0.2	3 bleeding,2 liver abscess,5 pulmonary,3 renal
Zhou Z 2014 ⁹³	Retrospect ive cohort	China	HCC	5	RES	21	15/6	42.2±7.6	1.7±0.3	0.81	NA	0.81	1 intraperitoneal hemorrhage
					RFA	31	20/11	46.7±9.8	1.7±0.4	0.81	NA	0.81	2 pleural effusion;2 fever;1 pneumonia;1 biloma
Kim JM 2014 ⁹⁵	Retrospect ive cohort	Korea	HCC	2.8	RES	66	48/18	58.	2.1(0.8-3.0)	0.89	NA	0.89	NA
					RFA	67	52/15	59	1.8 (1.0-2.9)	0.49	NA	0.49	NA
Ko S 2014 ⁹⁴	Retrospect ive cohort	China	HCC	5	RES	12	9/3	71.6±4.3	2.9±1.4	NA	NA	0.67	NA
					RFA	17	9/8	57.3±3.6	2.3±1.1	NA	NA	0.35	NA
Kang TW 2015 ⁹²	Retrospect ive cohort	Korea	HCC	5	RES	142	107/35	53(28-74)	2(1.1–3.0)	0.90	NA	0.90	1 intra-abdominal abscess,3 wound problem,1 abdominal bleeding,1 intestinal obstruction

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
					RFA	438	337/101	58(30-80)	1.9(1.1–3.0)	0.85	NA	0.85	3 tumor seeding,2 biloma,2 hepatic abscess,1 bile duct stricture,1 hepatic infarction
Lee YH 2015 ⁹¹	Retrospect ive cohort	China	HCC	3.63	RES	330	261/69	61±12	<5	NA	NA	0.76	NA
					RFA	369	244/125	66±11	<5	NA	NA	0.66	NA
Liu PH 2016 ⁸⁷	Prospectiv e cohort	China	HCC	3.7	RES	109	78/31	60±13	<2	NA	0.81	0.81	NA
					RFA	128	84/44	64±12	<2	NA	0.76	0.76	NA
Hof J 2016 ⁸⁹	Retrospect ive cohort	Nethe rlands	HCC	3.2	RES	261	151/110	63.4	<5	0.69	NA	0.69	NA
					RFA	75	55/20	65.7	<5	NA	0.33(3y)	0.33(3y)	NA
Lee HW 2018 ⁸⁵	RCT	Korea	HCC	5	RES	29	23/6	55.6±7.9	<5	NA	0.97(3y)	0.97(3y)	7 pleural effusion
					RFA	34	24/10	56.1±7.4	<5	NA	0.97(3y)	0.97(3y)	3 pain

Study Year	Design style	Countr /	Disease type	Follow-up (year)	Treatment style	Group n (Tumor n)	Male/ Female	Age	Tumor size, cm	5-year Survival rates (unless stated)			Complication
										<3cm	3-5cm	All	
Li W 2017 ⁸⁶	Retrospect ive cohort	China	HCC	5	RES	220(239)	37/183	61.8 (40-73)	2.1 ±0.5	0.75	NA	0.75	64 complications
					MWA	60(61)	14/46	65(45-71)	2.0 ±0.5	0.67	NA	0.67	13 complications
Vogl TJ 2015 ⁹⁰	Retrospect ive cohort	Germ any	HCC	5	RFA	25(32)	19/6	57 ±3.5	3.2(0.8-4.5)	0.72(3y)	NA	0.72(3y)	NA
					MWA	28(36)	23/5	60 ±4.2	3.6(0.9-5)	0.79	NA	0.79(3y)	NA
Liu H 2016 ⁸⁸	RCT	China	HCC	4.7	TR	100(114)	86/14	52(31-80)	2.8(0.6-5)	0.67	NA	0.67	8 pleural effusion,5 biliary fstula,4 abdominal ascites,2 liver dysfunction,2 pneumonia,1 wound infection,1 abdominal infection
					RES	100(109)	94/6	49(30-76)	3(0.6-5)	0.84	NA	0.84	4 pleural effusion,3 liver dysfunction,3 abdominal ascites,1 abdominal bleeding

HCC: hepatocellular carcinoma;
BCLC: Barcelona Clinic Liver Cancer;
RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection;
 RCT: randomized controlled trial;
 NA: not available.

Table S2.
Quality assessment of included studies using GRADE framework.

Intervention/Comparator	Illustrative comparative risks* (per 1000, 95% CI)			Relative effect of survival time (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Comparator	Assumed survival risk	Corresponding survival risk with intervention			
1-year OS rate						
RES/MWA	923	984 (932 to 997)		OR 5.25 (1.15 to 23.97)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
RES/MWA	947	944 (902 to 968)		OR 0.94 (0.52 to 1.71)	990 (6 studies)	⊕ ⊕ ⊖ ⊖ low
RES/PEI	835	802 (674 to 889)		OR 0.80 (0.41 to 1.58)	519 (3 studies)	⊕ ⊕ ⊖ ⊖ low
RES/PEI	944	963 (906 to 1000)		OR 1.02 (0.96 to 1.09)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
RES/RFA	932	945 (931 to 956)		OR 1.25 (0.99 to 1.60)	5006 (30 studies)	⊕ ⊕ ⊕ ⊕ high
RES/TR	939	904 (765 to 965)		OR 0.61 (0.21 to 1.79)	201 (2 studies)	⊕ ⊕ ⊖ ⊖ low
RES/TR	938	802 (310 to 978)		OR 0.27 (0.03 to 2.90)	31 (1 study)	⊕ ⊕ ⊖ ⊖ low

3-year OS rate

RES/MWA	712	734 (623 to 822)	OR 1.12 (0.67 to 1.87)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
2					
3					
REA/MWA	736	779 (717 to 828)	OR 1.26 (0.91 to 1.73)	987 (6 studies)	⊕ ⊕ ⊖ ⊖ low
4					
5					
6					
RES/PEI	499	536 (421 to 645)	OR 1.16 (0.73 to 1.83)	519 (3 studies)	⊕ ⊕ ⊖ ⊖ low
7					
8					
9					
10					
RFA/PEI	729	748 (657 to 822)	OR 1.10 (0.71 to 1.71)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
11					
12					
13					
RES/RFA	785	851 (823 to 875)	OR 1.57 (1.28 to 1.93)	15906 (30 studies)	⊕ ⊕ ⊕ ⊖ moderate
14					
15					
16					
RES/TR	798	760 (618 to 860)	OR 0.80 (0.41 to 1.55)	201 (2 studies)	⊕ ⊕ ⊖ ⊖ low
17					
18					
19					
20					
RFA/TR	737	611 (516 to 704)	OR 0.56 (0.38 to 0.85)	454 (4 studies)	⊕ ⊕ ⊕ ⊖ moderate
21					
22					
23					
24					
RES/MWA	545	607 (492 to 712)	OR 1.29 (0.81 to 2.07)	290 (2 studies)	⊕ ⊕ ⊖ ⊖ low
25					
26					
27					
28					
29					
30					
RES/PEI	293	436 (334 to 545)	OR 1.87 (1.21 to 2.90)	519 (3 studies)	⊕ ⊕ ⊕ ⊖ moderate
31					
32					
33					
RFA/PEI	533	496 (368 to 624)	OR 0.86 (0.51 to 1.45)	9187 (4 studies)	⊕ ⊕ ⊖ ⊖ low
34					
35					
36					
RES/RFA	601	744 (705 to 779)	OR 1.93 (1.59 to 2.34)	15154 (25 studies)	⊕ ⊕ ⊕ ⊖ moderate
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					

RES/TR	290	419 (251 to 607)	OR 1.76 (0.82 to 3.78)	117 (1 study)	⊕ ⊕ ⊕ ⊖ low
RFA/TR	464	356 (222 to 523)	OR 0.64 (0.33 to 1.27)	139 (1 study)	⊕ ⊕ ⊕ ⊖ moderate

The absolute and relative risk of survival with treatments*. GRADE: Grading of Recommendations, Assessment, Development and Evaluation. *The results presented in the Table S1 were built around the assumption of a consistent relative effect. The implications of this effect for populations were considered at different baseline risks. Based on the assumed risks, corresponding risks after an intervention were calculated using the meta-analytic risk ratio.

Table S3.
Ranking treatments of 1-, 3-year and 5-year survival rate of the lesions < 3 cm, 3-5 cm and ≤ 5 cm in RCT.

Treatment	1-year			3-year			5-year		
	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank
< 3cm	13			11			5		
RES		2	2.86		1	1.52		1	1.42
RFA		3	3.13		3	2.58		2	2.46
MWA		1	1.04		NA	NA		NA	NA
TR		4	3.59		2	2.35		3	2.89
PEI		5	4.43		4	3.55		4	3.23
3-5cm	4			4			2		
RES		1	1.17		1	1.19		1	1.69
RFA		3	2.88		3	2.91		3	2.60
MWA		NA	NA		NA	NA		NA	NA
TR		2	1.94		2	1.90		2	1.71
PEI		NA	NA		NA	NA		NA	NA
All tumours (≤ 5 cm)	20			16			7		

5cm)							
RES	3	2.53	1	1.85	1	1.62	
RFA	4	3.94	4	3.62	3	2.87	
MWA	1	1.67	3	2.88	NA	NA	
TR	2	1.93	2	2.38	2	1.78	
PEI	5	4.92	5	4.27	4	3.73	

RES: resection;
RFA: radiofrequency ablation;
MWA: microwave ablation;
TR: transcatheter arterial chemoembolization and radiofrequency ablation;
PEI: percutaneous ethanol injection.

Table S4.
Ranking treatments of 1-, 3-year and 5-year survival rate of the lesions < 3 cm, 3-5 cm and ≤ 5 cm in all studies.

Treatment	1-year			3-year			5-year		
	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank	Study numbers (n)	Rank	Meanrank
< 3cm	50			48			37		
RES		3	1.18		1	1.71		1	1.16
RFA		4	3.17		3	3.38		2	3.02
MWA		1	1.91		4	3.42		3	3.11
TR		2	2.63		2	2.73		4	3.61
PEI		5	4.62		5	3.76		5	4.11
3-5cm	19			18			12		
RES		1	1.12		1	1.04		1	1.93
RFA		4	3.54		4	3.58		3	3.18
MWA		3	3.45		3	3.50		4	3.43
TR		2	2.05		2	2.14		2	1.94
PEI		5	4.84		5	4.74		5	4.53

All tumours (≤ 5 cm)	72		68		50		
RES	2	2.07		1	1.50		1.11
RFA	3	3.48		3	3.68		3.34
MWA	4	3.57		4	3.84		3.23
TR	1	1.19		2	1.82		3.05
PEI	5	4.70		5	4.16		4.28

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection.

Table S5.

Survival rates (1-year, 3-year and 5-year) for small lesion (<3cm) treatment comparisons estimated by direct and network meta-analysis in RCT.

Intervention	OR (95% CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.97 (0.42-1.98)	0.98 (0.77-1.26)
MWA vs RES	152 (1.44-505.80)	NA
TR vs RES	1.08 (0.15-3.78)	0.99(0.67-1.47)
PEI vs RES	0.64 (0.18-1.61)	1.03 (0.54-1.94)
MWA vs RFA	173.30 (1.90-537.40)	1.42 (0.63-3.19)
TR vs RFA	1.25 (0.16-4.64)	1.00 (0.56-1.80)
PEI vs RFA	0.67 (0.28-1.35)	0.97 (0.78-1.19)
TR vs MWA	0.15 (0-0.80)	NA
PEI vs MWA	0.08 (0-0.38)	NA
PEI vs TR	1.17 (0.11-4.66)	NA
3-year OS rate for treatment vs reference		

RFA vs RES	0.75 (0.41-1.31)	0.92 (0.71-1.19)
MWA vs RES	NA	NA
TR vs RES	1.17 (0.16-4.17)	0.80(0.52-1.22)
PEI vs RES	0.58 (0.29-1.16)	1.21 (0.59-2.15)
MWA vs RFA	NA	NA
TR vs RFA	1.54 (0.25-13.43)	1.01 (0.55-1.87)
PEI vs RFA	0.79 (0.45-1.39)	0.91 (0.71-1.17)
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	1.02 (0.14-3.56)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.72 (0.10-2.47)	0.93 (0.62-1.37)
MWA vs RES	NA	NA
TR vs RES	0.84 (0.03-4.18)	0.88(0.69-1.12)
PEI vs RES	0.50 (0.04-2.04)	0.55 (0.26-1.15)
MWA vs RFA	NA	NA
TR vs RFA	2.87 (0.04-13.43)	NA
PEI vs RFA	0.94 (0.08-3.97)	0.97 (0.66-1.40)
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	3.93 (0.03-19.61)	NA

Table S6.
Survival rates (1-year, 3-year and 5-year) for lesion (3-5cm) treatment comparisons estimated by direct and network meta-analysis in RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.25 (0-1.47)	0.89 (0.45-1.77)
MWA vs RES	NA	NA

TR vs RES	1.00 (0-5.0)	NA
PEI vs RES	NA	NA
MWA vs RFA	NA	NA
TR vs RFA	3.40 (0.64-11.93)	1.10 (0.78-1.55)
PEI vs RFA	NA	NA
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.24 (0-1.25)	0.70 (0.34-1.45)
MWA vs RES	NA	NA
TR vs RES	1.14 (0-6.20)	NA
PEI vs RES	NA	NA
MWA vs RFA	NA	NA
TR vs RFA	3.98 (0.71-15.22)	1.29 (0.87-1.89)
PEI vs RFA	NA	NA
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA
5-year OS rate for treatment vs reference		
RFA vs RES	1.05 (0.03-5.33)	0.71 (0.32-1.57)
MWA vs RES	NA	NA
TR vs RES	12.87 (0.02-44.43)	NA
PEI vs RES	NA	NA
MWA vs RFA	NA	NA
TR vs RFA	7.64 (0.14-42.49)	1.93 (0.53-7.06)
PEI vs RFA	NA	NA
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	NA	NA

Table S7.
Survival rates (1-year, 3-year and 5-year) for lesion (≤ 5 cm) treatment comparisons estimated by direct and network meta-analysis in RCT.

Intervention	OR (95%CI)	
	Network Meta-analysis	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.57 (0.27-1.08)	0.96 (0.78-1.19)
MWA vs RES	2.01 (0.47-5.70)	0.98 (0.54-1.78)
TR vs RES	1.50 (0.48-3.67)	0.99 (0.67-1.47)
PEI vs RES	0.37 (0.13-0.82)	1.03 (0.54-1.94)
MWA vs RFA	3.84 (0.81-11.60)	1.42 (0.63-3.19)
TR vs RFA	2.69 (1.02-6.04)	1.09 (0.84-1.43)
PEI vs RFA	0.65 (0.33-1.13)	0.95 (0.80-1.14)
TR vs MWA	1.09 (0.16-3.50)	NA
PEI vs MWA	0.27 (0.05-0.84)	NA
PEI vs TR	0.29 (0.09-0.73)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.65 (0.31-1.29)	0.88 (0.71-1.10)
MWA vs RES	1.00 (0.16-3.30)	0.88 (0.39-1.98)
TR vs RES	0.98 (0.35-2.41)	0.80 (0.51-1.22)
PEI vs RES	0.55 (0.19-1.44)	1.12 (0.59-2.15)
MWA vs RFA	1.77 (0.22-6.24)	NA
TR vs RFA	1.56 (0.66-3.25)	1.20 (0.90-1.60)
PEI vs RFA	0.86 (0.39-1.79)	0.84 (0.66-1.07)
TR vs MWA	1.86 (0.21-7.59)	NA
PEI vs MWA	1.05 (0.12-4.56)	NA
PEI vs TR	0.64 (0.19-1.67)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.66 (0.20-1.62)	0.88 (0.65-1.18)
MWA vs RES	NA	NA

TR vs RES	1.35 (0.23-4.69)	0.80 (0.52-1.22)
PEI vs RES	0.41 (0.11-1.02)	0.55 (0.26-1.15)
MWA vs RFA	NA	NA
TR vs RFA	2.29 (0.41-7.61)	1.30 (0.70-2.41)
PEI vs RFA	0.74 (0.16-2.00)	0.97 (0.66-1.40)
TR vs MWA	NA	NA
PEI vs MWA	NA	NA
PEI vs TR	0.53 (0.06-1.90)	NA

OR: odds ratio;

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection;

NA: not available.

Table S8.

Survival rates (1-year, 3-year and 5-year) for small lesion (<3cm) treatment comparisons estimated by direct and network meta-analysis in all studies.

Intervention	OR (95% CI)	
	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.94 (0.39-1.91)	1.00(0.95-1.04)
MWA vs RES	1.49 (0.44-3.85)	1.02(0.72-1.43)
TR vs RES	1.30 (0.28-3.88)	1.01(0.74-1.39)
PEI vs RES	0.63 (0.22-1.44)	1.00 (0.93-1.07)
MWA vs RFA	1.59 (0.69-3.17)	1.02 (0.85-1.23)
TR vs RFA	1.48 (0.34-4.23)	1.00(0.56-1.80)
PEI vs RFA	0.68 (0.38-1.09)	0.99 (0.93-1.06)

TR vs MWA	1.08 (0.21-7.87)	NA
PEI vs MWA	0.49 (0.18-1.10)	NA
PEI vs TR	0.69 (0.14-2.13)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.72 (0.37-1.30)	0.94 (0.90-0.99)
MWA vs RES	0.73 (0.30-1.55)	0.95 (0.78-1.18)
TR vs RES	0.90 (0.31-2.10)	1.08 (0.64-1.33)
PEI vs RES	0.68 (0.30-1.39)	1.00 (0.71-1.40)
MWA vs RFA	1.02 (0.57-1.70)	1.00 (0.82-1.22)
TR vs RFA	1.31 (0.47-2.92)	1.01 (0.55-1.87)
PEI vs RFA	0.96 (0.59-1.50)	0.97 (0.90-1.03)
TR vs MWA	1.38 (0.42-3.40)	NA
PEI vs MWA	1.01 (0.47-1.95)	NA
PEI vs TR	0.90 (0.29-2.17)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.54 (0.24-1.05)	0.85 (0.81-0.90)
MWA vs RES	0.55 (0.19-1.25)	0.88 (0.61-1.30)
TR vs RES	0.49 (0.16-0.18)	0.77 (0.53-1.11)
PEI vs RES	0.43 (0.17-0.89)	0.79 (0.73-0.85)
MWA vs RFA	1.04 (0.50-1.77)	1.02 (0.78-1.33)
TR vs RFA	0.99 (0.32-2.39)	NA
PEI vs RFA	0.82 (0.48-1.29)	0.92 (0.85-0.99)
TR vs MWA	1.03 (0.28-2.73)	NA
PEI vs MWA	0.86 (0.39-1.65)	NA
PEI vs TR	1.07 (0.31-2.72)	NA

Table S9.
Survival rates (1-year, 3-year and 5-year) for lesion (3-5cm) treatment comparisons estimated by direct and network meta-analysis in all studies.

Intervention	OR (95%CI)
35	

	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.12 (0-0.63)	0.96 (0.81-1.14)
MWA vs RES	0.15 (0-1.00)	NA
TR vs RES	0.36 (0.01-2.08)	1.02 (0.55-1.88)
PEI vs RES	0.06 (0-0.31)	NA
MWA vs RFA	1.29 (0.32-3.60)	0.99 (0.60-1.64)
TR vs RFA	2.99 (1.14-6.58)	1.11 (0.80-1.54)
PEI vs RFA	0.49 (0.18-1.12)	0.89 (0.66-1.20)
TR vs MWA	3.39 (0.58-10.44)	NA
PEI vs MWA	0.55 (0.09-1.76)	NA
PEI vs TR	0.20 (0.05-0.54)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.11 (0.01-0.40)	0.72 (0.60-0.88)
MWA vs RES	0.12 (0.01-0.53)	1.02 (0.57-1.81)
TR vs RES	0.26 (0.01-1.10)	0.92 (0.48-1.75)
PEI vs RES	0.06 (0-0.28)	NA
MWA vs RFA	1.15 (0.39-2.65)	0.81 (0.45-1.43)
TR vs RFA	2.38 (0.93-5.38)	1.29 (0.87-1.89)
PEI vs RFA	0.55 (0.12-1.69)	0.71 (0.50-1.00)
TR vs MWA	2.62 (0.61-7.90)	NA
PEI vs MWA	0.61 (0.08-2.26)	NA
PEI vs TR	0.28 (0.04-0.96)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.69 (0.04-3.16)	0.53 (0.40-0.68)
MWA vs RES	1.24 (0.02-4.46)	0.90 (0.48-1.69)
TR vs RES	14.31 (0.04-21.06)	NA
PEI vs RES	3.02 (0.01-2.40)	NA
MWA vs RFA	1.26 (0.19-4.04)	0.57 (0.21-1.51)
TR vs RFA	6.16 (0.27-25.58)	2.36 (0.66-8.37)
PEI vs RFA	0.86 (0.06-2.68)	0.56 (0.37-0.84)

TR vs MWA	11.97 (0.19-46.76)	NA
PEI vs MWA	4.15 (0.04-5.18)	NA
PEI vs TR	5.77 (0.01-2.84)	NA

Table S10.
Survival rates (1-year, 3-year and 5-year) for lesion (≤ 5 cm) treatment comparisons estimated by direct, indirect and network meta-analysis in all studies.

Intervention	OR (95%CI)	
	Network Meta-regression	Pairwise Meta-analysis
1-year OS rate for treatment vs reference		
RFA vs RES	0.68 (0.35-1.17)	0.99 (0.95-1.04)
MWA vs RES	0.70 (0.29-1.39)	0.97 (0.77-1.23)
TR vs RES	1.72 (0.66-3.70)	1.01 (0.76-1.33)
PEI vs RES	0.52 (0.24-0.96)	1.01 (0.74-1.39)
MWA vs RFA	1.04 (0.55-1.76)	1.01 (0.85-1.20)
TR vs RFA	2.55 (1.20-4.85)	1.10 (0.85-1.43)
PEI vs RFA	0.77 (0.51-1.10)	0.98 (0.93-1.05)
TR vs MWA	2.69 (0.99-6.00)	0.91 (0.70-1.18)
PEI vs MWA	0.81 (0.38-1.51)	NA
PEI vs TR	0.34 (0.11-0.63)	NA
3-year OS rate for treatment vs reference		
RFA vs RES	0.63 (0.37-1.01)	0.96 (0.94-0.98)
MWA vs RES	0.62 (0.32-1.09)	0.94 (0.72-1.22)
TR vs RES	0.97 (0.48-1.79)	0.92(0.68-1.24)
PEI vs RES	0.59 (0.30-1.04)	0.93 (0.86-1.00)
MWA vs RFA	0.99 (0.64-1.47)	1.05 (0.86-1.26)
TR vs RFA	1.57 (0.89-2.57)	1.20 (0.90-1.60)
PEI vs RFA	0.94 (0.64-1.34)	0.95 (0.89-1.01)
TR vs MWA	1.65 (0.80-3.03)	NA
PEI vs MWA	0.98 (0.55-1.65)	NA

PEI vs TR	0.64 (0.32-1.16)	NA
5-year OS rate for treatment vs reference		
RFA vs RES	0.52 (0.29-0.88)	0.84 (0.80-0.88)
MWA vs RES	0.55 (0.25-1.05)	0.93(0.78-1.12)
TR vs RES	0.59 (0.25-1.20)	0.69 (0.34-1.42)
PEI vs RES	0.45 (0.23-0.82)	0.79 (0.73-0.85)
MWA vs RFA	1.06 (0.64-1.61)	0.97 (0.75-1.25)
TR vs RFA	1.16 (0.54-2.21)	1.30 (0.70-2.41)
PEI vs RFA	0.87 (0.57-1.26)	0.91 (0.84-0.98)
TR vs MWA	1.16(0.46-2.46)	NA
PEI vs MWA	0.87 (0.46-1.51)	NA
PEI vs TR	0.84 (0.35-1.74)	NA

OR: odds ratio;

RES: resection;

RFA: radiofrequency ablation;

MWA: microwave ablation;

TR: transcatheter arterial chemoembolization and radiofrequency ablation;

PEI: percutaneous ethanol injection;

NA: not available.

Table S11.

Posterior summaries from random effects consistency and inconsistency models for small lesion (<3cm) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.55	0.21	(0.15-1.00)	0.38	0.23	(0.02-0.88)
τ	12.40	65.04	(1.10-45.68)	109.40	620.40	(1.30-940.00)
resdev	90.04	13.04	(66.16-117.10)	94.65	12.94	(70.06-120.70)

pD	66.48			57.5		
DIC	453.18			404.59		
3-year OS rate for treatment vs reference						
σ	0.59	0.14	(0.34-0.88)	0.6	0.14	(0.36-0.91)
τ	3.26	1.62	(1.34-7.33)	3.28	1.90	(1.19-8.10)
resdev	92.02	14.19	(66.64-122.10)	90.7	13.92	(65.64-120.00)
pD	80.45			71.83		
DIC	589.01			517.44		
5-year OS rate for treatment vs reference						
σ	0.53	0.12	(0.32-0.80)	0.55	0.13	(0.34-0.84)
τ	4.06	2.02	(1.66-8.76)	3.80	2.05	(1.40-8.77)
resdev	63.99	11.47	(43.52-88.24)	63.55	11.37	(43.39-87.90)
pD	64.22			55.07		
DIC	488.23			412.10		

Table S12.
Posterior summaries from random effects consistency and inconsistency models for lesion (3-5cm) treatment in all studies.

Parameters	Network meta-regression (consistency model)			Inconsistency Model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.28	0.25	(0.01-0.92)	0.38	0.34	(0.02-1.28)
τ	3108.00	68630.00	(1.44-4879.00)	19500.00	720600.00	(0.62-4178.00)
resdev	28.90	6.96	(17.25-44.41)	484.70	5117	(0.63-2616)
pD	24.70			24.62		
DIC	166.90			157.30		
3-year OS rate for treatment vs reference						
σ	0.62	0.27	(0.17-1.24)	0.67	0.31	(0.14-1.40)
τ	5.34	12.61	(0.83-21.20)	41.87	585.80	(0.52-77.13)
resdev	32.36	8.17	(18.39-50.07)	32.62	8.22	(18.52-50.51)

pD	30.91			28.63		
DIC	212.30			188.69		
5-year OS rate for treatment vs reference						
σ	0.80	0.46	(0.14-1.94)	0.60	0.42	(0.04-1.64)
τ	337.00	11980	(0.30-20.22)	10100.00	258400.00	(0.37-691.30)
resdev	22.54	6.73	(11.29-37.43)	22.57	6.519	(11.45-36.90)
pD	22.61			19.88		
DIC	146.84			131.53		

Table S13.**Posterior summaries from random effects consistency and inconsistency models for lesion (≤ 5 cm) treatment in all studies.**

Parameters	Network meta-regression (consistency model)			Inconsistency Model		
	Mean	sd	CI	Mean	sd	CI
1-year OS rate for treatment vs reference						
σ	0.49	0.13	(0.26-0.77)	0.29	0.14	(0.05-0.58)
τ	6.00	6.24	(1.92-16.85)	116.80	1122.00	(2.96-419.40)
resdev	129.2	14.99	(101.40-160)	133.1	14.50	(105.70-162.80)
pD	95.71			78.20		
DIC	692.39			604.18		
3-year OS rate for treatment vs reference						
σ	0.50	0.09	(0.33-0.70)	0.47	0.096	(0.29-0.67)
τ	4.20	1.45	(2.15-7.71)	5.31	2.59	(2.24-11.80)
resdev	124	15.64	(95.16-156.40)	124.5	15.89	(95.35-157.50)
pD	111.54			93.41		
DIC	856.01			723.74		
5-year OS rate for treatment vs reference						
σ	0.44	0.10	(0.26-0.65)	0.44	0.1	(0.26-0.67)
τ	5.30	2.27	(2.38-14.90)	6.09	3.95	(2.29-14.87)

resdev	86.73	13.53	(62.35-115.40)	85.74	13.55	(61.39-114.40)
pD	84.53			68.81		
DIC	670.73			544.40		

sd: standard deviation;
CI: Credible Interval
 σ : between-trial standard deviation
 τ^2 : between-trial variance
resdev: residual deviance
pD: effective number of parameters
DIC: deviance information criterion

Figure S1.

Results of the consistency test for closed loop at 1-year, 3-year, and 5-year survival rate of the lesions < 3 cm, 3-5 cm and \leq 5 cm.

i Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions < 3 cm

- 1 ii Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions 3-5 cm
2
3 iii Results of the consistency test for closed loop at 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions ≤ 5 cm
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46





Figure S2.**Assessment of publication bias using funnel plot.**

- i Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions < 3 cm.
- ii Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions 3-5 cm.
- iii Assessment of publication bias using funnel plot for 1-year (A), 3-year (B), and 5-year (C) survival rate of the lesions ≤ 5 cm

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46



For peer review only

PRISMA NMA Checklist of Items to Include When Reporting A Systematic Review Involving a Network Meta-analysis

Section/Topic	Item #	Checklist Item	Reported on Page #
TITLE			
Title	1	Identify the report as a systematic review <i>incorporating a network meta-analysis (or related form of meta-analysis)</i> .	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: Background: main objectives Methods: data sources; study eligibility criteria, participants, and interventions; study appraisal; and <i>synthesis methods, such as network meta-analysis</i> . Results: number of studies and participants identified; summary estimates with corresponding confidence/credible intervals; <i>treatment rankings may also be discussed. Authors may choose to summarize pairwise comparisons against a chosen treatment included in their analyses for brevity.</i> Discussion/Conclusions: limitations; conclusions and implications of findings. Other: primary source of funding; systematic review registration number with registry name.	5,6
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known, <i>including mention of why a network meta-analysis has been conducted.</i>	7,8
Objectives	4	Provide an explicit statement of questions being addressed, with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	8

METHODS

Protocol and registration	5	Indicate whether a review protocol exists and if and where it can be accessed (e.g., Web address); and, if available, provide registration information, including registration number.	8,9
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale. <i>Clearly describe eligible treatments included in the treatment network, and note whether any have been clustered or merged into the same node (with justification).</i>	9,10
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	9
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	9,10, Figure 1, Additional file 1: Text S1
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	9,10
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	10
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	11
Geometry of the network	S1	Describe methods used to explore the geometry of the treatment network under study and potential biases related to it. This should include how the evidence base has been graphically summarized for presentation, and what characteristics were compiled and used to describe the evidence base to readers.	11
Risk of bias within individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	11,12

Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means). <i>Also describe the use of additional summary measures assessed, such as treatment rankings and surface under the cumulative ranking curve (SUCRA) values, as well as modified approaches used to present summary findings from meta-analyses.</i>	11,12
Planned methods of analysis	14	Describe the methods of handling data and combining results of studies for each network meta-analysis. This should include, but not be limited to: <ul style="list-style-type: none">• <i>Handling of multi-arm trials;</i>• <i>Selection of variance structure;</i>• <i>Selection of prior distributions in Bayesian analyses; and</i>• <i>Assessment of model fit.</i>	11,12
Assessment of Inconsistency	S2	Describe the statistical methods used to evaluate the agreement of direct and indirect evidence in the treatment network(s) studied. Describe efforts taken to address its presence when found.	10,11,12
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	10,11,12
Additional analyses	16	Describe methods of additional analyses if done, indicating which were pre-specified. This may include, but not be limited to, the following: <ul style="list-style-type: none">• Sensitivity or subgroup analyses;• Meta-regression analyses;• <i>Alternative formulations of the treatment network; and</i>• <i>Use of alternative prior distributions for Bayesian analyses (if applicable).</i>	11,12

RESULTS†

Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	11,12
Presentation of network structure	S3	Provide a network graph of the included studies to enable visualization of the geometry of the treatment network.	12,13,Figure2-3
Summary of network geometry	S4	Provide a brief overview of characteristics of the treatment network. This may include commentary on the abundance of trials and randomized patients for the different interventions and pairwise comparisons in the network, gaps of evidence in the treatment network, and potential biases reflected by the network structure.	12,13,
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	11,12, Table1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment.	
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: 1) simple summary data for each intervention group, and 2) effect estimates and confidence intervals. <i>Modified approaches may be needed to deal with information from larger networks.</i>	12,13, Figure2-5
Synthesis of results	21	Present results of each meta-analysis done, including confidence/credible intervals. <i>In larger networks, authors may focus on comparisons versus a particular comparator (e.g. placebo or standard care), with full findings presented in an appendix. League tables and forest plots may be considered to summarize pairwise comparisons.</i> If additional summary measures were explored (such as treatment rankings), these should also be presented.	12,13,Figure4-5, Additional file 1: Table S1-S13
Exploration for inconsistency	S5	Describe results from investigations of inconsistency. This may include such information as measures of model fit to compare consistency and inconsistency models, <i>P</i> values from statistical tests, or summary of inconsistency estimates from different parts of the treatment network.	12,13

Risk of bias across studies	22	Present results of any assessment of risk of bias across studies for the evidence base being studied.	12,13, Additional file 1: Figure S1-S2
Results of additional analyses	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression analyses, <i>alternative network geometries studied, alternative choice of prior distributions for Bayesian analyses</i> , and so forth).	12,13
DISCUSSION			
Summary of evidence	24	Summarize the main findings, including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy-makers).	14-16
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review level (e.g., incomplete retrieval of identified research, reporting bias). <i>Comment on the validity of the assumptions, such as transitivity and consistency. Comment on any concerns regarding network geometry (e.g., avoidance of certain comparisons).</i>	16
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	17
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. This should also include information regarding whether funding has been received from manufacturers of treatments in the network and/or whether some of the authors are content experts with professional conflicts of interest that could affect use of treatments in the network.	17

PICOS = population, intervention, comparators, outcomes, study design.

* Text in italics indicateS wording specific to reporting of network meta-analyses that has been added to guidance from the PRISMA statement.

† Authors may wish to plan for use of appendices to present all relevant information in full detail for items in this section.

For peer review only